

**INFECTION PREVENTION & CONTROL
PRIMARY CARE GENERAL PRACTICE
POLICIES AND SAFE PRACTICE
GUIDANCE**

Version:	1.1
Approved By:	Quality & Patient Safety Committee
Date Approved:	20 February 2020
Name of originator / author:	Christine Fisher
Name of responsible committee/ individual:	Quality & Patient Safety Committee
Name of executive lead:	Jayne Sivakumar

Date issued:	17/02/2020
Review Date:	17/02/2022
Target Audience:	Primary Care General Practices

INFECTION PREVENTION & CONTROL PRIMARY CARE GENERAL PRACTICE POLICIES AND SAFE PRACTICE GUIDANCE

DOCUMENT CONTROL

Version No	Type of Change	Date	Description of change
V1	New Policy	December 2015	
V1.1	3 year review (overdue)	February 2020	<p>Comprehensive rewrite to clarify purpose and scope of the policy and take into account the following:</p> <ul style="list-style-type: none"> • General Data Protection Regulations 2016 and: • The CCG's Policy on Policies issues in September 2017

CONTENTS		PAGE
1. Introduction		4 - 5
2. Purpose		5 - 6
3. The risks of not having this policy in place		6
4. Definitions		6
5. Principles		6
6. Roles and Responsibilities		6
7. Procedures		6 - 8
8. Monitoring the compliance and effectiveness of this policy		8 - 9
9. References		9
10. Review of the Policy		9
11. Appendices		
Policies		
Appendix 1	Definitions of terminology	
Appendix 2	Notifiable diseases & Notification of cluster infections	
Appendix 3	Recognition and management of an outbreak of infection	
Appendix 4	Residential Care Home	
Appendix 5	Antibiotic Prescribing Policy	
Safe practice guidance – general:		
Appendix 6	Infection Control Principles – the spread of infection	
Appendix 7	Standard Infection Control Precautions	
Appendix 8	Hand Hygiene	
Appendix 9	Personal Protective Equipment (PPE)	
Appendix 10	Safe use and disposal of Sharps	
Appendix 11	Management of healthcare waste	
Appendix 12	Environmental cleaning	
Appendix 13	Spillages of blood and body fluids	
Appendix 14	Pest control	
Safe practice guidance – clinical:		
Appendix 15	Decontamination of medical equipment	
Appendix 16	Single use and single patient use medical devices	
Appendix 17	Vaccine management	
Appendix 18	Aseptic technique and care of invasive devices	
Appendix 19	Isolation of infectious patients in general practice	

Appendix 20	Collection of Microbiological Specimens	
Appendix 21	Infections with specific alert organisms	
Appendix 22	Management of <i>Clostridium difficile</i> diarrhoea	
Safe practice guidance – personnel:		
Appendix 23	Management of occupational exposure to blood-borne viruses	
Appendix 24	Management of infections in staff	
Appendix 25	Vaccination programme for staff and service users	
Appendix 26	Definitions of terminology used in policy and guidance	
	Bibliography	

INFECTION PREVENTION & CONTROL PRIMARY CARE GENERAL PRACTICE POLICIES AND SAFE PRACTICE GUIDANCE

1. Introduction

1.1 The Health and Social Care Act 2008 (Regulated Activities) Regulations 2014 requires all providers of a regulated service to be registered with the Care Quality Commission (CQC). A service is regulated if it appears in a list of activities described in legislation. This includes the delivery of primary care to service users in registered primary care facilities including those providing Out of Hours services.

1.2 The Health and Social Care Act 2008 (Regulated Activities) Regulations 2014 comes into force in April 2015 for all providers and includes the following regulations:

'Regulation 12: Safe Care and treatment

- ☐ *(2)(h) assessing the risk of, and preventing, detecting and controlling the spread of infections, including those that are healthcare associated.....'*

And

'Regulation 15: Premises and equipment'

- ☐ *15 (1) All premises and equipment used by the service provider must be clean.....'*

1.3 These new regulations introduce the fundamental standards of quality and safety that all healthcare providers must comply with to prevent service users from receiving unsafe care and treatment and in order to prevent any risk of harm. The CQC will monitor organisations for the characteristics of 'good care'; against these fundamental standards; to ensure services provided do not fall below acceptable levels. The CQC will use key lines of enquiry (KLOEs) to assess these fundamental standards and check whether the services provided are: safe, effective, caring, responsive and well-led.

1.4 In relation to implementation of these regulations the provider should be able to demonstrate compliance with the recently updated 10 Criteria of The Health and Social Care Act 2008 *Code of Practice on the prevention and control of infections and related guidance (DH) 2015*. The Code contains statutory guidance about compliance with the registration requirement relating to infection control (regulation 12 (2) (h) of the Health and Social Care Act 2008 (Regulated Activities) Regulations 2014.

- 1.5 The law states that the Code must be taken into account by the CQC when it makes decisions about registration against the infection prevention requirement 12(h). Likewise providers must have regard for the Code when deciding how they will comply with registration requirements. By following the Code, registered providers will be able to show that they meet the regulation on cleanliness and infection prevention. However, they do not by law have to comply with the Code. They may be able to demonstrate a different way (equivalent or better) of compliance from that described in the Code.
- 1.6 The revised and updated Code of Practice does not make any new requirements of providers of services. However, it provides more explicit guidance on some elements of infection prevention and cleanliness with regard to recent changes in the evidence base as well as keeping pace with recent organizational changes and priorities. In particular, there is an additional requirement in relation to antimicrobial prescribing which is discussed in section 4 of this manual.
- 1.7 This section of the infection prevention and control (IPC) manual describes how IPC is managed within the organisation in accordance with the legislation and expert guidance as above.

2. Purpose

- 2.1 The purpose of infection prevention and control is to limit the acquisition and spread of pathogenic micro-organisms, by using scientifically based knowledge and through planning, surveillance, education and research as part of the overall policy of achieving high quality health and social care services.
- 2.2 The organisation supports the principle that infections should be prevented wherever possible and that effective arrangements for the surveillance, prevention and control of infection are provided throughout the organisation.
- 2.3 It is the organisation's policy to encourage the individual responsibility of every member of staff to participate in the prevention and control of infection and to comply with Health and Safety, COSHH and other legislation and regulations applying to the safe provision of health and social care.
- 2.4 This policy and guidance apply to all members of staff employed in the practice and includes agency, locum and bank workers as well as volunteers.

- 2.5 All adjustments to infection prevention and control arrangements and policy must be approved and assessed by the IPC lead.

3. The Risks of not having this Policy in place

- 3.1 Failure to comply with this policy may result in the following risks arising:
- 3.2 The CCG may not achieve its statutory obligations
- 3.3 The CCG may be unable to demonstrate that safe and effective Infection Prevention and Control Standards are in place wherever the CCG commissions
- 3.4 Risk of cross infection or contamination due to lack of clarity around measures staff should take to minimise the risk of infection to themselves and their clients

4. Definitions

- 4.1 Appendix 26 is a list and description of the meaning of the terms used in the context of the policy, procedures and guidance.

5. Principles

- 5.1 Infection prevention and control has a key role to play in the clinical governance framework of any health and care organisation. This comprehensive range of guidance documents have been developed to help with CQC requirements and to achieve compliance with the Health and Social Care Act 2008: Code of Practice on the prevention and control of infections and related guidance

6. Roles and Responsibilities

- 6.1 Chief Nurse, Barnsley CCG – Lead Officer
Assistant Director IPC - Operational
Root Cause Analysis Group – reports to:
Post Infection Review Group – BMBC, SWYPT, BHNFT & BCCG reports to:
Quality & Patient Safety Committee, BCCG reports to:
Governing Body, BCCG

7. Procedures

- 7.1 Training

Infection prevention and control training is a mandatory requirement at induction for all staff groups and as part of

mandatory updates for all staff involved in service users' care. Training attendance records must be maintained and reported through internal governance frameworks. All training delivered should be evaluated by delegates.

The Code of Practice does not specify the frequency of update training. However, the recently published UK wide Core Skills Training Framework (www.skillsforhealth.org.uk) specifies a frequency of annual training for mandatory core subjects, which includes infection prevention and control.

7.2 Governance

Infection prevention and control has a key role to play in the clinical governance framework of any health and care organisation.

The following activities should be considered an essential element of local IPC activity:

- Development of annual IPC programme and annual statement
- the implementation and monitoring of policies
- the education of all staff
- surveillance and reporting of occurrences of infection

Practices are required to have a nominated Infection Prevention & Control lead who will outline activities required to be undertaken to provide assurance under the Code of Practice.

The IPC lead may be the registered manager. If someone else takes this lead role they should report directly to the registered manager in this regard and produce an annual statement outlining the IPC arrangements and activities including policy compliance information.

Specialist advice on IPC should be available to all staff. This may be through the infection prevention and control team or other providers of expertise e.g. Public Health England.

7.3 Information Sharing

The Code of Practice requires primary care medical practices to share information on IPC activities and outcomes with patients. Involvement of patient liaison groups is recommended.

Practices are also required to share patient information as appropriate with other health and care providers having due regard to patient confidentiality requirements.

7.4 Uniform and Dress Code

The organisation supports the view that staff clothing should be such that it minimises risks of the transmission of infection. It is a requirement of the Code of Practice that all organisations have a written uniform and dress code policy. Compliance with this policy should form part of the annual audit programme.

In particular clothing must facilitate good hand hygiene practice. Staff must adhere to Bare Below the Elbow (BBE). Stoned rings and wrist jewellery should not be worn when performing clinical tasks. Long sleeves, if worn, should be rolled to the elbow for hand washing and clinical tasks.

7.5 See the list of appendices for all the procedures linked to this policy.

8. Monitoring the Compliance and Effectiveness of this Policy

8.1 The Lead Officer is responsible for ensuring the compliance and effectiveness of the policy.

8.2 Contact monitoring is undertaken quarterly.

8.3 A rolling audit programme is prepared annually by the IPC team with clear timescales for completion and progress and is monitored through governance frameworks.

8.4 The Infection Prevention Society audit tool is used by the community IPC team to audit GP practices and care homes.

8.5 The process for reviewing results and ensuring improvements is undertaken by the CCG and BMBC.

8.6 The monitoring of key performance indicators is undertaken by Contract monitoring.

8.7 Surveillance and data collection is a requirement of the Code of Practice but a specific policy on this is not required in primary care. However, it is recommended that a local system for monitoring infections is implemented. In particular post procedure surgical site surveillance is strongly recommended where minor operative procedures are undertaken.

8.8 Mandatory reporting of infections to Public Health England (PHE) is a requirement for MRSA bacteraemia, *Clostridium difficile* toxin and Gram-negative bacteraemia. Annual trajectories are set by the Department of Health to reduce the number of cases of MRSA bacteraemia and toxin positive cases of *C. difficile* diarrhoea. Trajectories are set for all acute

NHS Trusts and also (since 2011/12) for all Primary Care Organisations and now (since April 2014) for all Clinical Commissioning Groups (CCGs). Breaching trajectories carries financial penalties for the organisations concerned. Providers of primary health care services may be involved in the review of individual cases of these infections as part of the mandatory processes expected to be undertaken. Reviews are the responsibility of the acute Trust or CCG (depending on timing of positive specimen) and GPs / Practice Managers responsible for individual patients' will be informed if their participation is required on a case by case basis.

Monitoring of mandatory reporting of key infections is routinely undertaken by local commissioners of health care.

9. References

9.1 The references in relation to this policy will be reviewed two yearly.

10. Review of the Policy

10.1 The review will be two years from the date of approval of this policy.

10.2 Individual processes and guidelines will be updated as required, in response to new evidence, expert guidance or regulation.

INFECTION PREVENTION & CONTROL PRIMARY CARE GENERAL PRACTICE POLICIES AND SAFE PRACTICE GUIDANCE CONTENTS

Table of Contents

Ctrl + Click on each section heading to go directly to that section

[NOTIFIABLE DISEASES](#)

[RECOGNITION AND MANAGEMENT OF AN OUTBREAK OF INFECTION](#)

[MANAGEMENT OF AN OUTBREAK OF VIRAL GASTRO-INTESTINAL ILLNESS IN RESIDENTIAL CARE HOMES](#)

[ANTIBIOTIC PRESCRIBING POLICY](#)

[INFECTION CONTROL PRINCIPLES](#)

[STANDARD INFECTION CONTROL PRECAUTIONS \(SICPs\)](#)

[HAND HYGIENE](#)

[PERSONAL PROTECTIVE EQUIPMENT \(PPE\)](#)

[SAFE USE AND DISPOSAL OF SHARPS](#)

[MANAGEMENT OF HEALTHCARE WASTE - SUMMARY](#)

[ENVIRONMENTAL CLEANING](#)

[SPILLAGES OF BLOOD AND BODY FLUIDS](#)

[PEST CONTROL](#)

[ESTATES AND FACILITIES MANAGEMENT](#)

[DECONTAMINATION OF MEDICAL EQUIPMENT](#)

[SINGLE USE AND SINGLE PATIENT USE MEDICAL DEVICES](#)

[VACCINE MANAGEMENT GUIDELINES](#)

[ASEPTIC TECHNIQUE](#)

[ASEPSIS IN MINOR SURGICAL PROCEDURES](#)

[ISOLATION OF INFECTIOUS PATIENTS IN GENERAL PRACTICE](#)

[COLLECTION OF MICROBIOLOGICAL SPECIMENS](#)

[INFECTIONS WITH SPECIFIC ALERT ORGANISMS](#)

[MRSA](#)

[MANAGEMENT OF CLOSTRIDIUM DIFFICILE DIARRHOEA](#)

[MANAGEMENT OF OCCUPATIONAL EXPOSURE TO BLOOD-BORNE VIRUSES](#)

[MANAGEMENT OF INFECTIONS IN STAFF](#)

[DEFINITIONS OF TERMINOLGY](#)

[BIBLIOGRAPHY](#)

THIS PAGE HAS BEEN LEFT INTENTIONALLY
BLANK

NOTIFIABLE DISEASES

Some infectious diseases may spread easily in a community or may cause serious diseases in individuals. The requirement to notify some infectious diseases is contained within the Public Health (Control of Disease) Act 1984, updated 2010. Additional guidance and the list of notifiable diseases are contained in the Health Protection Legislation (England) Guidance 2010. This list is reproduced in on the next page to this section.

It is the responsibility of Registered Medical Practitioners (RMPs) to notify any suspected or confirmed instance of notifiable disease to the local authority and / or local health protection teams (local arrangements will vary). Laboratories also have notification responsibilities.

Additionally RMPs are required to notify instances of infection which, in the view of the clinician presents, or could present, significant risk to human health and this should include new or emerging infections.

The regulations concern only single incidents of infection or suspected infection. The guidance however does advise that RMPs continue to voluntarily notify clusters of cases of infection whether the disease in question is notifiable or not. This includes, for example, outbreaks of diarrhoea / vomiting.

Notification by clinicians should be to the 'Proper Officer' within the local authority or health protection team. This may in some cases be deemed urgent (see table below) in which case notification by telephone is required. In all cases notification should also be made on a written form within 3 days. Such written notification may be made by secure email. Notification forms can be acquired from the local authority or Public Health England health (PHE) protection team.

Local arrangements for health protection teams in your region can be found at:
www.gov.uk/government/organisations/public-health-england

NOTIFIABLE DISEASES

Notifiable diseases	Definition / comment	Likely to be Urgent?
Acute encephalitis		No
Acute meningitis	Viral and bacterial.	Yes, if suspected bacterial infection.
Acute poliomyelitis		Yes
Acute infectious hepatitis	Close contact of acute hepatitis A and B cases need rapid prophylaxis. Urgent notification will facilitate prompt laboratory testing. Hepatitis C cases known to be acute need to be followed up rapidly as this may signify recent transmission from a source that could be controlled.	Yes
Anthrax		Yes
Botulism		Yes
Brucellosis		No – unless thought to be UK – acquired
Cholera		Yes
Enteric fever (typhoid or paratyphoid fever)	Clinical diagnosis of a case before microbiological confirmation(e.g. case with fever, constipation, rose spots and travel history) would be an appropriate trigger for initial public health measures, such as exclusion of cases and contacts in high risk groups (e.g. food handlers)	Yes
Food poisoning	Any disease of infectious or toxic nature caused by, or thought to be caused by consumption of food or water (definition of the advisory committee on the Microbiological Safety of Food).	Clusters and outbreaks, yes For specific organisms see Table 2

Notifiable diseases	Definition / comment	Likely to be Urgent?
Mumps	Post-exposure immunization (MMR or HNIG) does not provide protection for contacts.	No
Plague		Yes
Rabies	A person bitten by suspected rabid animal should be reported and managed urgently, but if a patient is diagnosed with symptoms of rabies, they will not pose a risk to human health.	Yes
Rubella	Post-exposure immunization (MMR or HNIG) does not provide protection for contacts.	No
SARS		Yes
Smallpox		Yes
Tetanus		No, unless associated with injection drug use
Tuberculosis		No, unless healthcare worker or suspected cluster or multi drug resistance
Typhus		No
Viral haemorrhagic fever (VHF)		Yes
Whooping cough		Yes, if diagnosed during acute phase
Yellow fever		No, unless thought to be UK- acquired

NOTIFICATION OF CLUSTERS OF INFECTION

Although not Notifiable, clusters of infection (even single cases) can have significant public health implications. Such infections should be reported to the local HPU or proper officer (local authority) promptly. Examples include:

Notifiable diseases	Definition / comment	Likely to be Urgent?
Acute encephalitis		No
Acute meningitis	Viral and bacterial.	Yes, if suspected bacterial infection.
Acute poliomyelitis		Yes
Acute infectious hepatitis	Close contact of acute hepatitis A and B cases need rapid prophylaxis. Urgent notification will facilitate prompt laboratory testing. Hepatitis C cases known to be acute need to be followed up rapidly as this may signify recent transmission from a source that could be controlled.	Yes
Anthrax		Yes
Botulism		Yes
Brucellosis		No – unless thought to be UK – acquired
Cholera		Yes
Enteric fever (typhoid or paratyphoid fever)	Clinical diagnosis of a case before microbiological confirmation(e.g. case with fever, constipation, rose spots and travel history) would be an appropriate trigger for initial public health measures, such as exclusion of cases and contacts in high risk groups (e.g. food handlers)	Yes
Food poisoning	Any disease of infectious or toxic nature caused by, or thought to be caused by consumption of food or water (definition of the advisory committee on the Microbiological Safety of Food).	Clusters and outbreaks, yes For specific organisms see Table 2
Haemolytic uraemic syndrome (HUS)		Yes
Infectious bloody diarrhoea	See also HUS in Schedule 1 and VTEC in schedule 2.	Yes
Invasive group A streptococcal disease and scarlet fever		Yes, if IGAS . No if scarlet fever

Notifiable diseases	Definition / comment	Likely to be Urgent?
Legionnaires' disease		Yes
Leprosy		No
Malaria		No, unless thought to be UK- acquired
Measles		Yes
Meningococcal septicaemia		Yes
Mumps	Post-exposure immunization (MMR of HNIG) does not provide protection for contacts.	No
Plague		Yes
Rabies	A person bitten by suspected rabid animal should be reported and managed urgently, but if a patient is diagnosed with symptoms of rabies, they will not pose a risk to human health.	Yes
Rubella	Post-exposure immunisation (MMR or HNIG) does not provide protection for contacts.	No
SARS		Yes
Smallpox		Yes
Tetanus		No, unless associated with injection drug use
Tuberculosis		No, unless healthcare worker or suspected cluster or multi drug resistance
Typhus		No
Viral haemorrhagic fever (VHF)		Yes
Whooping cough		Yes, if diagnosed during acute phase

THIS PAGE HAS BEEN LEFT
INTENTIONALLY BLANK

RECOGNITION AND MANAGEMENT OF AN OUTBREAK OF INFECTION

INTRODUCTION

An outbreak of communicable disease/infection can be defined as the incidence of disease above that normally expected. Usually this means that there are two or more linked cases with the same illness/symptoms. In some instances, only one case may be sufficient to instigate investigation as an Incident, e.g. meningococcal meningitis. Outbreaks in community care settings where primary care medical practices may be involved will be similar to those experienced in acute hospital settings e.g. viral gastro-enteritis, influenza etc.

Outbreaks of infection may vary in extent and severity, ranging from a few cases of infestation to a large number of food poisoning cases, affecting hundreds of people. Recognition of an outbreak in the early stages may be difficult, therefore medical and nursing/care staff must remain vigilant.

The Consultant in Communicable Disease Control (CCDC) at the local PHE department has overall responsibility for outbreaks of infection in all health and social care provider settings (both NHS and independent sector) and the designated infection control advisor / senior manager on call in the provider setting must inform the local PHE of any suspected outbreak of infection. An on-call service is provided by the PHE, out-of-hours and at weekends.

STAFF RESPONSIBILITIES

Primary Care Medical practice staff should be able to recognise a potential outbreak of infection or food poisoning.

Outbreaks in community residential settings should be managed by the organisation providing the care with advice and support from the Infection Control and Prevention Team (IPCT) and/or PHE. It is recommended that Primary Care Medical practice staff responsible for an individual's care within that setting assure themselves that outbreaks are reported to IPCT and that processes to manage the outbreak are in place.

Practice staff should be aware of contact telephone numbers for IPCT and PHE and of local arrangements for managing outbreaks in community settings.

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

MANAGEMENT OF AN OUTBREAK OF VIRAL GASTRO-INTESTINAL ILLNESS IN RESIDENTIAL CARE HOMES

This guidance is provided for information to GPs to assist them when providing support and guidance to care home managers and staff.

The management of outbreaks of viral gastro-enteritis in primary care settings can be a significant challenge to staff. General Practitioners play an important role in supporting managers and staff during such outbreaks whilst also maintaining responsibility for those residents that are registered with their practice. In addition, the role of the IPCT for Community and Public Health England (PHE) in advising in the early detection and management of infective diarrhoea/Norovirus outbreaks is crucial to the success of local control measures.

Outbreaks of infective diarrhoea/norovirus infection in care homes and other healthcare environments can have a devastating effect on activity due to the numbers of affected individuals which includes staff. Business continuity can be adversely affected and acute NHS Trusts have been known to close due to the effect of widespread outbreaks. Often entire communities are affected with schools, nurseries, hospitals, care homes etc. being the most affected due to the large numbers of susceptible individuals in confined environments.

Prompt identification of possible cases of infective diarrhoea/ norovirus is crucial so that early interventions aimed at limiting spread can be implemented.

Norovirus is responsible for causing a high incidence of viral gastro-intestinal illness. This was previously called “winter vomiting”, Norwalk virus and small round structured virus (SRSV). Norovirus is a highly contagious gastro-intestinal virus that can be spread by a number of different routes:

- By direct contact with an affected individual.
- By aerosol droplets of virus particles in faces; and in particular when vomiting occurs.
- By aerosol droplets (from vomit / faeces) landing onto surfaces and equipment and then being transferred onto hands and then into the mouth.
- Food-borne either from contaminated food or water or by food handlers that are symptomatic;

Norovirus causes a short illness (24 – 48 hours) associated with nausea, profuse vomiting – often projectile, diarrhoea and abdominal pain. Infection is self-limiting but can cause dehydration and deterioration in the very young and elderly.

.CRITERIA FOR SUSPECTING NOROVIRUS OUTBREAK

- ☐ Vomiting in > 50% of cases (although sometimes diarrhoea is the prominent symptom)
- Duration of illness 24 - 48hrs hours
- ☐ Patients AND staff affected (this is a critical criteria)
- Cases often occur in clusters up to 48 hours apart due to incubation period of 24–48 hours.

REPORTING / RECORDING AN OUTBREAK OF VIRAL GASTRO-ENTERITIS

As soon as an outbreak is suspected, it is essential to report cases through the local incident reporting mechanism. The IPCT should be contacted to log the details of the outbreak and provide further guidance. In addition, clinicians including GPs must be made aware. It is essential that the local PHE is notified of the existence of an outbreak, irrespective of whether this is deemed to be trivial. PHE have a responsibility to be notified of all outbreaks of infection in the independent sector and to be actively involved in outbreaks occurring in the NHS. PHE provide 24hr cover over the weekend and Bank holidays if urgent advice is required.

Outbreaks of viral gastro-enteritis in care homes can have a significant knock-on impact of local acute NHS healthcare facilities if service users require admission to hospital for further care. It is *essential* that the IPCT are involved at the earliest opportunity so that they can communicate promptly and effectively with other healthcare providers (and ambulance personnel) to minimise the risk of spread and service disruption.

DOCUMENTATION

It is advised that staff accurately complete a DAILY outbreak record sheet to assist in managing the outbreak and for documentation purposes. It is essential to include sick staff details on the record sheets as well as service users. An outbreak toolkit can be found at the end of this section and can be modified and photocopied for local use. A Bristol stool chart is also included.

INVESTIGATION OF SPECIMENS.

Faecal specimens should be taken from symptomatic service users as soon as possible after symptoms develop. Ideally, staff should also submit faecal samples via their own GP. Only a small sample is required – do not fill container to top – and it is acceptable to obtain a specimen from a bedpan that also contains urine, as this will not affect results. Request forms should be sent for both C&S **and** virology, including Norovirus and forms should be marked “outbreak”. Remember to send specimens **PROMPTLY** for investigation as virus particles deteriorate rapidly leading to difficulty in detection. Unless specifically requested, do NOT send samples of vomit for investigation as these are not required.

OUTBREAK MANAGEMENT

The most important aspects of outbreak control are (a) outbreak recognition and reporting and (b) implementation of strict enteric precautions to minimise spread.

ENTERIC PRECAUTIONS

The three most important actions during an outbreak of diarrhoea and vomiting are:

- Effective hand hygiene

- Isolation of affected patients, restriction of movement of staff, service users and visitors and exclusion of affected staff
- Enhanced cleaning of the environment and equipment

EFFECTIVE HAND HYGIENE

Effective hand hygiene is vital to prevent transmission of infection and must be actively encouraged. Managers must ensure that staff are properly trained in hand washing technique and that they have easy access to hand hygiene facilities including warm water, liquid soap and disposable paper towels. Plain liquid soap is adequate; antiseptic agents e.g. 'Hibiscrub' are not required for routine hand hygiene even during an outbreak.

- Hand should always be cleaned:
- Before patient contact.
- Before placing apron and gloves on
- When in contact with blood or body fluids.
- Before and after performing an aseptic technique
- After patient contact, or contact with the service users environment.
- Before preparing food.
- Before serving food.

Remember to always provide service users with adequate hand-washing facilities i.e. washing hands at a sink with liquid soap, or an antibacterial/ detergent wipe if service user is not able to use a sink.

Please note:

Alcohol-based products e.g. alcohol gel should **NOT** be used as a primary means of hand decontamination as this has been found to be less effective in viral outbreaks of gastro-enteritis. Liquid soap and water should always be used initially and can be supplemented by the use of alcohol if required afterwards.

MANAGEMENT OF RESIDENTS (ISOLATION)

It is necessary to isolate residents with symptoms of diarrhoea and/ or vomiting immediately they become symptomatic. This means they have to remain in their own room, i.e. away from others who are well (asymptomatic), and with their own toilet facilities and designated cleaning equipment. If en-suite facilities are not available, specific toilet areas should be designated for their use only, or a commode allocated for individual symptomatic residents only.

All en-suite toilets / commodes must be cleaned after use to reduce the risk of environmental contamination. It is very important that strict isolation procedures are implemented by staff e.g. hand washing, environmental cleaning, and safe handling of infected linen/ waste etc. for the duration of the illness. Patients must remain isolated until 48 -72 hours after normal bowel habits have returned and/ or vomiting has stopped. Residents Rooms should have a full chlorine clean (1,000 per million).

All unnecessary items of equipment should be removed from rooms, to minimise the risk of contamination. This includes medical equipment and foodstuff such as fruit, biscuits and snacks.

Segregation (cohorting) may be necessary in an outbreak when single rooms may not be available for all affected persons, or the resident cannot be isolated due to dementia, or being at risk of falls. In general, however, it is important that symptomatic people are kept apart from those that are asymptomatic. In practice, this means nursing affected residents in the same area together and not admitting / transferring into empty beds in affected rooms unless the resident (being transferred) is already symptomatic or has recovered from symptoms. Staff caring for affected residents should, where possible not care for asymptomatic patients and should be allocated workload by rooms where staff numbers allow. Seriously ill patients may be particularly vulnerable to acquiring norovirus, which may worsen their underlying medical condition. In such cases it is advisable to isolate these vulnerable patients in their rooms (in effect, putting them into protective isolation) in order to minimise risk as much as possible.

All unnecessary items of equipment should be removed from rooms and bays to minimise the risk of contamination. This includes medical equipment and foodstuff such as fruit.

MOVEMENT OF SERVICE USERS IN AFFECTED AREAS

During an outbreak, service users should NOT leave the home to visit other areas unless it is essential for their clinical management. This includes attending day care facilities, rehabilitation etc. However a resident may require transferring to an acute hospital for further management of their condition. It is important that the receiving healthcare facility and transport is aware that the resident is infective and this information should be relayed prior to transfer. This will enable receiving hospitals to adequately isolate the resident on arrival thus minimising the risk of spread further.

TRANSFERS OUT OF AN AFFECTED HOME

Transfer to other hospitals

The transfer of service users to hospital during an outbreak of diarrhoea and vomiting should be avoided other than in a medical emergency, and ideally the clinician caring for the patient should agree such transfers. In such instances, staff MUST inform the receiving hospital and also the local ambulance Trust that they are transferring a resident from an area affected by diarrhoea and vomiting. This will allow ambulance personnel to take appropriate precautions and the receiving hospital to adequately isolate the resident on arrival thus minimising the risk of further spread.

TRANSFER TO OTHER CARE HOMES

No service users should be transferred out to other care homes during an outbreak unless they have been symptomatic and subsequently symptom-free for a minimum of 48 hours. If transfer is considered, it should be with medical approval only and in the full

knowledge of the receiving care home manager. Service users that have not been affected should NOT be transferred as they may be incubating the virus and could easily spread this to other healthcare settings. Occasionally, when beds are limited and an agreement has been made to receive a service user, it is advised that the service user remains in their room for the first 48hrs. Staff should use strict standard IPC precautions, with this person, during this time. This must ONLY be in exceptional circumstances and the Care Home manager must agree to the transfer. Please ensure that IPCT are aware of the transfer to allow support and advice to be given to the receiving care home.

DISCHARGES

Discharge to service users own home

Service users affected during an outbreak should not be discharged home until clear of symptoms for 48 hours. Those that have not been affected should ONLY be discharged home if the individual's carer(s) are fully aware of the likelihood of him / her becoming symptomatic and feel able to cope in such a situation. Any community care providers e.g. district nursing team should be fully informed that the individual has been discharged from an affected facility so that they can make suitable arrangements to minimise the risk of spread.

CARE HOME QUARANTINE / CLOSURE

If an outbreak is suspected/confirmed, then a decision may be made to close the home, or the affected floor, to new admissions. This must be communicated to relevant external agencies.

At such times, restrictions on movement of staff, service users and visitors are of paramount importance in order to limit spread. Staff should be allocated to asymptomatic OR symptomatic patients. Catering staff, domestics and laundry staff should not be moving from affected/unaffected floors, to reduce the risk of spread. Unaffected residents should receive their meals and have their rooms cleaned before symptomatic residents.

Care homes must remain closed for 72 hours after the detection of the last new case and the decision to re-open a home will be made by the IPCT and / or the local PHE. No home will be re-opened to admissions until thorough chlorine clean of the entire affected areas has taken place, including the changing of curtains, steam cleaning of carpets and thorough cleaning of service users' furniture and equipment especially seating, commodes, moving and handling equipment etc.

STAFF MOVEMENT

Certain groups of staff move between healthcare environments i.e. allied medical professionals, district nurses, agency nurses and medical staff. Such staff should be reminded of the importance of hand washing, both before and after care and should consider visiting affected homes / service users AFTER visiting non-affected facilities and service users. Uniforms should be changed DAILY and laundered at 60 oC or as per manufacturer's instructions. In particular, agencies providing staff should be notified of outbreaks of viral gastro-intestinal infection. This will enable them to take necessary actions to ensure their personnel do not inadvertently transmit infection to other facilities.

EXCLUSION OF AFFECTED STAFF

Exclusion is vital for any symptomatic staff member who should be sent home immediately they become affected. They should not return to work until 48 hours after symptoms have resolved. This includes bank and agency staff as well as visiting staff. It is the responsibility of the individual to ensure that they are fit to work. Staff should be advised to go to own GP and send a sample.

EXCLUSION OF VISITORS

It is important that visitors to homes during an outbreak are advised of the fact by displaying notices to all doors. If visiting more than one clinical area e.g. visiting clergy, they should be advised to visit affected areas at the end of their visit to avoid unnecessary transmission to unaffected areas / service users. In addition visitors should be advised that if they (or members of their household) have symptoms of diarrhoea and / or vomiting they should not visit the home until 48 hours after symptoms have resolved.

In certain circumstances it may be advisable to restrict / cancel all but essential visiting. This decision will be made on a case by case basis by the care homes in conjunction with PHE and IPCT. Visitors who insist on visiting must be advised and reminded to clean hands on entry to the building and wash their hands when exiting the resident's room. Gloves and aprons will only be required if they are providing direct care.

CARE HOME CLEANING

Cleaning / housekeeping staff should be made fully aware of the outbreak situation and supervisory staff / managers notified immediately there is the suspicion of an outbreak, to ensure that they are able to respond to the increased demand for cleaning in the affected areas and for additional demand for cleaning supplies etc. Cleaning should be increased to twice daily in all areas, especially paying attention to high touch areas and high risk areas, e.g. toilets taps, door handles. A standard clean using detergent should be undertaken. This should be followed by a further clean of all areas using a hypochlorite solution (Chlorine product) 1,000 parts per million. Alternatively a combined detergent / chlorine-based disinfectant solution such as Chlorclean can be used. This is more effective as only one full clean is required, and different products are not being mixed. All chlorine products should be made up as manufacturer's instructions. The majority of products can be made up once and lasts for 24hrs. Usually one tablet to 1,000 liter of water makes up 1,000 parts per million of chlorine. The chlorine solution should be made up in a designated container which is clearly labeled. Do not decant into an old bottle. The product requires to be securely locked away from vulnerable residents. Please refer to manufacturer's instructions.

- Staff must be aware of and comply with COSHH regulations when using a chlorine-based product
- All cleaning cloths must be disposable and discarded after each use. Strict attention should be paid to correct colour-coding of cleaning equipment. If possible, yellow equipment should be used in those rooms deemed to be isolation areas.
- Do NOT use the same cleaning equipment in rooms of both symptomatic and non-

symptomatic residents. A separate cloth, mop-head and bucket should be used for each area and every resident's room.

- Mop heads must be changed and laundered, or discarded after each room.
- Where service users are isolated or in communal areas, these areas must be cleaned LAST at the end of the cleaning schedule, and cloths disposed of in the infectious waste bin in that room / area.
- Aprons and gloves should be worn and removed in each room. For symptomatic patients. Apron and gloves must be disposed of in the infectious waste bin in that room when removed. Do not forget to wash hands prior to placing gloves on and when removing gloves.

SPILLAGES

(See also Spillages section)

Spillages should be dealt with immediately. Please follow manufacturer's instructions depending on the type of spill kit you have access to. Please ensure that you use the correct kit for the correct spillage. There are different types of kits for blood and for urine spillages. Chlorine fumes may be released if using hypochlorite. Protective clothing (gloves and apron) should be worn whilst cleaning spills, and discarded immediately afterwards as clinical waste.

DECONTAMINATION OF MEDICAL EQUIPMENT

(See also Decontamination of Medical Equipment section)

Where possible, all medical equipment should be dedicated for use by individual service users (or bays of affected residents) during an outbreak. Any shared equipment **MUST** be adequately decontaminated between patients. After use, with detergent and water. This should be followed by a chlorine-based disinfectant solution or a 2 in 1 product, then thoroughly dried with paper towels,

This is of particular importance for equipment such as commodes, wheelchairs, moving and handling equipment etc. that may come into contact with contaminated body fluids. Such items of equipment must be routinely decontaminated **after each and every use during an outbreak of gastro-intestinal infection.**

QUICK REFERENCE GUIDE FOR CARE STAFF DURING VIRAL GASTRO-ENTERITIS OUTBREAKS

- Ensure thorough and frequent washing of hands with liquid soap and water between all care activities and after contact with service user's immediate environment.
- Wear gloves and aprons for service user contact and environmental / equipment cleaning. Ensure hands are cleaned prior to putting gloves and aprons on and after removing them.
- Change gloves and aprons between service users / tasks, performing hand hygiene in between.
- Clean up and disinfect spillages of vomit and faeces immediately
- Pay particular attention to the cleaning of commodes, moving and handling equipment, seat raisers etc.

- Increase the frequency of routine bathroom and toilet cleaning and also cleaning of frequently touched areas (door handles, phones etc.) This also includes the dirty utility area.
- Disinfect surfaces and equipment using freshly prepared 0.1% (1000ppm) chlorine-releasing agent after cleaning with neutral detergent. Alternatively use a combined detergent/chlorine-based disinfectant e.g. Chlorclean for surfaces and equipment.
- Ensure offensive waste, clinical/Infectious waste and infected laundry are handled with care wearing protective clothing and removed promptly. Ensure hands are washed before and after glove use. Alginate bags to be used for fouled laundry
- Keep outbreak record sheet up-to-date on a daily basis
- Isolate symptomatic service users or cohort nurse groups of affected residents in the same area.
- Avoid movement of staff between affected and unaffected areas
- Exclude affected staff immediately and until asymptomatic for 48 hours
- Exclude non-essential personnel
- Avoid transfer of service users to other facilities Inc. day centers, rehabilitation etc. unless medically indicated and after consultation
- Do not re-open care home to admissions until agreed with IPCT /PHE - usually 72 hours after last new case.
- Please note that if samples are sent and all negative, the home may be able to open to admissions sooner.
- Communicate effectively and regularly to all who need to know including visitors. Provide notices indicating restrictions at entrance doors
- If in doubt, contact IPC team PHE for guidance and support.

ANTIBIOTIC PRESCRIBING POLICY

The increasing incidence of antibiotic-resistant micro-organisms and *Clostridium difficile* infections in conjunction with increased public concern about healthcare associated infections has resulted in two key initiatives being launched by the Department of Health in recent years.

Firstly, the Chief Medical Officer's annual report of 2013 and the subsequent publications of the UK Five Year Antimicrobial Resistance Strategy sets out priorities and objectives for addressing the very worrying prospect of a post-antibiotic future.

In 2017-The Chief Medical Officer-warned that if antibiotics lose their effectiveness it would spell the end of modern medicine and unless action was taken on antimicrobial stewardship the world was facing a "post-antibiotic apocalypse" and estimated about one in three or one in four prescription in UK primary care were probably not needed

Latterly, the revised *Code of Practice on the prevention and control of infections and related guidance (2018)* which provides guidance on compliance with Outcome 8 of Essential Quality Standards (Care Quality Commission) contains a criterion dedicated to addressing the issue of antimicrobial prescribing at local level. Criterion 3 states "Ensure appropriate antibiotic use to optimize patient outcomes and to reduce the risk of adverse events and antimicrobial resistance".....

Guidance on achieving compliance should include:

- ☐ Antibiotic use in primary care accounts for 80% of all NHS use and its use is expected to be both appropriate and necessary
- ☐ Local prescribing should be modified from national guidance and in particular PHE's guidance for primary care. Evidence to demonstrate adoption and adherence to policies and guidelines should be available to commissioners
- ☐ Prescribers should have access to advice on antibiotic use from prescribing advisors and microbiologists and their expertise should be used when required
- ☐ Primary care practices should participate in local and national activities designed to support antimicrobial stewardship such as back-up or delayed antibiotic prescribing and European Antibiotic Awareness Day.

Registered providers of primary care should work collaboratively with local clinicians at their acute NHS Trust(s), other local primary care providers' inc. Out of Hours providers, clinical commissioning groups and local specialists at Public Health England to ensure local antibiotic prescribing complies with the principles contained within these DH guidance documents.

Initiatives of relevance in primary care include the TARGET antibiotics toolkit. This has been developed as a national resource to help prescribers and commissioning organisations improve antibiotic prescribing in primary care.

The toolkit provides a range of resources whose ultimate aim is **Treat Antibiotics Responsibly Guidance, Education and Tools**.

Available from <http://www.rcgp.org.uk/targetantibiotics/> Last accessed 22.07.2019

Local Prescribers should have available and follow, where possible, locally produced formularies which should be modified from national guidance. Commissioning bodies may conduct antimicrobial prescribing audits and produce reports on prescribing trends. Local reduction targets may be set.

There are 4 overarching principles of good antibiotic stewardship which should be applied where applicable:

- ☐ Antibiotic therapy should only be prescribed where there is clear clinical need, e.g. the patient is showing signs of infection. The rationale for the prescription should be documented in the patient's notes. Wherever possible, antibiotics should be prescribed after clinical samples have been obtained. The prescription should be reviewed on receipt of microbiological results to ensure it is still appropriate
- ☐ Intravenous antibiotics, (rarely used in primary care settings), should only be prescribed where there is no oral alternative or where oral administration is not possible. A switch to oral antibiotics should take place at the earliest possible opportunity
- ☐ Except in rare situations antibiotics should not be prescribed for more than 7 days
- ☐ The use of broad spectrum antibiotics should be avoided where possible. These include Cephalosporins, Quinolones, broad-spectrum penicillin (including amoxicillin) and Clindamycin.

INFECTION CONTROL PRINCIPLES

The spread of infection

The spread of infection requires three elements:

- a source of infecting organisms (bacteria, viruses, fungi)
- a susceptible host
- a route of transmission of the organism from one person / site to another.

Source

The source may be service users, staff or visitors and may include persons with an obvious/unobvious symptomatic illness, or those who are asymptomatic or colonized by the infectious agent. Another source may be the service user's own microbial flora. Other potential sources are objects within the environment that have become contaminated, including health care equipment.

Susceptible Host (the individual service user, staff member, visitor)

It is important to remember that it is not only service users that may be susceptible to infection but also staff members and also visitors to the facility.

An individual's resistance to pathogenic micro-organisms can vary greatly. Some individuals may be immune to or able to resist colonization by an infectious agent, others may simply be colonized and become asymptomatic carriers, whereas others will develop a clinical disease. Persons with underlying disease such as diabetes, lymphoma, leukaemia, etc. or treated with certain antimicrobial agents, corticosteroids, irradiation or immunosuppressive agents are particularly prone to infection. Extremes of age, chronic debilitating disease, shock, coma, traumatic injury or surgical procedures and the presence of invasive devices can also make an individual more susceptible to infection.

Transmission

Micro-organisms can be transmitted by a variety of routes and the same micro-organism may be transmitted by more than one route. For example the *Varicella zoster* virus which causes chickenpox can spread via the airborne route as well as by direct contact and gastro-intestinal infections e.g. norovirus can spread by both indirect contact (with contaminated equipment and surfaces e.g. commodes and horizontal surfaces) as well as via the airborne route where virus particles are propelled through the air (and inhaled) and then drop onto surfaces where they contaminate hands and are then ingested.

There are four main routes of transmission:

- contact
- droplet/airborne
- infected food and water
- vectors

Contact transmission:

The most important and frequent means of transmission of infection can be divided into two main subgroups:

- **Direct contact:** Involves direct physical transfer of the micro-organism from person to person e.g. sexually transmitted diseases or from one site to another in the same individual e.g. bowel flora contaminating the urinary tract
- **Indirect contact:** This is the most significant route of spread in healthcare and involves contact with a contaminated object such as bed linen, instruments, equipment, dressings, etc. It is also the route by which the hands of healthcare workers transmit micro-organisms during service user care

Airborne /droplet transmission:

- **Droplet transmission:** by large droplets during coughing, sneezing, talking and during procedures which may generate droplets such as suctioning. The droplets are propelled only a short distance through the air
- **Airborne transmission:** caused by dispersal of smaller micro- organisms,e.g. viruses, contaminated water particles or airborne dust particles containing the infectious agent. These organisms can be widely dispersed by air currents before being inhaled or deposited on the susceptible host; by aerosolisation of water particles which are then inhaled e.g. in shower heads and in the case of dust particles, by airborne spread onto horizontal surfaces, equipment etc.

Food and water transmission:

Infection can occur via contaminated food or water supplies. Organisms can be transmitted via the food chain e.g. salmonella in eggs or by inappropriate handling of contaminated raw food or inadequate cooking. Secondary spread (cross- infection) can then occur if surfaces are contaminated by food-stuff e.g. chopping board used to cut contaminated poultry then used to chop salad vegetables. Additionally, infected food handlers can transfer micro-organisms on their hands to food.


Water provides an ideal breeding ground for some micro- organisms, which can then be colonized if the water supply has not been appropriately treated. In the case of *Legionella pneumophila* the bacteria forms a biofilm in pipes / shower-heads etc. and can then be dispersed in water particles and inhaled.

Vector borne transmission:

This occurs when vectors such as flies, mosquitoes, rats and other pests transmit infection. This route of transmission is rare in healthcare in the UK although it is a route of spread requiring containment in food preparation areas.

Breaking the chain of infection

The spread of micro-organisms from their source to a susceptible host is frequently referred to as the chain of infection.



The principles of infection control relate to the implementation of a series of basic control measures whose aim is to break the links in the chain thus reducing the likelihood of spread. These control measures are referred to as standard infection control precautions.

In the prevention of spread via the **direct or indirect contact** route, the following measures apply:

- Effective hand hygiene is the single most important measure in the prevention of the spread of infection
- Healthcare staff should wear suitable gloves, aprons and other protective clothing whenever there is any possibility of direct contact with infected blood, body fluids or contaminated material
- Strict adherence to the principles of aseptic technique will minimise the likelihood of contamination during the insertion and management of invasive devices and other clinical procedures such as wound care
- Effective environmental cleaning and good housekeeping techniques together with appropriate cleaning, disinfection and sterilization of medical equipment.
- Appropriate segregation and disposal of healthcare waste and contaminated laundry.

In the prevention of infection by **food and water** the following additional measures are important:

- Provision of adequate hand washing facilities, especially when handling or preparing food
- Strict adherence to food hygiene regulations. All staff who handle food should have undertaken food hygiene training.
- Healthcare environments are subject to strict flushing regime controls to minimise the risk of *Legionella pneumophila*.
- Food handlers suffering from septic conditions of the skin or gastro-intestinal infections **MUST** be excluded from work until proven to be microbiologically free from infection.

In the prevention of spread of infection by the **airborne** route the following additional measures are important:

- Adequate un-crowded housing
- Segregation of infected service users to minimise the risk of cross-infection. This is usually achieved by either physical segregation in a single room or by measures such as keeping affected service users together (cohort nursing)
- Vaccination/immunization programmes where appropriate.

In the prevention of infection by **vectors** the following information is relevant.

Whilst most people readily associate rats and mice with risks to health, the part played by cockroaches, flies and other insects is not always appreciated. They have been implicated in the transmission of infection in food stores and food preparation areas as well as in medical supplies and in the home.

STANDARD INFECTION CONTROL PRECAUTIONS (SICPs)

There is often no way of knowing which service users / clients are contaminated or infected with a transmissible micro-organism. It is essential that Standard Infection Control Precautions (SICPs) are used for all service users on every contact.

SICPs, often referred to as 'universal or standard precautions', are a single set of activities used by all staff for all service users at all times, in order to reduce the transmission of micro-organisms from both recognised and unrecognised sources of infection.

In many instances, pathogenic (disease-producing) organisms have already spread prior to the confirmation of a diagnosis. Furthermore, pathogenic organisms are frequently carried by individuals in their blood or body fluids or on the skin without signs of clinical infection – known as "colonisation". Therefore, it is important to apply appropriate precautions at all times, for all service users, rather than wait for confirmation of a diagnosis when it may be too late to prevent the spread of infection.

SICPs apply to the care of all service users regardless of diagnosis or presumed infection status, where there is possible contact with:

- blood
- all other body fluids
- secretions and excretions except sweat
- non-intact skin
- mucous membranes (conjunctivae, mouth, nose, vagina, rectum)

Standard IPC precautions include:

- Effective hand hygiene
- Wearing appropriate protective clothing- Apron, Gloves.
- Safe disposal of sharps and other healthcare waste
- Safe management of spillages.
- Prevention and treatment of sharps injuries
- Adequate and appropriate decontamination of the healthcare environment and service user-related equipment.
- Protecting cuts and abrasions on staff skin with an impermeable dressing, e.g. plaster and ensuring appropriate immunizations are up-to-date by means of routine pre-employment screening.

Guidance on implementation of specific SICPs is given throughout this document in the relevant sections.

Additional precautions

Additional precautions may be required in certain circumstances and are used in addition

to SICPs. For example service users with Pulmonary TB or Influenza may pose a risk of airborne transmission requiring respiratory precautions. Guidance on implementing additional precautions is given throughout this document in the relevant sections. Please contact the IPCT.

HAND HYGIENE

Effective hand hygiene is the single most important measure in reducing the risk of transmission of micro-organisms from one person to another or from one site to another on the same person. Decontaminating hands as per six step techniques between service user contacts and after contact with blood, body fluids, secretions, excretions and contaminated equipment/articles is essential in order to minimise the risk of cross-infection.

Hands are contaminated with both transient and resident flora:

- **Transient** flora are those micro-organisms that are not resident on the skin but are acquired by day-to-day activity including direct contact with service users, contaminated equipment and environmental surfaces. It is these micro-organisms that are responsible for the majority of episodes of cross infection. Transient flora includes the vast majority of bacteria, viruses and other pathogenic micro-organisms that our hands come into contact with during the course of daily living. This includes organisms such as *Staphylococcus aureus*, *Clostridium difficile*, gram negative bacilli and noro- viruses. Transient flora are loosely attached to the skin and are readily removed by the mechanical action of washing, rinsing and drying hands using liquid soap and water. Alcohol hand rub should not be used on hands that are contaminated, as it is ineffective. Alcohol hand rub can be used once hands have been washed as an extra precaution.
- **Resident flora** are those micro-organisms that live on the skin and provide a protective function. In the vast majority of instances these flora do not cause cross-infection and it is unnecessary to eradicate them from hands during most healthcare activities. However, in certain circumstances resident flora can pose a risk to susceptible individuals. They are a particular risk during surgery and the insertion of some invasive devices such as central venous cannulae etc. Resident flora are not easily removed by mechanical methods and require the application of skin antiseptics e.g. chlorhexidine or povidone iodine to reduce their numbers to acceptable levels. Thus the use of skin antiseptics is standard practice prior to surgical procedures and the insertion of some invasive devices

Basic hand care

To keep hands in good condition and to perform effective hand hygiene, staff should perform some basic hand care.

Use an emollient hand-cream twice a day. Use before and after shifts to help replace the skin's oils that can be lost through frequent hand hygiene. Hand creams should be for individual use or dispensed from either a wall-mounted container or from a pump dispenser. Pots / tubes of cream should not be used by groups of staff as they can be easily contaminated.

Observe the hands for any signs of damage to the skin as this can provide a portal for micro-organisms to enter the body. Cover with a waterproof plaster or dressing before the shift begins and replace if necessary. If cracks or breaks do not heal, then occupational health advice should be sought. Dermatitis can be caused by sensitivity to ingredients in hand cleansers. Always rinse thoroughly. Seek guidance from occupational health or local GP if skin problems on hands do not

clear.

All staff should be bare below the elbow when in clinical practice. Hand and wrist jewellery (including wrist watches) should not be worn by staff undertaking direct care. Rings containing stones or mounts should not be worn by care staff as micro-organisms are known to readily colonise such items providing an on-going source of potential pathogenic micro-organisms. Plain wedding bands are acceptable. Wrist watches are easily contaminated and can prevent thorough hand washing of wrists.

Nails should be kept short at all times to reduce the accumulation of micro-organisms. False nails nail extensions and nail jewellery should NOT be worn by care staff as they too are recognised sources of potential pathogenic micro-organisms and discourage staff from thorough hand decontamination.

Long sleeves should not be worn by staff undertaking direct care. In the event that long sleeves are worn, they must be rolled up above the elbows prior to hand washing and service user contact.

Types of hand hygiene/decontamination

Current research advocates a variety of processes to ensure effective hand hygiene and these are described below. The most appropriate of these processes must be used by healthcare workers depending on the work that is being undertaken.

General / clinical / social hand wash

This involves the use of liquid soap products, warm running water and disposable paper towels. This activity mechanically removes transient micro-organisms from the hands and is perfectly acceptable for the vast majority of healthcare interventions.

Alcohol-based general / clinical / social hand decontamination

Alternatively, an alcohol-based product can be used for general hand decontamination in the place of a hand-wash but only if hands are visibly clean and not soiled, or contaminated – see below.

Surgical / antiseptic scrub

This is an extended hand decontamination procedure using hand wash products containing antiseptic skin cleansers e.g. chlorhexidine or povidone-iodine. Alternatively, alcohol-based products can also be used. This type of hand wash is only required when removal of resident micro-organisms is required e.g. prior to surgical procedures and certain high risk invasive procedures.

Types of hand decontamination products

Liquid soap products

These products are used for the vast majority of hand decontamination interventions that require the removal of transient micro-organisms. Products should be purchased from an approved supplier of medical products e.g. NHS Supply Chain as these products have been independently evaluated and economies of scale will be achieved with regards to cost. Liquid soap should come in a single cartridge dispenser. Bar soap should not be used for hand decontamination by healthcare staff as it can harbour micro-organisms.

Soap impregnated/ antibacterial wipes should **not** be routinely used by health care workers who require a more thorough hand decontamination that is best provided by the use of liquid soap and running water. Soap impregnated wipes are useful in, for example, in care homes for service users, prior to meals and after using toilet facilities and in other circumstances where access to a hand wash basin is impaired.

Liquid soap products containing antibacterial agents (as are widely available in supermarkets) are not necessary for routine hand decontamination and should be avoided in health care environments.

Some soap formulations are also available as foams. These are acceptable.

Alcohol hand rub/gel

Alcohol based hand products – usually rubs or gels are currently recognised as being a known method of hand decontamination for most healthcare interventions where rapid hand decontamination is required at the point of use. However this is not effective if hands are dirty/contaminated. In this case hands must be washed to remove transient micro-organisms. Hand wipes are also useful where adequate facilities are not available, e.g. when caring for service users in their own homes.

There is a common misconception that hands should be washed after every four or five applications of **alcohol-based hand rub**. There is no reason to do this. (WHO 2009) Please assess what you have done and the risk of contamination.

Alcohol-based products should be purchased from an approved supplier of medical products e.g. NHS Supply Chain; thus ensuring that an appropriate product suitable for healthcare activities is supplied and of the required strength (usually 60-70%) and type (usually isopropanol). Alcohol products should be used from wall- mounted dispensers (see below) or can be provided for individual staff use in bottles that can be attached to uniforms thus ensuring that the product is available at the point of care.

Alcohol is not as effective as soap and water in removing *Clostridium difficile* spores or some viruses including Norovirus and must therefore not be used whilst caring for service users with diarrhoeal illness.

Antiseptic detergent products (e.g. Chlorhexidine, povidone iodine)

These products are designed for use when a higher level of antimicrobial kill is required e.g. when it is necessary to remove / reduce resident as well as transient micro-organisms. This is usually only necessary prior to surgical procedures and certain high risk invasive procedures.

In primary care facilities e.g. GP surgeries, health centres etc. antiseptic detergent products should be available where minor surgical procedures and / or Minimal Access Interventions (MAIs) are undertaken.

Hand wash facilities:

Soap and alcohol containers / dispensers

All soap and alcohol products should be dispensed from a sealed container, which delivers a measured amount of product. The nozzle must be cleaned regularly to prevent clogging and contamination. Open containers and refillable containers must not be used as they can become contaminated with micro-organisms.

Ideally, containers should be wall mounted with a pump-action and operated with heel of hand or wrist, not fingers.

Alcohol dispensers should be available at the point of care and not wall mounted adjacent to the sink

Paper towels

Good quality, absorbent paper towels should be available for use at all hand wash basins. Towels should be dispensed from wall-mounted dispensers to avoid contamination.

Hand cream

Hand cream should be available for staff use. Ideally, it should be provided in wall- mounted dispensers or from a pump-action container. Tubes or jars of hand cream must be avoided as they are easily contaminated. Nozzles must be cleaned regularly to prevent clogging and contamination.

Equipment required for effective hand hygiene in clinical settings

All hand wash basins and taps in clinical areas should conform to the requirements of Health Building Note (HBN) 00-10 (2013) *Part C Sanitary ware assemblies* which outlines the minimum requirements for such equipment. This includes the need for:

- elbow / wrist / automatically operated lever taps
- mixer taps ensuring that water is delivered at an appropriate temperature
- basins without plugs or overflows
- taps that are situated so that water does not flow directly into the waste outlet but are off-set

- taps without swan necks to minimise the potential for Legionella spp. biofilm formation in pipe-work

In primary care environments, the provision of adequate clinical hand wash basins is often overlooked. As a general rule, where-ever clinical care is provided e.g. in a clinical, treatment or consulting room as well as in dirty utility or decontamination rooms then a clinical hand wash basin should be fitted.

The following basic principles apply:

- A clinical hand wash basin compliant with HBN 00-10 should be available where-ever clinical activity takes place
- Clinical hand wash basins should be used for hand washing only and not for other purposes e.g. decontamination of equipment, disposal of body fluids including specimens or crockery/utensil cleaning.
- Clinical hand wash basins must be equipped with warm running water from a mixer tap. Separate taps are not acceptable as they do not allow for water to be delivered at the correct temperature
- Hand wash basins in clinical areas should be equipped with lever (wrist or elbow-operated) taps
- Disposable paper hand towels and liquid hand soap in wall mounted dispensers must be available at each clinical hand wash basin
- Alcohol hand gel should also be available in wall-mounted dispensers and as an individual container for each staff member
- A foot operated pedal bin should be available at each hand wash basin for the hygienic disposal of paper hand towels. (Used towels do not need to be disposed of as clinical waste unless contaminated by blood or body fluids)
- A hand washing poster demonstrating an effective hand washing technique should be displayed near hand wash basins in each clinical area

Equipment required for effective hand hygiene in home care settings

Many primary care interventions take place outside healthcare facilities e.g. in the patient's own home. Resources available for hand decontamination will vary significantly and should not be relied upon. Providing staff with personal alcohol gel dispensers facilitates hand decontamination at the bedside or in other locations where there is limited / no access to a hand wash basin. For nursing staff working primarily in patients own homes e.g. district / community nurses then a range of hand decontamination equipment should be available in portable form e.g. pouches or small cases which hold dispensers of soap and alcohol gel together with disposable paper towels. These are widely available from medical suppliers.

Hand hygiene methods

To ensure all surfaces of the hands are adequately decontaminated, it is helpful to use a standardised technique. To wash all surfaces thoroughly should take 20-30 seconds.

Some areas of the hands are more frequently missed than others during hand decontamination.

It is important to pay attention to all areas of the hands, whilst washing, but paying particular attention to the finger tips and nail area. These are the areas most in contact with the service user and can be heavily contaminated with micro-organisms.

Application of alcohol gel /rub

- ensure hands are not soiled – if necessary wash with soap and water beforehand
- dispense a measured dose of the gel / rub into the palm of one hand
- rub vigorously into all surfaces of the hand up to the wrist, using the same six technique for hand washing, until the product has dried,

Application of liquid soap

- Wet hands under running water
- Apply the recommended amount of hand cleanser
- Rub hands together vigorously to make a lather covering all surfaces up to the wrist. Follow the 6-step technique
- Rinse hands thoroughly under running water
- Dry hands thoroughly with clean paper towels
- Turn off taps using elbows or clean paper towels to prevent recontamination
- Discard paper towels into a foot operated pedal bin. Do not lift up the lid of the bin with hands as this will re-contaminate them
- If in service user's home, dispose of towels into domestic waste

Applying hand hygiene principles in clinical practice:

WHO “My five moments for hand hygiene” initiative

The World Health Organisation (WHO) concept of “5 moments for hand hygiene” has been adopted internationally as a means of providing a user- and patient- centred approach to hand decontamination with minimal complexity and across a wide range of health care settings and professions.

The concept of “5 moments” is intended to make it easier to understand the occasions (moments) when there is a risk of micro-organism transmission via the hands, to memorise these “5 moments” and to assimilate them into healthcare activities. The concept does not define specific and multiple procedures and care situations but helps focus on essential moments embedded within the care sequence that are essential for hand hygiene.

Applying the “5 moments for hand hygiene” in primary care

The need for hand hygiene is closely connected with the activities of healthcare workers within the geographical area surrounding the patient. This can be divided into two areas – the *patient zone* and the *healthcare area*.

The *patient zone* includes the patient and his / her immediate surroundings e.g. all surfaces that are touched by or in direct physical contact with the patient e.g. chair arms, walking aids, medical devices etc. It also includes all surfaces frequently touched by staff whilst caring for the service

user e.g. monitors, knobs and buttons, chair handles, computer keyboards, telephones etc. The patient zone is not static – it changes as the service user is moved from place to place and the zone accompanies the individual where-ever he / she goes e.g. from the chair to examination couch etc.

The *health-care area* corresponds to all surfaces in the health-care setting outside the patient zone i.e. other patients and their zones and the wider health-care environment. This environment still poses a risk – particularly from staff who may acquire micro-organisms within the wider health-care environment that are then transferred to service users when the staff member enters the patient zone to provide direct care. Examples include: dirty utility areas, treatment rooms, toilets, waste disposal areas etc.

In the primary care environment there are a number of occasions when clinical care is delivered in settings that are not deemed to be *health care areas* e.g. in the patient's own home. These environments cannot be controlled. However, hand hygiene CAN be controlled and should be used as the first line of defence against micro-organism transmission in ANY environment where clinical care is delivered.

When should hands be decontaminated?

Given what we now know about the environment within which the patient is cared for e.g. *the patient zone*, how do we decide **when** to decontaminate our hands?

There is a useful principle to apply:

- Assess what you have done and if you could have contaminated your hands?
- Assess what you are about to do if there is a risk of transferring micro-organisms from your hands to the patient.

5 Moments	Examples of care activity
1 Before touching a patient	<ul style="list-style-type: none"> • Before any direct contact with the patient
2 Before clean / aseptic procedure	<ul style="list-style-type: none"> • Before applying disposable gloves • Before examining a patient • Before undertaking an aseptic or clean wound dressing • Before handling / inserting an invasive device • If moving from a contaminated body site to another body site during examination / treatment of the same patient
3 After body fluid exposure risk	<ul style="list-style-type: none"> • After contact with body fluids, excretions, mucous membrane, non-intact skin or wound dressings • If moving from a contaminated body site to another body site during examination / treatment of the same patient • After removing gloves
4 After touching a patient	<ul style="list-style-type: none"> • After any direct contact with the patient • After removing gloves

5 After touching patient surroundings	<ul style="list-style-type: none">• After contact with inanimate surfaces and medical equipment in the immediate vicinity of the patient i.e. within patient zone
---------------------------------------	---

As these examples show, hand hygiene is required both **before** and **after** contact or procedure. Decontaminating hands **before** contact or procedure will protect the patient. Decontaminating hands **after** contact or procedure will protect the HCW and subsequent contamination of the healthcare environment.

Also remember that Hand hygiene should be performed before putting gloves on and after removing them.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Personal protective equipment is designed to protect the healthcare worker from coming into contact with potentially infectious body fluids. It may also protect the service user from the healthcare workers own microbial flora. Personal protective clothing includes:


- gloves
- water repellent aprons / gowns
- masks
- eye protection

Personal protective equipment is governed by Health and Safety Legislation including the Personal Protective Equipment Regulations and should only be used when risks cannot be averted by other work practices.

Risk assessment for PPE

The choice of PPE selected depends on the activity and the anticipated risk of exposure to body fluids. Many activities pose no risk of exposure to body fluids therefore there will be no need for any PPE. Risk assessment forms an integral part of Health and Safety legislation.

Assess risk of activity



Disposable Gloves

It must always be remembered that staff have a duty of care to protect their service users from risk as well as a responsibility to protect themselves. Gloves need to be changed between service users and also between tasks on the same service user to ensure that risk of transmission is reduced.

The use of latex-containing products including disposable gloves is not recommended. All healthcare providers should undertake a risk assessment relating to the provision of latex-free products to minimise the risk of inadvertent allergic reactions in those service users and staff known to be sensitive to latex and to prevent the acquisition of a sensitivity reaction in at-risk individuals e.g. those with known skin conditions such as eczema, dermatitis etc.

In addition to effective hand hygiene, disposable gloves of the recommended type play an important role in reducing the risks of transmission of micro-organisms.

Gloves are worn to:

- reduce the likelihood of micro-organisms being transmitted to service users during invasive or other care activities
- reduce the likelihood that hands of personnel contaminated with micro-organisms from a service user or equipment can transmit these organisms to another service user
- provide a protective barrier and to prevent gross contamination of the hands when anticipating contact with blood, body fluids, secretions, excretions, mucous membranes and non-intact skin
- protect staff from potentially harmful organisms

Glove use

Hand hygiene should be performed prior to donning gloves

Non sterile, powder-free latex or synthetic latex e.g. nitrile and vinyl gloves should be worn whenever contact with body fluids, contaminated equipment, non-intact skin or mucous membranes is anticipated.

Sterile, non-powdered, latex or synthetic latex e.g. nitrile gloves which provide greater dexterity and tactility are available for surgical and other invasive procedures requiring sterile gloves. These are supplied in a range of sizes for accurate fit.

For the majority of routine clinical tasks vinyl gloves provide adequate protection and should be the glove product of choice

Gloves are not required when handling unsoiled articles or for contact with intact skin in the absence of body fluids.

Gloves must be removed at the end of each individual procedure/healthcare activity, and hands

washed thoroughly.

It is essential to keep the time of wearing gloves to a minimum to avoid skin sensitization. Staff experiencing skin conditions which may be exacerbated by glove wearing should contact Occupational Health or their GP for further advice / assessment.

Disposable plastic aprons

Plastic aprons should be worn to protect staff uniform/clothing when contamination with body fluids is possible during healthcare procedures. This may include:

- testing urine specimens
- undertaking wound dressings

In addition, a plastic apron should be worn during the following activities to minimise microbial contamination of clothing:

- during environmental cleaning or decontaminating/cleaning equipment
- when handling used/soiled linen

Always remove the apron at the end of each care-giving procedure and discard into a waste bag, and wash and dry hands to reduce the likelihood of transferring organisms to another site.

Water repellent (sterile) surgical gowns

During Minimal Access Interventions (MAIs) and *some* minor surgical procedures (where a sterile device is being implanted) or when there is a risk of significant post- procedure infection then it is recommended that a sterile (water repellent) gown is worn to minimise the risk of surgical site contamination (Humphreys H., Coia J.E. et al (2012) *Guidelines on the facilities required for minor surgical procedures and minimal access interventions* Journal of Hospital Infection 80 103 – 109)

Face masks / eye protection

These are worn when there is a possibility of splashing of blood or body fluids or chemical/detergents into the eyes and/or mucous membranes. Face masks, goggles, safety glasses or shield masks are all suitable products and the most appropriate should be chosen and should be readily available for staff. If these products are disposable they should be disposed of as clinical waste or if non- disposable, cleaned as recommended in the disinfection policy/manufacture's recommendations, usually with detergent and warm water. Managers should ensure that appropriate masks and eye protection are available for staff use.

Face masks are not usually required during minor surgical procedures except when a sterile device is being implanted or when there are other issues predisposing to infection.

In certain circumstances, respiratory masks may need to be of increased efficiency in order to minimise the risk of transmission of highly infectious micro-organisms. Currently this includes pandemic influenza; some cases of sputum-positive pulmonary TB e.g. MDRTB. Current

guidance recommends the use of FFP3 respiratory masks which provide 99% particle filtration efficiency. These must conform to European Standard EN149 2001 (box is CE marked) and must be worn when exposed (within 3 feet of a service user). These masks are single use only. The Health and Safety Executive recommends that staff who are required to wear FFP3 masks are fit tested to ensure that masks adequately fit the individuals' face thus minimising the likelihood of infected respiratory droplets leaking through or around the facemask.

Influenza - FFP3 Masks are only required when performing aerosol generated procedures on a suspected/ confirmed Influenza (Flu) patient. For direct care/ close contact with suspected or confirmed flu, within one metre, a surgical face mask will suffice. Please note that these will become saturated after twenty minutes and may need to be changed. Staff who are immunocompromised and/or have not had a flu vaccination will need to seek further information.

Storage of PPE

All personal protective equipment (PPE) should be stored appropriately to minimise the risk of contamination prior to use.

Wall-mounted dispensers are available for the hygienic storage and dispensing of both disposable gloves and plastic aprons. These are recommended for use in primary care facilities where routine clinical interventions are undertaken for example examination, consulting or treatment rooms.

Hands should be washed and care should be taken when removing disposable gloves from boxes in order to minimise the risk of contaminating the contents with unwashed hands.

SAFE USE AND DISPOSAL OF SHARPS

See also Section 10 – Management of Healthcare Waste

Many needle-stick injuries are preventable providing staff are informed of the appropriate procedures which will minimise the risks associated with handling sharps. The following practices should be taught to all staff likely to handle sharps, at induction/orientation and regularly thereafter.

Non-compliance with these guidelines may carry medico-legal or health and safety implications.

DEFINITIONS

Clean / used sharp describes a sharp that has been used for a “clean” procedure such as drawing up injections. Such a sharp will not have had contact with a service user’s blood or body fluids and poses less of a risk to the HCW should a sharps injury occur, although from a Health and Safety perspective such injuries are still of significance.

Contaminated / dirty sharp describes a sharp that has been used invasively and has had contact with a service user’s blood or tissues thus posing a higher risk of potential cross-infection with a blood-borne virus should a sharps injury occur.

COLOUR-CODING OF SHARPS CONTAINERS

Sharps containers are colour coded to indicate their contents and the route for final disposal by incineration. In brief, the colours of sharps container lids show, by means of colour coding, the contents of the container and which waste stream they are required to enter for final disposal by incineration. Three colour-coded sharps waste streams apply:

- Yellow sharps container with a yellow lid. Sharps for incineration - partially discharged sharps including those contaminated with medicines other than cytotoxic and cytostatic, or where a mix of sharps is likely.
- Yellow sharps container with an orange lid. Sharps for incineration or alternative treatment.
- Sharps not contaminated with medicinal products, or cytotoxic residues. Contaminated with blood or saline / dextrose products e.g. in phlebotomy then an orange lidded container can be used.
- Yellow sharps container with a purple lid. Sharps for incineration Sharps including those contaminated with cytotoxic and cytostatic medicines.

At local level this requires providers of healthcare to assess the sharps that they generate and to ensure that appropriate colour-coded containers are used. This will usually be undertaken at a strategic (organisation-wide) level in discussion with the registered waste contractor responsible for the collection and ultimate disposal of clinical waste.

The rationale for colour coded lids relates to the fact that POMs and cytotoxic / cytostatic medicines are classed as hazardous waste and require separate licensing, transportation and final disposal arrangements.

*Many users are unaware of the wide range of cytostatic medicines currently in use. A comprehensive list of common drugs can be found in HTM 07-01 page 167.

ASSEMBLY OF SHARPS CONTAINERS

Only approved sharps containers must be used which comply with current standards. (BS 7320:1990, UN 3292)

Ensure that the sharps container is correctly assembled and that the lid is securely fitted. Follow the manufacturer's recommendations for assembly, as all containers differ. Label the sharps container with the date of assembly, the name of the member of staff who assembled it and location e.g. GP practice name.

PROVISION AND LOCATION OF SHARPS CONTAINERS

Adequate sharps containers must be available in all healthcare facilities where sharps are in use. Ideally, they should be available in all places of regular use i.e. at the point of use such as treatment and consulting rooms.

Containers should be available in a range of sizes appropriate to the number of sharps generated and where they will be used. For example, small containers (designed for disposal of needles only) are available for portable use at the bedside or for home care. Many can be fitted to sharps trays specifically for this purpose. Large containers should only be used when a high volume of sharps are generated e.g. phlebotomy.

All sharps containers must be stored out of the reach of children and others who may be at risk.

Sharps containers must never be stored on the floor or above shoulder level. They should be ergonomically positioned between waist and shoulder height to allow ease of access and to ensure the lid of the container can be visualized to avoid sharps injuries from over-full containers.

Sharps containers should be placed on a secure, stable surface and away from the edge of work surfaces. Most manufacturers can supply brackets to mount them on the wall or trolleys for ease of movement e.g. in minor surgical procedure rooms.

Wherever possible, sharps containers must be taken to the point of use to ensure immediate disposal. Small, portable containers, ideally mounted on trays provide a suitable mechanism for such use.

The temporary closure mechanism should be activated when sharps bins are left unattended and whenever a sharps bin is transported.

SAFE DISPOSAL OF USED SHARPS

It is the responsibility of the individual who has used the sharp equipment, to safely dispose of it in an approved container. Sharps must not be left for others to clear away.

Place all disposable sharps into an approved (BS 7320:1990, UN 3292) puncture proof sharps

container immediately at the point of use. Always ensure you have a bin ready to use prior to using a sharp. All containers have a temporary closure, which should be activated between uses

Re-sheathing sharps should NEVER occur. Recent EU legislation has banned re-sheathing with immediate effect (2010/32/EU).

Do not attempt to remove the needle from the syringe. Discard the needle and syringe as a single unit, into an approved sharps container.

Fill sharps containers to the 'fill' line only. Do not overfill any sharps container, as this is a significant risk to both you and others.

When full to the "fill" line or the bin has been opened for 3 months the permanent locking mechanism should be activated and the container then labelled with the date, name of the person disposing of the full container and the location details e.g. GP practice name.

Full sharps containers should be kept in a dedicated, lockable, area. Full containers must NOT be placed inside clinical waste bags.

SERVICE USERS OWN SHARPS

Many service users self-administer medications e.g. diabetics. A variety of administration and monitoring systems are available including pens as well as needles, lancets and syringes. All systems involving the use of sharps have the potential to cause injury if handled inappropriately.

Service users self-administering medication must be supervised and trained in safe practices prior to being allowed to self-medicate.

Appropriate equipment must be provided for the service user either by their GP or hospital consultant / nurse specialist (now on prescription). Small portable sharps boxes complying with relevant standards should be used. These must be returned to the service user's GP practice / pharmacy if distributed from there for disposal as hazardous waste or arrangements for collection should be made by the GP responsible for the patient. Care must be taken to ensure returned sharps boxes are transported appropriately by the service user to minimise risk to the individual and members of the public.

Service user's own sharps must not be disposed of into the household waste stream. This is no longer acceptable. This includes lancets used for blood glucose analysis.

Care must be taken by staff using self-administration systems on behalf of service users. An assessment of risk must be undertaken especially regarding needle disposal.

TRANSPORTING SHARPS CONTAINERS

Healthcare workers producing sharps waste in non-NHS environments e.g. in the patients' own home may be required to transport the sharps waste back to base in some circumstances (e.g. where such interventions are temporary and the householder does not have a waste collection arrangement in place).

Sharps waste must be transported in suitable UN-approved rigid sharps containers (as would be used in healthcare environments). 0.6litre bins should be used with a temporary mechanism that clips into place. The bin should be carried in a sturdy transport container. Sharps bins should be taken into the house to allow disposal of the sharp at the point of care. These must be provided by the healthcare provider. If the healthcare worker is travelling by public transport (or bicycle) then arrangements must be made to collect such sharps boxes from a suitable location. They should not be transported by such means.

PROTECTIVE CLOTHING AND VENEPUNCTURE

Gloves must be worn when handling or using sharps.

Gloves cannot prevent needle-stick injuries but they may reduce the likelihood of infection by reducing the volume of blood inoculated during the incident.

Some individuals highly experienced in venepuncture may prefer not to wear gloves because of a perceived reduction in manual dexterity. However, all experienced staff and new trainees, including doctors, should be trained and advised to wear gloves whilst taking blood in line with expert guidance.

The following is advised:

- Gloves must always be available for venipuncture. Hands should be cleaned prior to putting gloves on, and after removing gloves, after each patient
- All staff should receive education to wear gloves from the beginning of their training.
- If staff have cuts, abrasions or skin lesions on their hands which cannot be covered by waterproof dressings they should be referred to occupational health.
- Gloves should also be worn if the service user is uncooperative or restless

(Expert Advisory Group, 1998)

RISK ASSESSMENT OF WORK PRACTICES SAFETY ENGINEERED PROTECTION MECHANISMS

EU legislation (2010/32/EU) requires healthcare providers to undertake a risk assessment of all situations where there is injury, blood or other potentially infectious material. This includes measures designed to eliminate exposure risks and the consideration of possible alternative systems. This legislation came into force in England from 11 May 2013 as the Health and Safety (Sharp Instruments in Healthcare) Regulations 2013.

With regard to sharps use, there is a requirement to eliminate unnecessary use of sharps and, if risk assessment identifies that risks exist, then the provision of medical devices incorporating safety engineered protection mechanisms must be considered. These include:

- needleless intravenous systems
- syringes with advanceable needle guards or retractable needles
- self-sheathing trocars etc.

The Health and Safety Executive have produced an information sheet for employers and employees on the new legislation which can be found on their website at www.hse.gov.uk

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

MANAGEMENT OF HEALTHCARE WASTE - SUMMARY

INTRODUCTION

This waste summary has been written to provide guidance on managing waste in health and social care environments. Guidance is drawn from Health Technical Memorandum (HTM) 07-01: *Safe management of healthcare waste (2013)*. This guidance provides a number of changes designed to introduce cost savings, safer working practices and a reduction in carbon emissions related to managing waste. Relevant changes include:

- Updates to legislation, specifically for environmental permitting and transport/carriage regulations;
- A focus on the waste hierarchy through procurement practices, and the elimination, minimisation, recycling and recovery of waste;
- A drive to address the carbon impact related to waste through resource efficiency, transport impacts and disposal arrangements;
- The integration of new sector guides on GPs and dental practices as well as incorporating HTM 07-06: Disposal of pharmaceutical waste in community pharmacies as a sector guide
- A focus on practical advice and examples for classifying waste, in particular the infectious and offensive waste streams, including case studies to highlight best practice;
- A review of the terminology used for healthcare, clinical and non-clinical wastes.

This protocol is intended to provide organisations and their staff with insight into the legislation and regulation that applies to waste management and in particular provides guidance on infection control related elements of clinical waste management. **This document is NOT intended as a waste management policy.**

Organisations/independent contractors should produce a Waste Policy and Strategy which will ensure compliance with the requirements outlined below.

Good waste management is important for the following reasons:

- to reduce the health and safety risk to staff, service users and visitors from waste;
- to manage waste disposal costs and reduce where appropriate;
- to ensure compliance with environmental legislation which includes the reduction of carbon impacts of managing waste.

LEGISLATION AND REGULATION

To effectively manage waste generated, those responsible for the management of the waste should understand and comply with the requirements of different regulatory regimes:

- Health and Safety;

- Environment and waste;
- Medicines Management;
- Infection Prevention & Control;
- Transport.

The management of healthcare waste is directed by statute and regulation from the United Nations, European Union, UK parliament and devolved national parliaments. Such legislation and regulation is regularly reviewed and re-issued. For waste management practices to comply with these requirements, appropriate waste management services need to be procured. Organisations procuring such services should be aware that, under the Environmental Protection (Duty of Care) Regulations (England Scotland and Wales) contained within the Environmental Protection Act 1970, they have a duty of care for the safe management of waste “from cradle to grave” and not just within their own premises

Organisations that produce waste are required to register with the Environment Agency as a waste producer. This registration process should commence with an assessment of the types of waste to be produced and audit of same (pre- acceptance audit). Specialist advisors (Dangerous Goods Safety Advisors, DGSA) may be required depending on the volumes and types of waste generated.

It is recommended that a Waste Manager is identified to lead the production of a Waste Policy and Strategy which will include sourcing appropriate advice. Guidance on policy production can be found in HTM 07-01 - *Safe Management of Healthcare Waste section 6: Managing Compliance*.

RESPONSIBILITIES OF THE GENERAL PRACTICE

This section is reprinted directly from: Health Technical Memorandum (HTM) 07-01: Safe management of healthcare waste - sector guide: General Practices and Health Centres 141-151.

General medical practices have a statutory duty of care. This applies to everyone in the waste management chain from producer to disposer. It requires the practice to manage the waste and to take all reasonable measures to ensure that the waste is dealt with appropriately from the point of production to the point of final disposal. The general practice’s responsibilities do not end when it hands its waste to a waste collector.

The practice is solely responsible for ensuring that waste is:

- Correctly segregated;
- Appropriately labelled;
- Packaged appropriately for transport;
- Stored safely and in a secure place away from areas of public access within the premises (that is, taking all reasonable precautions to prevent waste escaping and to prevent the public getting access to it – this could be a fenced, locked compound);
- Described accurately and fully on the accompanying documentation when removed;
- Transferred to an authorised person for transport to an authorised waste site.

In addition the general practice should ensure that:

- Each of its premises is registered as a hazardous waste producer (unless exempt from registration); and
- It keeps a register of the necessary records and returns in the appropriate location (normally the practice's premises)

The practice manager should also ensure that staff are trained and aware of the local waste procedures.

The waste management contractor should be willing to provide advice on fulfilling the requirements for the above responsibilities. However:

- **It remains the legal responsibility of the practice**, not the waste contractor, to ensure full compliance; and
- the waste contractor will have less knowledge than the practice about what is in the waste

HEALTH AND SOCIAL CARE ACT (2008) and CODE OF PRACTICE

The Code of Practice for the prevention and control of infections and related guidance (2015) applies to the safe handling and disposal of waste (criteria 2 and 9). This can be achieved by the following:

1. Risks from waste disposal are properly controlled by:
 - Assessing risk
 - Developing appropriate policies
 - Putting arrangements in place to manage risks
 - Monitoring, auditing and reviewing the way in which arrangements work and
 - Being aware of statutory requirements, legislative change and managing compliance
2. Precautions should be in place when handling waste including:
 - Training and information
 - Personal hygiene; immunization and PPE
 - Segregation and storage of waste
 - Appropriate procedures for handling waste
 - Appropriate packaging and labelling
 - Suitable transport on-site and off-site
 - Clear procedures for accidents, incidents and spills and
 - Appropriate treatment and disposal of waste

3. Systems should be in place to ensure that the risks to service users from exposure to infections caused by waste present in the environment are properly managed, and that duties under environmental law are discharged. The most important of these are:
- Duty of care in the management of waste
 - Duty to control polluting emissions to the air
 - Duty to control discharges to sewers and
 - Obligations of waste managers
 - Collection of data and obligations to complete and retain documentation including record keeping
 - Requirement to provide contingency plans and have emergency procedures in place.

There is a unified methodology and definitions that will allow everyone who handles waste to determine whether the waste fits in to one of the following defined categories;

- Infectious clinical waste.
- non infectious clinical waste
- hazardous waste
- offensive/hygiene waste
- waste that is dangerous for carriage

This unified approach has been developed to enable those involved with waste management to comply with waste regulations. While it is not mandatory to comply with this unified approach it is considered best practice.

SEGREGATION OF WASTE

Segregation of waste into separate streams ensures appropriate and safe disposal in order to reduce costs and treat waste appropriately.

It is essential that all staff are aware of and comply with safe methods of disposal which should be clearly documented in local procedures.

Segregation can be easily achieved by careful use of the correct receptacles (bags and bins), together with appropriate storage prior to collection.

It is the responsibility of the person who disposes of an item to ensure that it enters the waste stream in the correct receptacle.

DEFINITIONS OF WASTE

Clinical and Hazardous Waste

The definition of clinical waste (as defined by the Controlled Waste Regulations – issued under the Environmental Protection Act) is:

1. ...any waste which consists wholly or partly of:

- human or animal tissue
- blood or other body fluids, excretions
- drugs or other pharmaceutical products
- swabs or dressings
- syringes, needles or other sharp instruments

....being waste which unless rendered safe may prove hazardous to any person coming into contact with it; **AND**

2. ...any other waste arising from medical, nursing, dental, veterinary, pharmaceutical or similar practice, investigation, treatment, care ...being waste which may cause infection to any person coming into contact with it.”

Clinical waste can be divided into three broad categories of materials:

- any healthcare waste which poses a risk of infection (and thus by definition possesses a hazardous property categorised as H9 infectious)
- certain healthcare wastes which pose a chemical hazard
- medicines and medicinally contaminated waste containing a pharmaceutically active agent

Offensive/hygiene waste describes waste that is non-infectious and which does not require any specialist form of treatment or disposal. In the past this has been described as Human Hygiene or Sanpro Waste. Offensive/hygiene waste is healthcare waste (or similar from municipal sources) which meets the following criteria

- It is not clinical waste
- It is not dangerous for carriage
- The producer has identified, after segregation at source, that it is suitable for disposal at a non-hazardous landfill site without further treatment
 - It may cause offense to those coming into contact with it

Items that are considered to be offensive/hygiene waste are;

- incontinence and other waste produced from human hygiene
- sanitary waste
- disposable medical items and equipment that do not pose a risk of infection, including PPE (that is items that are not clinical waste)
- nappies

Such waste must be assessed for medicinal, chemical or infectious properties before being assigned to this category.

Sharp waste is defined as any item that could pierce the skin. This includes: needles, broken crockery and glass

Items that may explode on incineration must not be disposed of as clinical waste, but must be decontaminated before disposal as per local authority guidance. This includes aerosol cans (even if empty) and batteries.

Other Waste streams

There are other waste streams which do not carry infection risks but are covered by regulation. These streams should be defined in the organisations Waste Policy.

National colour-coding approach

Segregation of waste at the point of production into suitable colour-coded packaging is vital to good waste management. Health and Safety, carriage and waste regulations require that waste is handled, transported and disposed of in a safe and effective manner. The following colour-coded waste segregation guide represents best practice and ensures, at minimum, compliance with current regulations.

Proper segregation of different types of waste is critical to safe management of healthcare waste and helps control management costs. The use of colour-coded receptacles is an essential element of good segregation practice.

The national waste colour-coded segregation system identifies and segregates waste on the basis of waste classification and suitability of treatment/disposal options.

Appendix A summarises the colour-coding system currently in use.

Waste Minimisation and Carbon impact

The guidance on which this chapter is based stresses the importance and need to minimise both the volume of waste produced and also the carbon impact of waste disposal methods used. Thus consigning all waste as clinical for incineration is not considered acceptable. Waste assessments and strategies should be devised to allow minimisation of both waste quantities and carbon impact. This potentially benefits the organisation in cost savings as well as the environment. Further guidance on achieving this can be found in the source document (HTM 07-01 - Safe Management of Healthcare Waste section 5). Additionally advice may be sought from the Waste Contractor.

APPENDIX A WASTE STREAMS – INTERPRETATION

Anatomical Waste – yellow stream.

Anatomical waste – Yellow stream requires disposal by incineration in a suitably licensed or permitted facility. This waste stream includes anatomical waste and may include other types of waste which require incineration to comply with national or regional policy, including un-autoclaved waste from clinical laboratories. This waste stream also includes waste that is, or may be contaminated with infectious micro- organisms *but which also has an additional characteristic that means it must be incinerated*. For example: anatomical waste; medicinally-contaminated infectious waste etc. **This waste stream should NOT be used solely for known/suspected infectious waste. Such waste should be treated as infectious waste and placed into orange bags**

Yellow-stream infectious waste is hazardous waste and is subject to the controls of the Hazardous Waste Regulations.

Anatomical Waste – red-lidded receptacles.

Waste which contains recognizable body parts should be incinerated in suitably licensed premises. Containers for such waste are yellow with red lids.

Infectious Waste – orange stream

Infectious waste – orange stream may be treated to render it safe prior to final disposal to landfill. Treatment may only take place in a suitably licensed facility. Orange-stream infectious waste is known or suspected to contain pathogens and is hazardous waste subject to the controls of the Hazardous Waste Regulations. ***The orange clinical waste stream should NOT contain waste that is non-infectious***
e.g. offensive and domestic waste or that has additional characteristics that require incineration e.g. medicinal, chemical, anatomical characteristics.

Under the Landfill Regulations it is prohibited to send infectious waste direct to landfill for disposal without prior treatment.

Infectious Liquid Waste – yellow or orange receptacles

Infectious liquid waste should be contained in rigid receptacles for disposal. Some contractors require such waste to be solidified before removal.

Offensive / Hygiene Waste

Offensive/hygiene waste is disposed of by deep landfill. Such waste is collected in yellow / black striped bags – so-called “tiger stripe” bags.

Sharps Waste

Sharps are items that could cause cuts or puncture wounds including needles, syringes with needles attached, broken glass ampoules, scalpels and other blades and infusions sets. Sharp items such as needles attached to syringes that contain, or may potentially contain residues of Prescription Only Medicines (POMs) are also subject to classification under the Special Waste Regulations as Pharmaceutical waste (see below) and must be discarded into appropriate sharps bins with colour- coded lids. See Safe Management of Sharps section of this Manual.

Domestic (household) Waste

Domestic waste is waste that is similar to the waste generated at home. It should not contain any infectious materials, sharps or medical products and may be placed in either black or clear bags for disposal.

Pharmaceutical Waste

Pharmaceutical waste is described as waste containing a pharmaceutically active agent. This may include expired or unused medicinal product, and discarded items associated with medicines e.g. bottles, connecting tubing, syringes etc.

Pharmaceutical waste is further divided into Cytotoxic/Cytostatic waste and non-Cytotoxic/Cytostatic waste.

All pharmaceutical waste must be disposed of into an appropriately coloured pharmaceutical waste container. This is blue for non-Cytotoxic / Cytostatic waste and Purple for Cytotoxic / Cytostatic waste.

WASTE RISK ASSESSMENT AND SEGREGATION

In England and Wales, mixing of waste is prohibited by law. This means that waste **MUST** be segregated into appropriate waste streams prior to disposal. This requires waste to be risk assessed *on a case by case basis* and thus requires a waste provider to ensure that a full range of waste streams is available for use when required.

On a daily basis this means that it is not acceptable to dispose of all clinical waste into an orange waste stream or into a yellow/black offensive waste stream but that both streams must be available for use. Both streams are acceptable for the disposal of waste contaminated with body fluids **BUT** the orange stream should only be used *if the body fluids are suspected / known to be infectious*.

This aspect of waste disposal is the most commonly misunderstood element of the waste cycle and requires a comprehensive understanding of waste management by those responsible for waste policy, together with easily understood local protocols supported by staff training.

Comprehensive guidance and further explanations can be found in HTM 07-01 – Safe management of healthcare waste: sector guide on General Practices and Health Centres page

141.

EFFECTIVE DISPOSAL OF WASTE

For effective disposal of waste it is important for consideration to be given to the placement of waste receptacles. Waste must be disposed of as close to source as possible and bins must be positioned where they are easily accessible to staff. Clinical waste bins should not be placed where visitors/service users may use them for the disposal of domestic waste.

Bins should be colour-coded or clearly labelled, fire retardant and fully enclosed with lids which must be foot-operated. All bins should be in good working order.

When bins are two-thirds full the bags must be removed, securely tied and, if appropriate, labelled in accordance with the legal requirements for transporting and packaging waste (to ensure traceability) and removed to a designated waste storage area or bin. In healthcare facilities clinical waste bags should be secured with a tie and not by knotting.

The storage area or bin must be lockable (for clinical waste) and free from access to the public, pests or vermin. Waste streams should be clearly segregated in storage areas.

Domestic waste bags must also be changed when two-thirds full, secured and stored in a designated area separate from clinical waste.

Sharps bins, when full, must be closed securely and labelling completed prior to disposal. Sharps bins must NOT be placed inside yellow / orange bags but should be stored in a locked storage area.

When handling any waste bag the bag must only ever be held by the neck.

STAFF PROTECTION

When handling clinical or hazardous waste staff should always wear appropriate protective clothing i.e. apron/overalls and gloves.

When such waste handling is complete protective clothing must be disposed of in to the clinical waste stream.

Hands must be thoroughly washed and dried after protective clothing has been removed.

All staff handling clinical waste must be offered a programme of vaccinations for Hepatitis B, Hepatitis A and Tetanus. (See section - Vaccination Programme for Staff and Service Users)

All staff must be aware of the policy for exposure to blood-borne viruses and take the appropriate action after an incident. (See section - Management of Occupational Exposure to Blood Borne Viruses)

SPILLAGE

All spillage must be regarded as potentially hazardous and dealt with immediately.

Under no circumstances should service users or members of the public be allowed to assist, or be involved in any way in the clearing or cleaning up of spillage.

When dealing with spillage, protective clothing (gloves and apron) must be worn.

If it is possible, ask another member of staff to assist in keeping unauthorised persons away, until the area can be barricaded off.

If dealing with a broken or split bag, re-bag the contents and ensure that the area is free of waste.

If sharps are present, puncture proof gloves/gauntlets must be worn. A pair should be available in all areas where clinical waste is handled.

If the area has been contaminated with blood or body fluids clean the area well with a solution of detergent and warm water, followed by a hypochlorite (Chlorine) based disinfectant. (See section – Spillages of Blood and Body Fluids).

After any spillage always thoroughly wash and dry your hands.

Spillages of clinical/hazardous waste should be reported using the organisation's incident reporting processes with an investigation being undertaken to identify risks and allow risk reduction actions to be implemented.

AUDIT AND MONITORING

The Code of Practice on the prevention and control of infections (2010) requires a programme of audit to demonstrate and ensure compliance with policies. Waste management guidance also requires an audit programme of waste segregation and storage arrangements. This should include quarterly observation of waste containers (without handling the waste itself) as a minimum. Additional and more detailed audits of container contents are advised at intervals determined by the volume and types of waste produced. Such audits require careful risk assessment and the application of control measures to ensure the safety of auditors. Such control measures will include, but not be limited to, the use of Personal Protective Equipment.

TRAINING

All staff having contact with waste whether through the production of waste or disposal must have training in safe management of waste and local policies. Staff should be trained at induction and regularly thereafter.

RECORD KEEPING

The manager with designated responsibility for waste disposal must keep records that include

details of the waste disposal contract and records of all clinical waste collections from the healthcare premises. Waste transfer and consignments notes for hazardous waste should be retained for 3 years.

ENVIRONMENTAL CLEANING

INTRODUCTION

All staff have a responsibility to promote and safeguard the wellbeing and interests of service users. A dirty cluttered environment is not a standard on which any health care organisation wishes to be judged.

Cleaning is necessary to maintain the appearance, structure and efficient function of the environment and equipment. It is also required to control the microbial population and to prevent the transfer of certain micro-organisms. Cleaning, when performed effectively and regularly, is often all that is necessary to minimise the risk of cross- infection.

Standards of environmental cleaning services should be audited monthly to ensure compliance with local schedules and processes as laid down in the National Specifications for Cleanliness in the NHS: *Guidance on setting and measuring performance outcomes in primary care medical and dental premises* (NPSA 2010).

Where environmental cleaning services are out-sourced to a third party contractor, local arrangements for regular audit against the contract should be undertaken by the healthcare provider. Where primary care services are undertaken in third party shared / rented premises i.e. within health centres the Practice should satisfy itself that appropriate standards are being maintained in accordance with relevant national specifications.

STAFF PERSONAL HYGIENE

Personal hygiene is important. Hands should be washed frequently and especially after each cleaning operation, to ensure that harmful organisms are not spread.

It is important that domestic staff report to their line manager any infections which they have or have come into contact with. (See section: Management of Infections in Staff).

Adequate and appropriate protective clothing must be available for domestic staff at all times including household gloves and plastic aprons. Staff should be trained in the use of PPE and the frequency for change of equipment.

GENERAL HYGIENE

Regular cleaning and attention to cleaning processes does more to remove environmental bacteria than any other activity, including the type of cleaning agent used.

Stained, dusty or unhygienic surroundings combine to produce an unattractive and sometimes high-risk health care environment.

Cleaning equipment should be cleaned thoroughly after use and stored dry in a clean secure place. Mops should not be left soaking as the water acts as a reservoir for micro-organisms. Mops must be disposable or changed after each use a fresh mop should be stored head uppermost ideally using wall-mounted brackets. Mop heads should be either disposable or

laundered regularly dependent on local risk assessment.

Appropriate protective clothing should be worn when carrying out cleaning processes, appropriate gloves (powder-free) and plastic aprons. Face protection should be available to staff handling disinfectants in compliance with Health and Safety and COSHH regulations.

COLOUR CODING OF EQUIPMENT

The aim of colour coding is to ensure that cross-infection does not occur when cleaning equipment is used in more than one type of area. Using a cloth in a consulting / treatment room following its use in the toilet would provide considerable risk of cross- contamination on environmental surfaces.

Colour coding should be applied to all housekeeping equipment in all areas of the organisation. All staff, especially domestic and healthcare staff should be familiar with the colour coding in use. Posters demonstrating this should be available for staff as a reference tool. Ideally, colour coding of housekeeping equipment should reflect the guidance issued by the National Patient Safety Agency (NPSA 2007). A chart is provided at the end of this section.

USE OF DISINFECTANTS

Disinfectant solutions must only be used by staff that have been trained in their use and are aware of how to prepare the solution (including dilution), how to use the solution, what protective clothing must be worn and how to dispose of the solution after use. They must be aware of the COSHH regulations for the disinfectants used and have access to data sheets which are available from the product manufacturer. A folder containing COSHH data sheets must be kept in all areas and be available for staff to refer to at all times.

Research has shown that efficient routine cleaning using a general purpose liquid detergent will remove a high proportion of micro-organisms, including bacterial spores and in most situations thorough cleaning will be adequate. Chemical disinfectants are not cleaning agents and to use them as such is unnecessary and wasteful as well as potentially harmful. Chemical disinfectants would be used in an outbreak or cluster situation.

All disinfectants must be adequately labelled with the active ingredients in case of accident/splash/ingestion in accordance with COSHH regulations.

Gloves and plastic aprons must always be worn when handling disinfectants. Eye protection must also be available.

A decision should be made by the facility to use the same disinfectant preparations throughout the building to ensure consistency and economies of scale.

Decisions relating to the use of disinfectant solutions should be made in collaboration with the local IPCT to ensure use is appropriate.

Preparations should be available in the correct concentration. designated containers should be labelled accordingly. A Hypochlorite concentration of 10,000 ppm (parts per million) is necessary for use on blood and body fluid spillages. A weaker concentration of 1,000 ppm is used for

environmental disinfection (where appropriate).

Usually, the type of disinfectant solution required to deal with high risk situations can be restricted to a specific chlorine-releasing agent that is highly effective against bacteria, bacterial spores, viruses and other relevant pathogens.

An alternative system of environmental cleaning is the use of Microfibre cleaning systems which negate the requirement for the use of environmental disinfectants. Many commercial companies provide Microfibre systems which are widely used in NHS premises and are currently being evaluated with regards to their efficacy.

Where Microfibre systems are used there should be protocols in place. These should include, as a minimum:-

- Colour coding of cloths/mop heads
- Frequency of change of cloths/mop heads i.e. per room/bed space
- Maximum time of use/reprocessing of cloths/mop heads
- Method of laundering of cloths/mop heads
- Management of Microfibre laundry facilities

Microfibre systems should not be used during an outbreak situation, as the evidence is still not conclusive

STORAGE OF CLEANING EQUIPMENT

Cleaning equipment kept on site should be stored in a separate, lockable area ideally with a slop hopper and hand wash basin. If a separate area is not available then cleaning equipment may be located in dirty utility / sluice facilities. Ensure they are clearly segregated, where they are not at risk of environmental contamination. Under no circumstances should cleaning equipment be stored in clinical areas which are used for patient care, examination or treatment.

FREQUENCY OF CLEANING / CLEANING SCHEDULES

Environmental cleaning should be undertaken at a clearly defined frequency dependent on the risks associated with the specific environment. For example, clinical/treatment rooms require more frequent cleaning than office areas. (National specifications for cleanliness in the NHS: *Guidance on setting and measuring performance outcomes in primary care medical and dental premises* (NPSA 2010) both contain comprehensive guidance on cleaning frequencies and provide schedules for local modification and use. Cleaning schedules should be available for both domestic service staff and healthcare staff. Both different staff groups should be aware of who is responsible for which task. The schedule should clearly define which items require cleaning, who is responsible and what product is used. A log should accompany the schedule for staff to complete when the said task is completed. The monthly audit should give assurances that effective cleaning has taken place. Cleaning schedules should be available for public/service user inspection. This enhances public/service user confidence and is a requirement of the Hygiene Code of Practice.

STAFF TRAINING

It is essential that all housekeeping staff receive a fully documented induction and orientation programme including:

- Cleaning methods;
- Cleaning products and their safe use and storage;
- Use of appropriate protective clothing;
- Disposal of waste, including bagging, labelling and storage;
- Sharps safety;
- Cleaning of equipment, including care and storage;
- Personal and environmental COSHH safety;
- Hand Hygiene
- Food hygiene, if necessary;
- Incident/accident and illness reporting.

SPILLAGES OF BLOOD AND BODY FLUIDS

Blood and body fluid spillages must be dealt with immediately. In clinical areas this is usually a healthcare worker responsibility. In public access areas, e.g. corridors, lifts, public toilets, this is usually a domestic staff responsibility. However, in premises without domestic staff on site during working hours, this responsibility must be clearly defined. The registered provider should ensure that local staff are aware of their responsibilities which should be included in staff induction and infection control training.

Adequate and appropriate cleaning equipment, disinfectant preparations, protective clothing and clinical waste bags must be readily available. Floor signs indicating danger of slippage must be used where appropriate.

Spillages of blood and other high risk body fluids, e.g. faeces, should be dealt with immediately using a chlorine releasing agent e.g. sodium hypochlorite or one containing Na DCC (Sodium Dichloroisocyanurate). These are available as solutions and tablets (which require diluting to reach the correct concentration) or as powders and granules which contain an appropriate concentration. Powders and granules are available as spillage kits which often contain all the equipment required for the spill including yellow bags and card/scoop for removal of spill. Blood and body fluid spill kits are also available containing absorbent mats containing peracetic acid which can be used safely on all types of body fluid spillage but are not suitable for use on carpeted areas.

Urine and vomit spills should not be treated with chlorine-releasing products as these body substances are usually acidic (with a low pH) and can react with chlorine releasing noxious gases which may be inhaled (particularly in confined spaces such as toilets). Urine and vomit should be dealt with using method 3 unless a suitable spill kit is used.

Liquid preparations should be available in the correct concentration. A hypochlorite concentration of 10,000 ppm (parts per million) is necessary for use on blood and body fluid spillages. Preparations must be diluted immediately before use and any unused liquid must be discarded. Do NOT store reconstituted solution as it rapidly loses its efficacy.

WHY MANAGE BLOOD AND BODY FLUID SPILLAGES?

- Exposure to blood and other body fluids, such as faeces, vomit, pus and urine, poses a potential risk for transmission of infection to those providing care.
- Exposure to viruses such as HIV, hepatitis B and hepatitis C through blood or other body fluids can have severe consequences.
- Therefore, quick and effective management of spillages, regardless of the setting, is essential for the health and safety of all.

MANAGEMENT OF BLOOD AND BODY FLUID SPILLAGES

- All necessary equipment to deal with a spillage must first be gathered (see below)
- All items used during a spillage must be disposed of appropriately.
- Hand Hygiene must be performed following management of spillages.

- Control of Substances Hazardous to Health (COSHH) sheets and product data sheets should also be referred to so as to ensure safe management of spillages. Disinfectant must be used in accordance with manufacturers instructions, for storage, contact times and expiry dates refer to COSHH sheets.
- Wear PPE in accordance with manufacturers instructions for cleaning solutions/wipes and ensure adequate ventilation
- Wear disposable gloves, apron and eye protection when making up chlorine solutions and ensure adequate ventilation

Items to manage the spillage:

- Blood and/or body fluid spillage kit/or disposable absorbent paper towels and a solution of chlorine made up to 10,000ppm
- Disposable apron and gloves
- Face/eye protection if splashing is likely and/or cleaning with chlorine solution above waist height
- Clinical waste bag
- Wet floor signage as necessary

INCIDENT REPORTING

Consider local incident reporting of spillages to ensure future incidents or exposures to blood and other body fluids from spillages can be avoided and appropriate measures can be put in place.

Who should manage spillages?

- Those trained in the safe and effective management of blood and other body fluid spillages.
- All those working in health or social care who may be exposed to spillages of blood or other body fluids must receive training, e.g. through induction, information updates to ensure they are safe and effective in the management of such spillages. Training records must be held to reflect this.
- Responsibilities for the cleaning of blood and body fluid spillages should be clear within each area/care setting.

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

PEST CONTROL

There are a number of animals that can be considered pests within the health care setting and have the potential to cause disease or harm. These can range from mammals, such as cats, foxes, mice, rats and squirrels; insects such as ants, Pharaoh ants, cockroaches, beetles, wasps and spiders; parasites such as bedbugs, mites, lice and some birds, including pigeons.

Apart from the possibility of disease transmission, food may be tainted and spoiled, fabric and building structure damaged. Furthermore, Pharaoh's ants have been responsible for the penetration of sterile packs.

Pest control is a specialist problem, which requires immediate attention. The registered provider should have a contract in place for the routine management of pest control. Alternatively, the local council (pest control officer) may provide guidance.

Reporting and responsibilities

All staff sighting a pest within the healthcare practice should report the incident immediately by referring to the local protocol for pest control (which should be located with estates management policies). The information required will include:

- the location including, where possible the room number
- the type of pest if known
- the possible numbers and frequency of sighting
- the name of the person reporting
- If feasible, insects etc. can be captured and kept in a clean container, e.g. specimens' pots. It may be possible to take a picture using a digital camera for identification purposes.

If the infestation is noted in a clinical or food area, then it should not be used until further assessment and an appropriate inspection has been undertaken. Any food stuffs must be disposed of.

General control measures

Pests require somewhere to live, food, warmth and a means of entry.

Food needs to be kept covered and in rigid, impermeable containers and any spilt food must be cleared up as soon as possible.

Ensure that there are no areas of static water, such as puddles, either in the building or in the immediate grounds.

Do NOT feed pigeons, wild cats etc. with leftover food as this encourages pests and results in soilage from droppings.

Treatment with insecticides and rodenticides, by themselves, is rarely enough and it is essential that attention be paid to good general hygiene and structural maintenance.

Buildings should be well maintained, drains covered, damaged surfaces repaired, access holes sealed and leaking pipe work repaired. All of these can provide access to pests.

Close fitting windows and doors, fly screens and bird netting all help to reduce pest access.

All food preparation, storage and serving areas are subject to compliance with national and EU food hygiene legislation. Pest control in all food areas is subject to stringent controls under these regulations.

Waste storage areas should be well maintained and secure to minimise the likelihood of access by pests especially foxes, rats and pigeons. Clinical, household and food waste in particular will attract pests and should be stored off the ground in rigid, covered containers and, in the case of clinical / hazardous waste kept locked. Regular waste collections must be in place.

ESTATES AND FACILITIES MANAGEMENT

Increases in the incidence of Healthcare Associated Infections, and rising public concern, has highlighted the importance of appropriate management of healthcare environments.

Research has consistently shown that the environment can be a secondary reservoir for organisms with the potential for infecting patients. Good standards of basic hygiene, cleaning and regular planned maintenance can assist in preventing healthcare associated infections. This is more easily achieved if the built environment supports best practice.

The Code of Practice requires organisations delivering care to “provide and maintain a clean and appropriate environment in managed premises that facilitate the prevention and control of infection”. Criterion 2 of the guidance states:-

“Premises and facilities should be provided in accordance with best practice guidance. The development of local policies should take account of infection prevention and control advice given by relevant expert or advisory bodies or by the ICT, and this should include provision for liaison between the members of any ICT and the persons with overall responsibility for the management of the service user’s environment. Policies should address but not be restricted to:-

- Cleaning services
- Building and refurbishment, including air handling system
- Waste management
- Laundry arrangements for used and infected linen
- Planned preventative maintenance
- Pest control
- Management of drinkable (potable) and non-drinkable (non-potable) water supplies
- Minimising the risk of Legionella by adhering to national guidance
- Food services, including food hygiene and food brought into the care setting by service users, staff and visitors.”

Information and guidance is provided in sections of this Manual on some of these matters. This section offers guidance on the built environment (build and refurbishment works); planned preventative maintenance and water safety.

Local policies concerning waste management, food hygiene, environmental cleaning, provision of laundry and other facilities matters must include the requirement for liaison with Infection Prevention & Control specialists when service arrangements are made or changed.

Building & Refurbishment works

Technical guidance is produced by the Department of Health for healthcare building projects covering a range of health and social care provision. These include Health Building Notes (HBNs) and Health Technical Memoranda (HTMs). A key document covering IPC aspects of buildings is HBN 00-09, *Infection Control in the Built Environment* (2013). Planning guidance for primary and community care facilities is also available (see Bibliography)

When planning builds or refurbishment of primary care facilities, or when planning additional clinical services, the appropriate guidance must be consulted and IPC advice sought.

Areas or rooms where clinical activities are to be undertaken (e.g. wound dressings, insertion of urinary catheters, or other invasive procedures) or where medical consumables are to be stored should incorporate IPC requirements. If the disposal of blood and body fluids is likely to be undertaken in primary care then a dirty utility facility (sluice), is required. Consideration should be given to providing a suitable area for the testing and disposal of urine samples. Samples should never be disposed of into a hand wash sink.

Carpets are not acceptable in areas where clinical procedures are undertaken.

If enhanced services are provided consideration must be given to the suitability of the environment in which these will be conducted. Local Commissioners policies detailing this must be sourced and followed. If no such policy is available advice should be sought from IPC specialists. Details of requirements will depend on what type, or levels of procedures are to be undertaken. Minimal Access Interventions (MAIs) and ophthalmic procedures for example required mechanical ventilation systems whilst minor procedures can be conducted in a naturally ventilated room. Windows, if opened, must be protected with mesh screens.

Clinical services may be provided in sites managed by other organisations. In such situations the organisation must assure itself that the environment is appropriate for the care being delivered and is managed in accordance with the principles outlined in this policy and with published guidance.

Water Safety

Legionella spp. which causes Legionnaires' disease is found naturally in water supplies. If appropriate control measures are not in place, the bacterium may multiply to a pathogenic level and outbreaks may follow. HTM 2040 and the Health & Safety Commission Approved Code of Practice (L8) give detail on the required management arrangements to reduce this risk. Processes should include routine, and repeated, risk assessment and the adoption of advice from suitably qualified specialists. There should be local policies detailing this. As stated above, if the facility is managed by a host organisation, the practice should seek assurance that water safety is appropriately managed.

Legionella risks increase where water outlets are used infrequently and thus Legionella can multiply. Staff should monitor the use of water outlets, all staff should report infrequently used outlets i.e. those not used daily, and these should be documented. Identified low use outlets should be subject to regular (usually weekly) flushing regimes. These should also be routinely documented using a simple log.

Guidance has recently been issued (March 2012 and 2013) on reducing risks of Pseudomonas infections from tap water. This is of particular relevance to Augmented Care Units (renal, burns, critical care, haematology, neonatal units) however all health care providers are asked to assess the risks to their patient groups. Advice should be sought from the water advisor or from the organisation managing the facility as to whether a formal Water Safety Group is required and established and what measures practice staff should take to protect patients.

Planned Preventative Maintenance (PPM)

Most equipment used in health and social care carries PPM requirements as recommended by manufacturers. Good equipment management can prolong the life of the equipment, prevent costly breakdown, and ensure the equipment is fit for purpose. Failure of some equipment in healthcare may pose IPC risks. This would include, but is not limited to:-

- Bed Pan Washers/macerators
- Laundry equipment e.g. washers/dryers
- Vaccine/specimen Fridges
- Catering equipment e.g. fridges/dishwashers

Policies or processes should be in place to ensure this equipment is maintained in line with manufacturer's instructions and this maintenance should be documented.

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

DECONTAMINATION OF MEDICAL EQUIPMENT

INTRODUCTION

Decontamination requires the implementation of a number of processes, from purchasing equipment through to delivery and use, cleaning and disinfecting, packing, sterilising, repair and disposal. To be effective it needs standards to be set for all elements of the device life cycle.

DEFINITION OF A MEDICAL DEVICE

A medical device is any instrument, apparatus, appliance, material or other article used alone or in combination and intended by the manufacturer to be used for humans for any of the following purposes:

- Control of conception
- Monitoring, diagnosis and investigation
- Treatment, alleviation or compensation for injury or incapacity
- Replacement or modification of anatomy and physiology

The vast majority of equipment used in patient care will be defined as a medical device.

DUTIES

It is the responsibility of healthcare staff to ensure that all medical devices used in patient care are appropriately decontaminated and fit for purpose. Duties of key personnel are clearly defined in HTM 01-01 (2016).

DECONTAMINATION OF RE-USABLE MEDICAL DEVICES

The re-processing of medical devices required to be sterilised prior to re-use (either at point of use or prior to storage) is subject to stringent process controls. Since 2007 there has been a requirement for all such devices e.g. surgical instruments to be re-processed in a registered facility. It is no longer acceptable for local re-processing to be undertaken in any provider service *with the current exception of primary dental decontamination which is subject to the requirements of HTM 01-05.*

Primary care practices have a duty to ensure that re-usable medical devices required to be sterilized have arrangements in place to ensure that they:

- Use a decontamination service registered with MHRA who are compliant with the Medical Device Regulations (2002) and who use a Notified Body as their third party auditor or:
- Use a decontamination service which is subject to Care Quality Commission (CQC) audit or:
- Use CE marked single use medical devices or:
- Employ a strategy featuring a combination of the above

RISK ASSESSMENT

Medical equipment is categorised according to the risk that particular procedures pose to patients and by assessing the microbial status of the body area being manipulated during the procedure. For example, items that come into contact with intact mucous membranes are classified as intermediate risk and require disinfection between each use as a minimum standard. Items that enter normally sterile body areas, or items that come into contact with broken mucous membranes, are classified as high risk and must be sterile before use.

Risk Assessment for Decontamination of Equipment

Risk	Application of Item	Minimum Standard
Low	<ul style="list-style-type: none"> In contact with healthy skin e.g. furniture, mattresses, surfaces, or no contact 	Clean
Intermediate	<ul style="list-style-type: none"> In contact with intact mucous membranes Contaminated with virulent or readily transmissible organisms (body fluids) e.g. commode pans / bed pans For use on immuno-compromised patients 	Disinfect, or single use
High	In contact with broken skin or mucous membrane <ul style="list-style-type: none"> For introduction into sterile body areas 	Sterilize, or single use

Adapted from Medical Devices Agency, Part 2 (1996) now MHRA

DECONTAMINATION - DEFINITIONS

Decontamination of Re-Usable Devices

Decontamination is the term widely used to collectively describe the combination of processes of cleaning, disinfection and sterilisation to make a re-usable device safe for further use on patients and safe for the user. The effective decontamination of re-usable devices is essential to reduce these infection risks. Decontamination methods used will depend on the nature of the micro-organisms present and the infection risk associated with the surface, equipment, device or procedure.

Cleaning

Cleaning is a process which physically removes contamination but does not necessarily destroy micro-organisms. The reduction of microbial contamination will depend upon many factors including the efficiency of the cleaning process and the initial contamination. A further reduction will occur on drying, as some micro-organisms cannot multiply on a clean dry surface. Cleaning is the first step in the decontamination process. It must be carried out before disinfection and sterilisation to make these processes effective. Thorough cleaning is extremely important in

reducing the possible transmission of all micro-organisms, including the prion protein that causes vCJD.

Liquid detergent and warm water is an effective cleaning agent. Hot water should not be used, as it will coagulate proteins (body fluids) making it more difficult to remove from the equipment. Hard surface detergent wipes are also available for equipment cleaning.

Disinfection

Disinfection is defined as a process used to kill or remove pathogenic micro-organisms but which cannot usually kill bacterial spores (Lawrence and May, 2003). Disinfection processes, if used appropriately will reduce micro-organism counts to safe levels.

Disinfection processes can be used on both equipment and environmental surfaces and usually involves the use of either a disinfectant solution or a structured process using equipment such as bed-pan washer-disinfectors, dishwashers and washing machines where temperature of water or steam provides the disinfection process.

Disinfection using antiseptic solutions is the process used to reduce microbial contamination of the skin, mucous membranes and other body tissues and cavities.

Sterilisation

Sterilisation is a range of processes used to render the device free from viable micro-organisms, including spores. Processes include moist and dry heat using autoclaves and / or hot air ovens; low temperature steam and formaldehyde; ethylene oxide and gas plasma.

In healthcare, sterilisation processes are usually confined to the application of moist heat using autoclaves.

DECONTAMINATION ADVICE FOR HEALTHCARE STAFF

If the method of decontamination is in doubt, then advice may be sought from:

- The device supplier and/or manufacturer of the equipment
- Local decontamination lead and / or infection control lead

CLEANING AND DECONTAMINATION SOLUTIONS

The following products are suitable for the decontamination of the majority of health care equipment and surfaces. Specialised equipment should be decontaminated following manufacturers' instructions.

Neutral detergent (washing-up liquid) – a mild detergent that is adequate for most cleaning of equipment and surfaces and will mechanically remove (by cleaning) the majority of micro-organisms contaminating equipment. Refer to bottle before use, but usually 5ml in 1 litre of warm water is sufficient.

Hard surface wipes – There are two main types of wipe. Some contain 70% alcohol or other disinfectants and others contain a detergent. Wipes are cheap and effective and are portable (in drums/packs) and require no access to water. They can also be used on large items of equipment. Detergent wipes can be used instead of detergent and water. Alcohol wipes can be used for surfaces requiring disinfection as well as cleaning e.g. dressing trolleys. Where gross soilage/contamination is present a detergent wipe is preferable as alcohol is inactive in the presence of soil. When using alcohol wipes it is important to remember that the item must be cleaned with a detergent based product first.

Chlorine-releasing agents – These contain a chlorine-releasing agent and are often referred to as bleach solutions or hypochlorite. They are used for spillages of blood and high risk body fluids such as faeces and can be used to disinfect service user contact surfaces in an isolation room and also during outbreaks of infection diarrhoea and vomiting for cleaning of both the environment and equipment such as commodes. Staff using such products must be familiar with COSHH regulations. Aprons, gloves and facial protection must be worn for preparation and use. The product must be mixed in a well ventilated area. Refer to bottle for correct dilution.

Non-abrasive cream cleaner – Mild cleaner for general hard surface use e.g. sinks

Toilet cleanser/sanitizer – cleanser which can contain bleach and/or lime-scale remover

Thermal washer/disinfector e.g. bedpan washer/dishwasher – designated machinery for thermal (heat) disinfection of articles where a higher temperature and controlled method of cleaning are required e.g. bedpans and / or cutlery / crockery.

Decontamination of equipment prior to loan, servicing or repair

It is the responsibility of the person/department using the equipment to ensure that it is visibly clean and free of surface contamination with blood and/or body fluids if being sent for service, maintenance or repair either on or off site (HSG(93)26).

A decontamination notice must be attached to the equipment to warn others of the type of contamination it may have been exposed to and whether it has been possible to decontaminate it. Many manufacturers provide their own decontamination certificates with their equipment and will not accept returned equipment without an accompanying certificate. This is appropriate practice and should also apply to equipment being repaired or serviced on-site.

MEDICAL DEVICE CLEANING PROCESSES – GUIDANCE NOTES

- Comprehensive guidance on cleaning of both medical devices and other patient specific equipment is available in two publications - National specifications for cleanliness in the NHS: a framework for setting and measuring performance outcomes (NPSA 2007) and revised healthcare cleaning manual (NPSA 2009)
- Cleaning of medical devices and other patient specific equipment should be subject to regular, on-going monitoring of the standard of cleaning

- Cleaning schedules specifying the frequency of cleaning should be devised incorporating all medical devices / equipment used locally. These schedules should be available for all staff and a simple check-list should be devised for staff to sign after completion of cleaning
- Re-usable medical devices must be decontaminated between each patient use. Some larger items of equipment e.g. notes trolleys etc. should be cleaned weekly. Frequency of cleaning is specified in the documents above.
- The user of the device is responsible for ensuring that it is visibly clean and free from contamination with blood and/or body fluids following each procedure or care episode and prior to sending for service or repair internally and externally.
- Dirty equipment awaiting cleaning, should be stored separate from clean items and should be cleaned as soon as possible after use and then stored appropriately
- Cleaning of equipment should take place in a designated area e.g. dirty utility or away from clean items that could become contaminated during the cleaning process
- Personal protective equipment (PPE) should be worn when cleaning medical devices. Disposable gloves (or household gloves) together with a plastic apron should be worn to protect hands and clothing. Please note that facial protection is required when using some products; please refer to manufacturer's guidance.

A to Z and decontamination of patient equipment and medical devices

Device	Cleaning method	Method for contaminated/infected equipment	Frequency
Airways	Single use only	N/A	N/A
Arm rests	Wash with neutral detergent, rinse and dry	Wash with chlorine based product or approved disinfectant, rinse and dry	On discharge or if visibly soiled
Audiometer headphones	Wipe with neutral detergent and dry		After each use
Auroscope and ear pieces	Wash ear pieces with neutral detergent. Wipe scope with detergent wipe	Consider disposal or wash with chlorine approved disinfectant Ensure ear piece is free of wax, particularly down the lumen. Do not immerse the auroscope.	After each use
Baby bottles and teats	Single use only or use a chlorine releasing solution (e.g. Milton)	N/A	After each use
Baby changing mats	Wipe with neutral detergent, rinse and dry.	Wash with chlorine based product or approved disinfectant	After each use
Baby Scales	Wipe with neutral detergent, rinse and dry	Wash with chlorine based product or approved disinfectant rinse and dry	After each use
Blood pressure cuffs	Clean with a detergent wipe	Single patient use cuff for infectious patients	After each use
Blood pressure machine	Clean with detergent wipe	If not able to use dedicated equipment for infectious patients, clean after each use with chlorine based product or appropriate detergent	
Breast pumps	Single patient use accessories, these should be washed in neutral detergent, rinsed and then immersed in a chlorine releasing agent (e.g. Milton). Pump should be cleaned with neutral detergent and dried	Disposable	After each use
Crockery and cutlery	Dishwasher or hand wash using hot water and detergent	Dishwasher only	After each use.
Dressing trolley	Clean with neutral detergent and dry. Clean from top to bottom using an 'S' shape motion.		Before and after each use
Drip stands	Clean with neutral detergent and dry	Wash with chlorine based product or appropriate disinfectant, rinse and dry	Weekly, after discharge or when visibly dirty.

Device	Cleaning method	Method for contaminated/infected equipment	Frequency
Electronic devices e.g. Ipad	Clean with a disinfectant wipe, or use commercial wipes for electronic equipment. Keyboards – cover with wipeable cover or use soap and water. Dry with a paper towel.		After each use
Examination couch	Clean with neutral detergent, rinse and dry. Use paper couch roll and change between each patient use.	Wash with chlorine based product or appropriate disinfectant, rinse and dry	Daily or if visibly dirty
Fans	Clean with neutral detergent.	Wipe with chlorine based product or appropriate disinfectant rinse and dry	After each use or if visibly dirty
Furniture and fittings	Damp dust with detergent solution	Clean where possible with chlorine based product	
Hair brushes	Single patient use	N/A	N/A
Medicine Pots	Single use	N/A	N/A
Medicine trolley/cupboard	Clean with neutral detergent, rinse and dry		Weekly or if visibly soiled
Mops	Disposable mops for ward use.	N/A	After each use
Nail brushes	Single use	N/A	N/A
Nebuliser compressor	Clean with detergent wipe		Weekly, following discharge or if visibly soiled
Nebuliser pots	Clean with hot water. Change weekly	N/A	After each use
Oxygen masks	Single patient use.	N/A	Weekly or if soiled
PCs, printers and keyboards	Wipe with detergent wipe		Weekly
Peak flow meters	Wipe with detergent wipe. Mouth pieces single use.	Wipe with disinfectant wipe	After each use
Pillows	See mattresses'		
Pulse oximeter units	Wipe with detergent wipe	Wipe with chlorine based product or approved disinfectant	After each use
Pulse oximeter probes	Wipe with detergent wipe	Wipe with chlorine based product or approved disinfectant	After each use
Shaving brushes	Patients own. Not to be used for clinical shaving	N/A	N/A
Resuscitation trolley	Clean with neutral detergent, rinse and dry.	Clean with chlorine based product or appropriate disinfectant rinse and dry	Weekly or if visibly soiled
Sputum container	Disposable only	N/A	N/A
Stethoscopes	Wipe with detergent wipe	Wipe with chlorine based product or approved disinfectant	After each use

Device	Cleaning method	Method for contaminated/infected equipment	Frequency
Suction equipment	Clean and dry containers with neutral detergent. Use disposable liners	Clean with Wipe with chlorine based product or approved disinfectant or approved disinfectant rinse and dry	On discharge or if visibly soiled
Syringe drivers	Wipe with detergent wipe	Clean with disinfectant wipe	On discharge or if visibly soiled
Therapy equipment	Wipe with detergent wipe	Wipe with Wipe with chlorine based product or approved disinfectant or approved disinfectant	After each use
Thermometers	Use disposable probe cover Wipe monitor with detergent wipe	Clean monitor with disinfectant wipe	After each use
Tympanic thermometers	Use disposable ear piece Wipe monitor with detergent wipe	Clean monitor with disinfectant wipe	After each use
Toys and play equipment	Wash with neutral detergent, rinse and dry. Do not soak soft toys.	Clean with Wipe with chlorine based product or approved disinfectant or approved disinfectant. Heavily contaminated soft toys may have to be destroyed.	After each use
Walking aids	Wash with neutral detergent, rinse and dry	Clean with Wipe with chlorine based product or approved disinfectant or appropriate disinfectant, rinse and dry	After each use
Wheel chairs	Clean with neutral detergent	Clean with with chlorine based product or approved disinfectant rinse and dry or use disinfectant wipe	After each use
Weighing scales	Clean with neutral detergent or use detergent wipe	Clean with Wipe with chlorine based product or approved disinfectant , rinse and dry or use disinfectant wipe	After each use

This list contains common use equipment only and is not exhaustive.

EQUIPMENT	RECOMMENDED DECONTAMINATION PROCEDURE
Auriscopes ear pieces	Wash with neutral detergent and dry then wipe with 70% alcohol wipe and air dry
Baby changing mats	Always replace mat if ripped or damaged Protect with disposable paper and change after each use. Clean mat at the end of the session or when contaminated with neutral detergent and hot water Follow with a hard surface disinfectant wipe if contaminated with blood or body fluids Protect mat during use with disposable paper roll
Bed pans, commode pans, urinals	Disposable or Decontaminate in washer-disinfector
Bed pan shells (holders for disposable bed pans)	Wash in warm detergent and water, rinse and dry with paper towels
Buckets (used to soak dressings)	Ideally use disposable liner and change after each patient. Always wash after removal of liner. Wash with neutral detergent and warm water, rinse and dry thoroughly. Store inverted and separated.
Computer keyboards	Cover with wipeable cover. Clean with detergent and warm water. Dry with paper towels. Alternatively use commercial wipes for electronic equipment
Pillows	Cover with impermeable cover and decontaminate using detergent and water. Dry thoroughly.

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

SINGLE USE AND SINGLE PATIENT USE MEDICAL DEVICES

Single use medical devices are manufactured to be used on a single occasion and then discarded. They are not designed or manufactured for re-use even on the same service user. The re-use of single use devices is dangerous and has legal implications under the Medical Devices Regulations (2002) and Medical Devices (Amendment) Regulations (2008).

The Medicines and Healthcare products Regulatory Agency (MHRA) guidance 2018 – *DB2006 (04) Single-use Medical Devices: Implications and Consequences of Reuse*.

The MHRA state that “to reuse a single-use medical device without considering the consequences could expose the patient and staff to risks which outweigh the perceived benefits of using the devices”.

The MHRA advises against the reuse of any single-use medical device. MHRA Key points:

- A device designated for ‘single-use’ must not be reused. It should only be used on an individual patient during a single procedure and then discarded. It is not intended to be reprocessed and used again, even on the same patient
- The reuse of a single-use device can affect their safety, performance and effectiveness, exposing patients and staff to unnecessary risk.
- The reuse of single-use devices has legal implications:
 - Anyone who reprocesses or reuses a device intended by the manufacturer for use on a single occasion, bears full responsibility for its safety and effectiveness;
 - Anyone who reprocesses a single-use device and passes it to a separate legal entity for use has the same legal obligations under the Medical Device Regulations as the original manufacturer of the device.

Manufacturers are required to clearly identify single-use devices by displaying a “do not reprocess” symbol as shown below.

Figure 1: “Do not reprocess” symbol



TECHNICAL ISSUES

Reprocessing single-use devices may affect the capabilities and/or the materials from which the device is made.

Many single use devices are unable to withstand the decontamination and sterilization processes used in health care.

The manufacturer will provide a warranty for a medical device made for reuse if the recommended reprocessing is carried out. If a single use item is reprocessed, the manufacturer's warranty will not apply and the re-processor will take on this responsibility

PROBLEMS ASSOCIATED WITH RE-PROCESSING

Inadequate cleaning and decontamination - the cleaning process must be able to access all parts of the device to enable complete decontamination, the cleaning agents must be completely removed at the end of the process and this process must be validated by the processor. Many single-use devices have inaccessible angles and narrow lumens making cleaning and validation impossible

Material alteration - Exposure to chemicals and other processes may cause corrosion or alteration of the device materials making it unsafe to use e.g. plastics may become brittle and break during subsequent use.

Mechanical failure - Some devices if repeatedly processed may over time become stressed and fail or break in use e.g. single-use drills, burrs and blades, etc.

Potential for cross infection – Cross infection is a major risk associated with the re-use of single-use items due to failure to clean, decontaminate, disinfect or sterilise adequately.

Reactions to endotoxins - These are residues of bacteria which withstand exposure to heat and chemicals and may remain after re-processing and sterilization. The sterilization process may not inactivate the toxins even when cleaning and sterilization is effective in killing the bacteria.

Residues from chemical decontamination agents - Some materials used in the device's manufacture may absorb the chemicals used in the decontamination process resulting in chemical burns or sensitization of the patient.

Reprocessing a medical device designed or designated as single use requires the device to undergo an extensive validation process to ensure that it is safe to reuse. The majority of organisations do not have the finances or the facilities to carry out this process as the re-use of these devices is likely to carry a significant risk.

Prion disease (inc. CJD)

The abnormal proteins associated with prion diseases are highly resistant to conventional methods of decontamination and sterilisation. It is therefore an even greater risk to reprocess equipment that may have been exposed to patients known or suspected of being infected with

this agent. (See section – Infections with Specific Alert Mechanisms for further information).

CONCLUSION

To re-use a single-use device without considering the consequences to the organisation, the professional and the patient could expose each or all of these individuals to significant levels of risk both personal and financial.

MEDICINES

Medicines, including topical medical products must be considered as single use items unless the label and / or supporting manufacturers' guidelines clearly state they the item has been prepared as a multi-dose item.

A risk assessment must be carried out (in conjunction with Medicines Management) for each individual product.

THE USE AND RE-PROCESSING OF SINGLE PATIENT USE DEVICES

There are a number of medical devices that are manufactured for limited re-use by the individual to whom they are initially supplied. The majority of these devices are non-invasive and do not require sophisticated reprocessing to ensure they are safe for re-use.

It is essential that when these devices are re-used there are written manufacturer's guidelines available for their use, cleaning, decontamination and disposal. All staff should have access to manufacturer's guidelines which must be retained in a suitable folder / location.

Professional staff who use or supply these devices to patients must understand the requirements for safe use, decontamination between uses and disposal.

TYPES OF SINGLE PATIENT USE DEVICES

Patient self-administered intermittent urinary catheters

These are issued to an individual patient for their own use. They should be washed under running water after each use and stored clean and dry. Each catheter should be replaced according to the manufacturer's instructions or at least once a week, sooner if damaged. If used by a healthcare professional on behalf of the patient they must be treated as single use items and disposed of after a single use.

Face masks for oxygen administration

These items should be kept with the individual patient, particularly if the oxygen cylinder is shared. The facemask should be washed daily, and if soiled, with warm water and detergent, dried and stored dry. The mask should be replaced weekly. Tubing must also be single patient use, changed if wet and replaced weekly.

Feeding syringes for patient with well established PEG feeding tubes

Specific oral syringes are manufactured for use with PEG feeding tubes e.g. Baxa syringes. They are supplied as a clean not sterile product. Manufacturers' guidelines for re-use must be followed. Alternatively, they should be thoroughly cleaned after each use with warm water and detergent, rinsed in running water, shaken to remove water particles from the barrel of the syringe and dried externally with disposable paper towels prior to storing in a dry, covered container e.g. plastic food container with lid. These are for use by an individual patient, and must be replaced daily (or in accordance with manufacturers' instructions). Please note that single use syringes must not be re-used, even on the same patient.

Newly inserted PEG feeding tubes are classed as surgical wounds and thus feeding syringes should be used once only and discarded after single use until such times as the stoma is healed.

Nebulisers

These items should be kept with the individual patient. The nebuliser should be rinsed after each use with warm water ONLY (no detergent), shaken to remove water particles and drug residues and then dried with disposable paper towels and stored dry in a clean, dry, covered container. The nebuliser should be replaced weekly provided it maintains its efficacy or as per manufacturer's instructions. A label can be attached to the storage container indicating the date for change. Masks (if used) should be decontaminated as above.

Placebo inhalers (for prescribed inhaled therapy)

Currently there is no evidence upon which to base local protocols for decontamination of these devices, when a mouthpiece cannot be used, therefore manufacturer's instructions must be followed. Ideally, devices that can be fitted with a disposable mouthpiece should be used.

Where these products are in use, guidance should be sought from the local respiratory nurse or clinician who prescribed the device. As a minimum, patients with a known or suspected respiratory infection should not use communal inhalers.

Appropriate methods of decontamination (in the absence of manufacturer's guidance) include: thorough washing with liquid detergent and warm water followed by shaking to remove water particles and drying with paper towels. In addition, disinfecting in a freshly prepared solution of sodium hypochlorite (1,000 ppm) followed by rinsing under running water, shaking to remove water particles and then drying with paper towels can be undertaken after initial washing in detergent.

Other items

There may be other items that can be designated single patient use. Each of these must have written guidelines for use, decontamination and frequency of replacement, preferably supplied by the manufacturer.

VACCINE MANAGEMENT GUIDELINES

Vaccines are biological substances that may lose their effectiveness quickly if they become too hot or cold at any time, especially during transport and storage. Vaccines naturally biodegrade over time and storage outside of the recommended temperature range – including during transport and storage – may speed up loss of potency, which cannot be reversed. This may result in the failure of the vaccine to protect, as well as resulting in vaccine wastage.

It is essential that all those handling vaccines follow local policies to ensure cold chain compliance. This guideline is not a substitute for local policy (see below) but provides key information relating to the management of the vaccine cold chain which is a key element of local immunization policy. This guideline has been written incorporating the guidance provided in: *Immunization against infectious diseases (the Green Book) chapter 3 – storage, distribution and disposal of vaccines Department of Health (2013)*

POLICIES AND PROCEDURES IN PRIMARY CARE

Local immunisation policies are developed by key professionals involved in Medicines Management and Public Health, as well as clinicians, commissioners and Immunisation Co-ordinators at Public Health England. Primary care organisations should ensure that local practice is in accordance with national policy (as per reference above) for the ordering, storage, stock control, distribution, transport and disposal of vaccines. Details of local contacts for professional advice should be clearly provided in local policies.

Each practice should have one trained individual with at least one trained deputy, responsible for the receipt and storage of vaccines and the recording of refrigerator temperatures. Regular audit (at least annually) should be undertaken. Any other staff who may be involved with vaccines must be trained appropriately.

THE IMPORTANCE OF THE COLD CHAIN

The cold chain is standard practice for vaccines throughout the pharmaceutical industry. The cold chain commences with the vaccine manufacturer and continues through specialist pharmaceutical distribution companies to the Pharmacy / GP surgery and finally to the vaccine recipient. It will also incorporate transportation in refrigerated vans and / or cool boxes and the use of alternative refrigerators in case of failure. In other words the cold chain **MUST** be maintained throughout the life of the vaccine.

Maintaining the cold chain ensure that vaccines are transported and stored according to the manufacturer's recommended temperature range of +2°C to +8°C until the point of administration.

STORAGE

Vaccines should be stored in the original packaging at +2°C to +8°C and protected from light, as exposure to ultraviolet light will cause loss of potency. All vaccines are sensitive to some extent to heat and cold. Heat speeds up decline in potency of most vaccines, thus reducing their shelf life. Effectiveness cannot be guaranteed for vaccines unless they have been stored at the correct

temperature. Freezing may cause increased reactogenicity and loss of potency for some vaccines. It can also cause hairline cracks in the container, leading to contamination of the contents.

ORDERING AND MONITORING OF STOCK

Care must be taken in ordering vaccines, especially as certain vaccines are packaged in multiple quantities. Incorrect ordering can result in wastage and unnecessary costs. Vaccine stocks should be monitored by the Designated Person(s) to avoid over-ordering or stockpiling.

Generally surgeries should have no more than two to four weeks supply of vaccines at any one time. This will be sufficient for routine provision. Best practice is to order small quantities on a regular, scheduled basis.

Excess stock may:

- Increase the risk of vaccination with out-of-date vaccines
- Increase wastage and the cost of disposal by incineration
- Increase the dangers of over-packed refrigerators, leading to poor air flow, potential freezing and poor stock rotation
- Delay the introduction of new vaccines until local supplies have been used
- Increase the cost of replacement of stocks if the refrigerator fails
- Increase the pressure on clinic refrigerators in periods of high demand e.g. during the influenza vaccination season

Vaccine stocks should be placed within the refrigerator so that those with shorter expiry dates are used first.

Any out-of-date stock should be labelled clearly, removed from the refrigerator and destroyed as soon as possible according to the local procedure.

Vaccines must never be used when past their expiry date

RECEIPT OF VACCINES

On receipt of vaccines, staff must check them against the order for discrepancies and leakage or damage before signing for them. Pharmaceutical distributors and manufacturers will not accept any vaccine for return once it has left their control.

Vaccines must be refrigerated immediately on receipt and must not be left at room temperature.

Vaccine types, brands, quantities, batch numbers and expiry dates should be recorded with the date and time at which the vaccines were received.

THE VACCINE REFRIGERATOR

Specialised refrigerators are available for the storage of pharmaceutical products and must be used for vaccines and diluents. Ordinary domestic refrigerators must NOT be used. Food, drink and clinical specimens must NEVER be stored in the same refrigerator as vaccines. Opening of the refrigerator door should be kept to a minimum in order to maintain a constant temperature.

All vaccines are Prescription Only Medicines (POMs) and must be stored under locked conditions. Refrigerators must either be lockable or within a room that is locked when not occupied by a member of staff. Vaccines should never be left unattended at outlying clinics.

The accidental interruption of the electricity supply can be prevented by using a switchless socket or by placing cautionary notices on plugs and sockets.

Refrigerators should not be situated near a radiator or any other heat source that could affect their working and should be appropriately ventilated.

Ice should not be allowed to build up within the refrigerator as this reduces effectiveness. Records of regular servicing, defrosting and cleaning should be kept.

An approved cool box (with appropriate temperature monitoring (see below) or an alternative refrigerator should be available and used to store vaccines during defrosting of the main refrigerator (and in the event of refrigerator failure). Vaccines should only be replaced once the refrigerator has returned to the correct temperature after defrosting.

REFRIGERATOR THERMOMETERS

The temperature within the vaccine refrigerator must be continually monitored with a maximum-minimum thermometer. This will identify when the temperature may have been outside the recommended range. Digital thermometers are the most reliable. More sophisticated temperature-recording devices are now available, including alarmed digital maximum-minimum thermometers.

Thermometers should be reset and replaced according to the manufacturers' guidance.

Temperatures in the refrigerator must be monitored and recorded at least once each working day, and documented on a chart for recording temperatures (see Appendix). The calibration of thermometers should be checked annually to ensure that they are working correctly. This will require a contract to be in place for Planned, Preventive Maintenance (PPM). The records should be readily accessible for easy reference and during annual audit. Records should ideally be retained for two years after last entry.

REFRIGERATOR FAILURE OR DISRUPTION OF THE COLD CHAIN

Local arrangements should be in place for back-up facilities to be available in the event of the refrigerator failing or breaking down. This should include the following:

- Record temperature fluctuations outside of normal range on temperature record
- Estimate the time period the refrigerator has been out of range / failing
- Contact local personnel for remedial action in accordance with local immunisation policy
- Contact your pharmacist or a member of the medicines management team and report the problem and estimated time period
- Removal of vaccine in the refrigerator and storage in another refrigerator or cool box *that is within correct temperature parameters*
- Quarantine the stock and label “not for clinical use” until authorization to use has been given by your pharmacist / medicines management team
- Complete an Incident Report in line with local arrangements

STORAGE OF VACCINES

Vaccines must be kept in their original packaging when stored, so that they retain information on batch numbers and expiry dates. The packaging is also part of the protection against light and changes in temperature.

Vaccines must NOT be stored in the door, in the bottom drawers or adjacent to the freezer plate of the refrigerator. If there are temperature variations outside of the recommended range (+2°C to +8°C), they usually occur in these parts of the refrigerator. Sufficient space should be allowed in the refrigerator so that air can circulate freely.

Patients, parents and carers should not normally be asked to store vaccines. Exceptionally, they may be asked to transport vaccines and / or immunoglobulin and to store them for short periods of time. Should this need arise, there should be a local written protocol on appropriate storage that is given to the patient / parent / carer *which must be read and understood*.

PACKAGING AND TRANSPORTING OF VACCINES TO OUTLYING CLINICS / SITES

Domestic cool boxes should not be used to store, distribute or transport vaccines. Validated cool boxes (with maximum – minimum thermometers) and ice packs from a recognised medical supply company should be used. Individual manufacturers’ instructions should be strictly adhered to. These are usually validated for use up to 8 hours after packing, provided the manufacturer’s instructions have been followed and it essential that staff are knowledgeable about specific manufacturer’s instructions.

Vaccines must be kept in the original packaging, wrapped in bubble wrap (or similar insulation material) and placed into a cool box with cool packs as recommended by the manufacturer (of the cool box). This will prevent further contact between the vaccine and the cool packs and will protect the vaccine from any damage, such as being frozen.

Avoid exposure to high temperatures during transit and at vaccination clinic / site. Do not place cool box near any areas which may have high temperatures e.g. radiator, in full sunshine.

Always keep vaccine cool box within sight at all times when it is off premises and in transit / use elsewhere.

STOCK CONTROL IN TRANSPORTED VACCINES

Those vaccines that have been returned to the refrigerator from a previous session from the cool box should be used first. These should have been marked to indicate that they need to be used first.

It is important to take only the required amount of vaccines to the session, together with reasonable additional contingency stock.

At each immunisation session, ensure the minimum number of vaccine containers are removed for each vaccinator to ensure no wastage takes place. Ensure that the cool box is fully closed following each opening.

Vaccines removed from the cool box and not used must be discarded.

AT THE END OF A VACCINATION SESSION

Mark vaccines that have been returned to the refrigerator from a previous session from the cool box to indicate that they should be used first at the next vaccination session.

Avoid exposure to high temperatures during return transit.

Go straight back to the site of the storage refrigerator and replace vaccines back into the refrigerator immediately on arrival.

DISPOSAL OF VACCINES

There should be locally written procedures for the disposal of vaccines by incineration at a suitably authorised facility. These procedures must be followed.

All reconstituted vaccines and opened single or multi-dose vials must be used within the period recommended by the manufacturers or should be disposed of at the end of the immunisation session by sealing in a proper, puncture-resistant sharps box (UN approved BS 7320)

The sharps container should be replaced once the contents have reached the 'fill' line and should not be accessible to any unauthorised individual.

Out of date stocks of vaccines should be returned to the manufacturer for disposal.

STORAGE OF IMMUNOGLOBULINS

Immunoglobulins should be protected from light and stored at +2°C to +8°C. Although these products have a tolerance to ambient temperatures (25°C) of up to one week, they should be refrigerated immediately upon receipt. They can be distributed in sturdy packaging outside the cold chain if needed. They should not be frozen.

SPILLAGE

Locally written procedures should be used in conjunction with manufacturers' Control of Substances Hazardous to Health (COSHH) safety data sheets.

Spillages must be cleared up quickly and gloves should be worn. The spillage should be soaked up with paper towels, taking care to avoid skin puncture from glass or needles. The area should be cleaned according to the local chemical disinfection policy or COSHH data sheets. Gloves, towels etc. should be discarded at clinical waste for incineration.

Spillages on skin should be washed with soap and water. If a vaccine is splashed in the eyes, they should be washed with sterile 0.9% sodium chloride solution and medical advice should be sought.

APPENDIX B

REFRIGERATOR TEMPERATURE RECORD

Record on Working Days

Date	Current Temp	Minimum Temp	Maximum Temp	Memory Cleared Tick ✓	Checked By (Signature)	Thermometer Reset Tick ✓

Signature of Designated Person monitoring:

Refrigerator defrosted and cleaned by: Date:

ASEPTIC TECHNIQUE

A: CARE OF INVASIVE DEVICES

INTRODUCTION

The following section provides guidance for the most commonly performed nursing procedures and clinical practices in relation to the control of infection. The following advice reflects current expert opinion and guidance incorporating relevant research and best practice recommendations.

Expert advice should always be sought should staff require it. Further guidance can be obtained from the following specialists:

- Nutritional Support Team
- Tissue Viability/Wound Management Nurse Specialist
- Respiratory Nurse Specialist
- Continence Advisor

PRINCIPLES OF ASEPSIS

Asepsis means “without micro-organisms” thus an aseptic technique is a method used to prevent contamination of wounds and other susceptible body sites or invasive device insertion sites by potentially pathogenic organisms which may lead to infection. This can be achieved by ensuring that clinical staff understand the principles, follow the recommended practices and that only sterile equipment is used during invasive procedures.

All staff performing invasive procedures or managing wounds should receive appropriate training.

INFECTION RISKS IN IMMUNOCOMPROMISED PATIENTS

Infection is caused by micro-organisms which invade the host’s immunological defence mechanisms, although susceptibility to infection may vary from person to person. The risk of infection is increased if the patient is immunocompromised by:

- Age – neonates and the elderly are more at risk due to less efficient immune systems
- Underlying disease – for example those patients with a severe debilitating or malignant disease or conditions such as diabetes
- Prior drug therapy – for example immunosuppressive drugs, steroids or broad-spectrum antimicrobials
- Patients undergoing surgery

In addition, the following factors should be considered when undertaking aseptic procedures on immunocompromised patients:

- Classic signs and symptoms of infection are often absent
- Untreated infection may disseminate rapidly
- Infections may be caused by unusual organisms or organisms which, in most circumstances are non-pathogenic i.e. do not cause disease
- Some antibiotics are less effective in immunocompromised patients
- Repeated infections may be caused by the same organism
- Super-infections, where a patient acquires a more pathogenic organism (of the same or a different species) than the one already causing infection, require nursing care of the highest standard, including strict adherence to aseptic technique to prevent such infections

WHEN TO USE AN ASEPTIC/NON-TOUCH (ANTT) TECHNIQUE

An aseptic technique should be used during any invasive procedure which breaches the body's natural defences e.g. the skin, mucous membranes, or when handling equipment which will enter a normally sterile area. The principles of asepsis should be applied to:

- Wound dressings
- Insertion *and manipulation* of invasive devices e.g. urinary catheters, all intravenous devices, PEG tubes etc.

THE PRINCIPLES OF ASEPSIS

Action	Rationale
Hand hygiene	Hand washing is the single most important procedure for preventing cross infection. Transient bacteria can be almost completely removed by effective hand hygiene techniques. In addition, resident bacteria (which can cause infections following highly invasive procedures) can be reduced by the use of an antiseptic detergent or the application of an alcohol hand gel following a social handwash. Hands should always be washed before and after contact with susceptible sites. Hand Hygiene may be required several times during a procedure.
Gloves	Gloves should be worn for all contact with mucous membranes and invasive devices e.g. urinary catheters. Sterile gloves should be worn for the insertion of invasive devices and minor surgical procedures. Clean, non-sterile gloves are acceptable for most wound care procedures and on-going device-related care.
Protective clothing	Water repellent plastic aprons will need to be worn to prevent staff clothing from becoming contaminated with bacteria from wounds or invasive devices. It will also protect the wound/invasive device from bacteria that may be present on staff uniform/clothing. Sterile impermeable gowns may be required for some minor surgical procedures.
Non-touch technique	The susceptible site should not come into contact with any item that is not sterile.
Equipment	All instruments, fluids and materials that come into contact with a wound, surgical site or during the insertion/manipulation of an invasive device, must be sterile to reduce the risk of contamination. This includes not only products used during the procedure but any final dressing (s). The sterility of the device/fluids/materials must be protected from contamination.
Dressing trolley	The trolley should be cleaned with detergent and water if it becomes physically contaminated. Alcohol wipes may be used between uses if necessary. The sterile field will normally protect the trolley from contamination. Ensure sticky tape residues are removed from the trolley rails. (Ideally these trolleys should not be used for other purposes). Alternatively, for some procedures, plastic trays may be used. These must be cleaned before and after each use.

CHRONIC WOUND MANAGEMENT

This section is written using the following guidance documents:

1. Department of Health (2011) *High Impact Intervention – reducing the risk of infection in chronic wounds care bundle*
2. Dougherty L and Lister S (editors) (2015) *The Royal Marsden Hospital Manual of Clinical Nursing Procedures Ch. 48 wound management 9th Edition*

Comprehensive advice on the management of wounds should be sought from specialist tissue viability nurses as this is a complex and constantly evolving practice. This section refers to those aspects of chronic wound care that may contribute to infection / cross-infection.

A chronic wound is defined as a wound that does not heal within an expected time frame i.e. 6 weeks despite optimal correction of any underlying pathological processes interfering with the body's normal process of wound healing. The majority of chronic wounds are:

- Venous ulcers
- Pressure ulcers
- Diabetic ulcers

Other types of chronic wounds include arterial leg ulcers and wounds from fungating carcinoma. Acute wounds may also become chronic.

In chronic wounds there is a clear increase in colonisation, bacterial burden and infection caused by micro-organisms, including MRSA. Chronic wounds colonised with MRSA are at increased risk for both wound infection and systemic infection (especially blood stream infections) particularly if another acute illness occurs requiring hospitalisation. Patients with MRSA-colonised wounds present an increased cross-infection risk to others and the environment. Early referral of patients with chronic wounds to specialist health professionals e.g. tissue viability teams and, in the case of diabetic foot ulcers, urgent referral to a multidisciplinary foot care team, is indicated to promote healing and reduce the risk of infection.

PREVENTING CONTAMINATION AND CROSS INFECTION

Wound care should only be carried out by those who are deemed competent to do so and have received training in the principles of asepsis and appropriate wound management. The principles of asepsis should be applied to all wounds irrespective of causation or type e.g. surgical wound, trauma wound, chronic wound etc.

Personal protective equipment – disposable apron and gloves – must be worn and changed between each patient

Wounds must be assessed as per local policy at every dressing change

The wound must be dressed creating an optimum wound healing environment according to the local wound management formulary

The use of systemic antibiotics is considered, as per local formulary, for non healing or progressive ulcers with clinical signs of localised and / or systemic infection
Dressing type and frequency of change, wound assessment and next wound review date must be routinely documented

There must be clear communication – between team members and with other health or social care providers – of those service users known to be infected or colonised with pathogenic micro-organisms inc. MRSA. This is a requirement of the Code of Practice 2018.

Service users with pressure ulcers must be placed on appropriate pressure relieving / reducing mattresses and cushions

Pressure is offloaded in service users with diabetic foot ulcers, including provision of appropriate footwear and insoles

In addition, the following may help to reduce wound contamination/cross-infection:

- Wound dressings are best performed in a designated treatment room, which is subject to regular cleaning
- Dirty dressings should be placed immediately into a clinical / offensive waste bag for disposal
- Wounds and any sterile equipment should be exposed for the shortest possible time. Wound temperature can fall by 12°C if the procedure is prolonged or the cleansing lotion is cold. It can take 3 hours or longer for the wound to return to normal temperature during which time cellular activity is reduced and therefore the healing process slowed. During exposure of the wound there is a much higher risk of environmental contamination of tissues particularly if wound care is undertaken in a well ventilated, draughty or high activity area
- Sodium chloride 0.9% (normal saline) is a physiologically balanced solution that is compatible with human tissue it dilutes bacteria is nontoxic to tissue and used at body temperature it is the safest and best cleaning solution for non-contaminated wounds
- Evidence has demonstrated there is no significant difference in the healing and infection rates in wounds irrigated with tap water or 0.9% sodium chloride and that although swabbing wounds may be effective in removing foreign bodies from the surface of a wound, irrigation is far less harmful to wound tissue. However, even if using tap water the principles of asepsis still apply.
- That tap water can be used for cleaning chronic wounds e.g. leg ulcers and pressure sores. However, even if using tap water the principles of asepsis still apply.
- Wound dressings should be kept dry at all times when in situ. Leakage from wounds e.g. leg ulcers will be contaminated with bacteria even if not clinically infected. “Strike through” can contaminate surfaces and hands leading to cross-infection.

INFLAMMATION AND INFECTION or bacterial burden

All chronic wounds are known to harbour a variety of bacteria to some degree and this can range from contamination through colonization to infection. When a wound becomes infected it will display the characteristic signs of heat, redness, swelling, pain, heavy exudate and malodour. The patient may also develop generalized pyrexia. However, immunosuppressed patients, diabetic patients or those on systemic steroid therapy may not present with the classic signs of infection. Instead they may experience delayed healing, breakdown of the wound, presence of friable granulation tissue that bleeds easily, formation of an epithelial tissue bridge over the wound, increased production of exudate and malodour and increased pain. Careful wound assessment is essential to identify potential sites for infection, although routine swabbing is not considered beneficial. Methods available for the management of wound infection or to decrease the bacterial burden in the wound include debridement, antimicrobial dressings e.g. those containing iodine or silver, topical negative pressure therapy and antibiotic therapy. Honey and essential oils have also been used. Appropriate antibiotic treatment of the infection should be determined from a positive wound swab.

WOUND SWABS

Routine wound swabs are not recommended unless there are clinical signs of infection or when non-healing persists. Many chronic wounds will be colonised with a variety of bacteria, the presence of which may not be clinically significant. Swabs, if indicated, should be taken from the base or margin of the wound following the removal of dressing residues and slough if present.

URINARY CATHETER MANAGEMENT – long-term urinary catheters

This section has been written taking account of the evidence base for practice published by the National Institute for Clinical Excellence (NICE) *Prevention of Healthcare-associated Infection in Primary and Community Care (2012 updated 2017)*. Where relevant, references to this guidance have been included in this text.

Urinary tract infections (UTI) account for approximately 23% of all Healthcare Associated Infections (HCAI). Most are associated with the use of an indwelling urinary catheter. Catheter-associated urinary tract infections (CAUTI) are a common complication occurring in over 90% of patients within 4 weeks of catheterization. The risk of catheter-associated bacteriuria (the presence of bacteria in the urine but not necessarily infection) increases by 5-8% a day during catheterisation and is inevitable in long-term catheterised patients *but does not necessarily require antibiotic treatment*.

Residents / patients in community and primary healthcare settings e.g. care homes, hospices or domiciliary care may require either short or longer term catheterisation. Long term catheterisation is defined as > 28 days. Infection prevention and control aspects of catheter management will be similar for both short and long term urinary catheterisation.

CAUTIs are caused by microbial contamination which is acquired by one of two routes – from urine becoming contaminated within the drainage system e.g. from the drainage bag (back-tracking up the system) or when the closed system is interrupted by disconnecting components of the system e.g. disconnecting the catheter from the bag; or via the space between the

catheter and the urethral mucosa (which can occur during catheterisation or subsequently as a result of poor management of the indwelling catheter / poor meatal care). In other words, bacteria travel up into the bladder along the inside or the outside of the catheter. Once bacteria invade one part of the system, all other areas are at risk.

CAUTIs are difficult to treat as bacteria adhere to the surface of the catheter in the form of a biofilm. Whilst antimicrobials may successfully kill bacteria in the urine, bacteria in the biofilm are generally less susceptible to these agents; consequently, they may still persist following treatment and potentially restart the cycle of infection.

As yet there is no catheter material that resists biofilm formation in the clinical setting; however, all-silicone catheters have been found to take longer to become encrusted than the silicone-coated, Teflon and latex catheters.

Practices to prevent infection e.g. asepsis should be applied to the insertion of the catheter, the management of the urinary drainage system and the care of the urethral meatus.

EDUCATION OF SERVICE USERS AND CARERS (Intervention 1.2.1 NICE)

If appropriate, teaching service users or family members to care for their own urinary catheters can minimise the risk of cross-infection. Education should include advice on careful hand hygiene, perineal cleansing and positioning of the drainage bag (and other catheter management issues of relevance).

EDUCATION OF HEALTH AND CARE PERSONNEL (Intervention 1.2.1 NICE)

Community and primary health and care personnel must be trained in catheter insertion, including suprapubic catheter replacement and catheter maintenance.

ASSESSING THE NEED FOR CATHETERISATION (Intervention 1.2.2 NICE)

Indwelling urinary catheters should be used only after alternative methods of management have been considered e.g. penile sheath.

The service user's need for catheterisation should be reviewed regularly and documented in the care plan / service user's notes and the urinary catheter removed as soon as possible.

Catheter insertion, changes and care should be documented.

CATHETER DRAINAGE OPTIONS (Intervention 1.2.3 NICE)

Following assessment, the best approach to catheterisation that takes account of clinical need, anticipated duration of catheterisation, patient preference and risk of infection should be selected. This may be indwelling (urethral or supra-pubic); intermittent catheterisation or a penile sheath. Intermittent catheterisation should be used in preference to an indwelling catheter if it is clinically appropriate and a practical option for the service user. A choice of either single-use hydrophilic or gel reservoir catheter should be offered to those patients for whom intermittent self catheterisation is the preferred option

For indwelling urinary catheters, the type and gauge will depend on an assessment of the patient's individual characteristics including: age; allergy or sensitivity to catheter materials; gender; history of symptomatic urinary tract infection; patient preference and comfort; previous catheter history and reason for catheterisation.

In patients for whom it is appropriate, a catheter valve can be used as an alternative to a drainage bag. Consideration needs to be given to mental acuity, manual dexterity, clothing preferences and use of night drainage bags when considering the use of catheter valves. Smaller gauge urinary catheters (12-14 Ch) with a 10 ml balloon (3-5 ml in children) inflated with sterile water minimise urethral trauma, mucosal irritation and residual urine in the bladder which are all factors which predispose to CAUTI.

NB manufacturers' instructions for inflation should always be followed as a priority.

URINARY CATHETER INSERTION – IPC aspects – (Intervention 1.2.4 NICE)

Catheters should only be inserted by staff that have been trained and assessed as competent to undertake this procedure.

All catheterisations should be aseptic non-touch (ANTT) procedures.

The urethral meatus should be cleaned with sterile water or saline (or in accordance with local policy) before inserting the catheter

Instil single-use lubricating gel into the urethra prior to insertion to minimise urethral trauma and infection

Insertion technique:

- Hands should be washed and sterile gloves and a plastic apron should be worn
- Any gross contamination of the perineal area should be removed using soap and water prior to meatal cleaning, cleaning from front to back to avoid transference of bowel flora to the urethral area particularly in female service users
- Catheter insertion, changes and care should be documented in the service user's notes

CATHETER MAINTENANCE – leave the closed system alone! – (Intervention 1.2.5 NICE)

Indwelling urinary catheters should be connected to a sterile closed urinary drainage system or catheter valve.

Staff should ensure that the connection between the catheter and the urinary drainage system is not broken except for good clinical reasons e.g. changing the bag every 5 – 7 days (or in line with manufacturers' recommendations)

Disconnecting any part of the closed system can contribute to contamination and cross-infection. The use of drainage systems without a drainage port should be avoided where possible as this requires the bag to be changed daily.

Staff must decontaminate their hands and wear a new pair of clean, non-sterile gloves before manipulating a patient's catheter, and must decontaminate their hands after removing gloves. *A new pair of gloves must be used for each task on a patient.*

Urine samples should be obtained from a sampling port using an aseptic technique.

Using a drainage system without a sampling port requires disconnection of the system to obtain a specimen of urine which increases the likelihood of contamination/infection. The alternative is to collect directly from a clean drainage bag which is an inaccurate method which may lead to false results.

Urinary drainage bags should be positioned below the level of the bladder, and should not be in contact with the floor i.e. should be hung on an appropriate stand. If the bag is raised above bladder height *even temporarily such as during moving and handling* this can result in backflow and increased risk of infection.

A link system should be used to facilitate overnight drainage, to keep the original system intact. The urinary drainage bag should be emptied frequently enough to maintain urine flow and prevent reflux, and should be changed in line with manufacturers' recommendations (usually every 5 – 7 days) or when clinically indicated e.g. malodorous or damaged.

Drain the bag either into the toilet or receptacle / jug. If using a receptacle / jug ensure it is either disposable or kept for that individual and decontaminated in a bed pan washer *after each use*. When opening and closing the drainage tap ensure there is no spillage of urine, dry the tap outlet to prevent this and clean with an alcohol wipe. Avoid contact between the drainage tap and the receptacle / jug as this can increase the risk of contamination.

Bags that are non-drainable should be used **once** e.g. overnight and emptied before disposal. Meatal cleansing is no longer recommended. However it is advisable to keep the urethral meatus clean and free from debris. The use of soap and water once or twice a day is recommended. This can usually be undertaken whilst showering or bathing.

Catheters should be changed only when clinically necessary or according to the manufacturer's current recommendations.

To minimise the risk of blockages, encrustations and catheter-associated infections for patients with a long-term indwelling urinary catheter: a patient-specific care regimen should be developed; consider approaches such as reviewing the frequency of planned catheter changes and increasing fluid intake; document catheter blockages.

Bladder instillations or washouts must not be used to prevent catheter-associated infections. When changing catheters in patients with a long-term indwelling urinary catheter: do not offer antibiotic prophylaxis routinely; consider antibiotic prophylaxis for patients who have a history of symptomatic urinary tract infection after catheter changes OR experience trauma* during catheterisation. Patients should always be assessed for signs of infection before a catheter change

*Defined as frank haematuria after catheterisation or two or more attempts of catheterisation.

MANAGEMENT OF VASCULAR ACCESS DEVICES

INTRODUCTION

This section has been written taking account of evidence-based practice contained in the following:

1. Department of Health (2009) High Impact Intervention 2: *peripheral intravenous cannula care*
2. Dougherty L, Lister S (editors) (2008/15) *The Royal Marsden Hospital Manual of Clinical Nursing Procedures Ch. 45: vascular access devices: insertion and management*
3. National Institute for Clinical Excellence (NICE) *Prevention of Healthcare-associated Infection in Primary and Community Care (2012 Updated 2017)*.

Intravenous cannulae (IVCs) pose a risk of direct microbial entry to the bloodstream. Cannulae can become contaminated at the insertion site by skin micro-organisms (that can gain access to the bloodstream by migrating down the body of the device) or by other micro-organisms via the cannula hub or injection port.

EDUCATION

Healthcare workers caring for a patient with a vascular access device should be trained, and assessed as competent, in using and consistently adhering to evidence-based guidelines. It is recommended that competency is re-assessed regularly at defined intervals. This assessment should be documented.

Patients and their carers should be taught any techniques they may need to use to prevent infection and safely manage a vascular access device prior to discharge from hospital. They should also have access to follow-up training and support.

ASEPTIC -NON TOUCH (ANTT) TECHNIQUE

An aseptic non touch (ANTT) technique must be used for vascular access device catheter site care and when accessing the system. This will include:

- Insertion / cannulation
- IV drug / fluid administration
- changing of administration set / extension tubing
- securing of site and dressing

The technique used should ensure:-

- Hands are decontaminated before accessing or dressing a vascular access device
 - Equipment used is effectively decontaminated e.g. procedure trays
 - Key parts of sterile equipment are managed to avoid contamination
- Access points (e.g. cannula ports) are effectively decontaminated before accessing

SITE CARE

- Decontaminate the skin at the insertion site with chlorhexidine gluconate in 70% alcohol prior to insertion. A combined chlorhexidine with alcohol preparation has the advantage of residual activity for 4 – 6 hours following application. Solutions should be applied with friction for up to 1 minute and allowed to air dry for 30 – 60 seconds. It is essential to allow time for drying in order for disinfection to be completed.
- Use a sterile transparent semi-permeable membrane dressing to cover the device insertion site.
- Consider using a sterile gauze dressing covered with a sterile transparent semi-permeable membrane dressing **only** if the patient has profuse perspiration or if the insertion site is bleeding or oozing. If used, a gauze dressing must be changed every 24 hours (or sooner if soiled) **and** its use discontinued as soon as possible and replaced with a sterile transparent semi-permeable membrane dressing.
- Leave the transparent semi-permeable membrane dressing for the life of the cannula, provided that the integrity of the dressing is retained.
- If required, individual sachets of antiseptic solution or impregnated swabs or wipes should be used to disinfect the dressing site.
- The insertion site should be checked regularly – at least twice a day - for signs of phlebitis (erythema, pain and / swelling) or for signs of infection. The inspections should be routinely documented. Ideally, a standardised system should be used e.g. the Visual Infusion Phlebitis (VIP) score which uses a colour-coded system to score inspection details.

GENERAL PRINCIPLES FOR MANAGING DEVICES

- Decontaminate the infection port or catheter hub before and after accessing the system using chlorhexidine gluconate in 70% alcohol. Consider using an aqueous solution of chlorhexidine gluconate if the manufacturer's recommendations prohibit the use of alcohol with their catheter.
- If needleless devices are used, the manufacturer's recommendations for changing the components should be followed.
- When needleless devices are used, healthcare workers should ensure that all components of the system are compatible and secured, to minimise leaks and breaks in the system.

When needleless devices are used, the risk of contamination should be minimised by decontaminating the access port with either alcohol or an alcoholic solution of chlorhexidine gluconate before and after using it to access the system.

Avoid the use of multidose vials in order to prevent the contamination of infusates.

REPLACEMENT OF CANNULAE / ADMINISTRATION SETS

The need for intravenous access devices should be reviewed regularly and devices removed as soon as possible. Cannula/dressing labels are recommended to record the date of review/replacement.

Peripheral cannulae should be replaced every 72 – 96 hours. If this is not possible due to lack of access to veins then this should be recorded together with documentation relating to the condition of the site e.g. no inflammation noted.

In general, administration sets in continuous use need not be replaced more frequently than at 72-hour intervals unless they become disconnected or a catheter- related infection is suspected or documented.

Administration sets for blood and blood components should be changed every 12 hours, or according to the manufacturer's recommendations.

AUDIT

Regular audit of the management of intravenous devices should be undertaken using Dept. Of Health High Impact Intervention tools available at www.hcai.dh.gov.uk

MANAGEMENT OF RESPIRATORY EQUIPMENT

INTRODUCTION

Respiratory equipment such as nebulisers and humidifiers may act as potential sources of infection. Bacteria may colonise respiratory equipment and deliver contaminated air directly into the lungs leading to respiratory tract infection or may be transmitted to other residents on the hands of staff. It is essential that all respiratory equipment is appropriately managed and decontaminated in order to prevent this.

This chapter does not give guidance on the management of invasive ventilation or tracheostomy tube management. If invasive ventilation / tracheostomy care is undertaken there should be local policies on this which should include the prevention of ventilator associated infections. Guidance should be sought from the local respiratory specialist team.

For further information on respiratory equipment refer to Section – Single Use / Single Patient Use Devices

NEBULISERS

The majority of nebuliser systems are used to deliver drugs, which open up the lungs and improve breathing. The type of nebuliser and / or drug used depends on the service user's needs; the choice should be based on the medication to be administered and the ease of use by staff and/or service users.

Nebuliser mask and tubing may be single use or single patient use. Manufacturer's instructions must be followed.

HUMIDIFIERS

Humidifiers saturate inspired air with water, in order to prevent drying of the airways during oxygen therapy. Either heated or unheated humidifiers can achieve this.

RISK FACTORS FOR THE CONTAMINATION OF RESPIRATORY EQUIPMENT

- Fluid residues left in the nebuliser after use can provide an ideal medium for the multiplication of bacteria
- Shared use of equipment between service users can lead to cross- infection
- Non-sterile fluids cannot be guaranteed to be free of contamination from harmful micro-organisms
- Condensation accumulating in the tubing may become colonised with harmful bacteria

MINIMISING THE RISK OF INFECTION

- Staff must wash their hands before and after handling respiratory-therapy equipment. Gloves should also be worn if contamination with respiratory secretions is anticipated.
- Only sterile single-dose fluids / medications are recommended for nebuliser therapy. If vials of multi-dose medication are used, handle, dispense and store according to the manufacturer's recommendations
- Only sterile water should be used to fill humidifiers. They should be emptied daily, washed, dried and re-filled.
- Nebulisers must be washed in warm water (without detergent) and dried thoroughly after each treatment. They should be stored covered after use.
- Nebulisers should be routinely changed as per manufacturer's instructions.

Humidifier tubing should be changed regularly in accordance with manufacturer's recommendations.

MANAGEMENT OF ENTERAL FEEDING / ESTABLISHED PERCUTANEOUS ENDOSCOPIC GASTROSTOMY (PEG)

INTRODUCTION

This section has been written taking account of the evidence base for practice contained in:

1. Department of Health (2011) *High Impact Intervention: Enteral feeding care*
2. National Institute for Clinical Excellence (NICE) *Prevention of Healthcare-associated Infection in Primary and Community Care (2012 updated 2017)*

Enteral tube feeding is an accepted method for the provision of nutrition in individuals who are not able to take any foods orally or whose daily oral intake is not sufficient to meet their nutritional requirements.

The majority of individuals receiving an enteral feed have the product administered directly into their stomach via a gastrostomy or percutaneous endoscopic gastrostomy (PEG) tube or via a naso-gastric tube. Enteral feeding is the preferred and most physiologically normal method of artificial feeding, however, the risks of bacterial contamination of the feed and the possible risks of infection around a PEG site can lead to complications and need to be considered and addressed.

Contamination of feeds is a key concern in long term feeding / home enteral tube feeding as it has been found that more than 30% of feeds in hospital and home are contaminated with a variety of micro-organisms, largely due to the preparation or administration of feeds, and this has been linked to serious clinical infection. The elderly are particularly vulnerable to the effects of food poisoning. Therefore, it is essential to have robust policies and procedures to minimise the risk of food poisoning as a result of enteral feeding and to meet the requirements of current Food Hygiene Regulations.

All staff who handle enteral feeding systems should be trained in feed delivery and management of the administration system. Training should be documented.

MICROBIAL HAZARDS OF ENTERAL TUBE FEEDING

Infection associated with enteral feeding has been reported on numerous occasions. Enteral infections have been reported e.g. Salmonellosis but of greater concern is the incidence of pneumonia and bacteraemia associated with contaminated enteral feeds. Expert guidance reinforces the need for rigorous infection control procedures in the handling and delivery of enteral tube feeds.

SOURCES OF CONTAMINATION OF ENTERAL FEEDING SYSTEMS

There are a number of possible sources of contamination in an enteral feeding system. These can be summarised as:

PROBLEM	POSSIBLE SOURCE
Touch contamination of equipment	Lack of, or poor hand hygiene, poor non-touch technique and excessive manipulation of system
Inappropriate storage of feed	Stored in a contaminated area; failure to refrigerate where appropriate; opened feeds kept for too long
Poorly designed equipment	Multiple / exposed parts prone to touch contamination during assembly / use
Misuse of equipment	Prolonged use or re-use of administration sets and syringes
Contaminated additive	Medications or flush solutions
Site problems	Colonisation or infection of the PEG site
Cross-infection	Failure to adequately decontaminate equipment (e.g. pumps) between service users; lack of hand decontamination
Contaminated feed	During reconstitution, decanting, handling, manufacture, transportation or storage

SELECTION OF EQUIPMENT / SYSTEMS TO REDUCE HAZARDS

When selecting an enteral feeding system, it is important that the possible risks of introducing bacterial contamination are considered. The following issues should be considered when selecting equipment:

- Pre-packaged, ready-to-use feeds, to which a giving set can be directly attached, are preferable to those that need decanting, reconstitution or dilution
- The system selected should require minimal handling to assemble and be compatible with the service user's enteral feeding tube
- The feed container should have a lid that can be removed without hands touching the neck of the container to which the set will be attached
- A system that requires a minimum number of connections is recommended; 3 way taps should be avoided
- Feeds which come in collapsible bags and, therefore, are non-air dependent are preferable as they reduce the risk of airborne contamination
- Pre-filled containers with larger volumes, e.g. 1000ml or 1500ml, reduce the number of container changes and therefore reduce the risk of handling-associated contamination
- Enteral feeding pumps with flush panels are easier to wipe clean than pumps with lots of grooves and knobs

FEED PREPARATION

Always decontaminate hands thoroughly either with soap and warm running water or the application of alcohol hand rub before commencing preparation of feed.

When decanting, reconstituting or diluting feeds a clean, dry working area should be prepared and equipment dedicated for enteral feed use only should be used. Feeds to be attached to feed equipment should be taken to the patient using either a clean trolley or tray. Items should NOT be placed on beds or other surfaces which are not capable of being cleaned with detergent prior to use.

Raw foods such as meat, fish, eggs and vegetables should never be handled in close proximity to enteral feeding solutions or equipment. Pets should not be allowed near the feeding solutions or feeding equipment.

Bottle openers should be decontaminated prior to use (e.g. in a dishwasher). If scissors are required, they should be sterile prior to first use. Bottle openers and scissors used for opening sterile feeds should be dedicated for use with enteral feeding products only. They should be decontaminated after use, preferably in a dishwasher or by washing with hot water and detergent, rinsed, dried and stored covered.

Feeds should be mixed using cooled *freshly boiled* water or freshly opened sterile water and a no-touch technique with minimal handling of all component parts. Water should NOT be stored but should be discarded after each use.

ASSEMBLY OF FEEDS

There are three methods of assembling an enteral feed system:

- Ready-to-use sterile feeds
- Decanting sterile feeds
- Special or Modified feeds

Each requires a slightly different set of handling procedures and hanging times:

READY-TO-USE STERILE FEEDS

Prior to use, store feeds in a clean, dry environment according to the manufacturers' instructions and where applicable, food hygiene safety. The temperature in the storage area should not drop below 5°C or rise above 25°C.

Prior to use, check the feed expiry date and for any sign of damage to the container. Never use feeds that have expired or are in damaged containers.

Do not add any water, medication or other substances directly to the feed unless prescribed for this purpose.

Sterile feeds that have been opened, but not immediately connected to a sterile giving set can be resealed and stored on the top shelf in the body of a refrigerator labelled with date and time of storage. The feed must be stored at below 5° C and only for up to 24 hours (the fridge temperature should be checked regularly with a fridge thermometer). Once a sterile feed has been opened it must be used within 24 hours or discarded.

Feeds should never be stored near or below products such as raw or thawing meat or fish because cross contamination may occur

DECANTING STERILE FEEDS

Follow the above guidelines on ready-to-use sterile feeds AND consider the following additional points.

Sterile feeds should only be decanted when:

- the feed cannot be directly attached to a giving set, e.g. when using ring- pull cans or when there is no suitable ready-to use preparation in the size/volume required, e.g. for overnight feeding
- it is necessary to make additions to a sterile feed (see special or modified feeds)
- The feed is going to be given as a bolus via a syringe. (Feeds given as a bolus should be administered at room temperature. Therefore, if a feed has been stored in the fridge it should be taken out 30 minutes prior to administration and allowed to warm up. Keep the feed in its resealed container until it is poured into the syringe.)
- The following should be considered if decanting sterile feeds:
- Sterile feeds should be decanted into a sterile container.

- Sterile feeds that have been decanted into a sterile container can hang for a maximum of 24 hours before being discarded
- If decanting, decant the full volume required for the 24 hours. Do not top up the reservoir
- Decanting should be undertaken in a clean environment e.g. kitchen.
- When decanting feeds, use a non-touch technique. This means avoiding touching (with hands, objects or surfaces) any openings or connecting parts of the feed container or reservoir container.

The design of some feed containers causes the feed to come into contact with the outside of the container when it is being decanted, i.e. a ring-pull can. For this reason, wipe the outside of the feed container with a wipe impregnated with 70% isopropyl alcohol. Allow the alcohol time to dry before decanting.

ADMINISTRATION OF FEEDS

Use minimal handling and an aseptic technique to connect the administration system to the enteral feeding tube

Administration sets and feed containers are for single use and must be discarded after each feeding session.

MAXIMUM HANGING TIME FOR ENTERAL FEEDS

FEED TYPE	HANGING TIME
Modified or mixed feeds decanted into a sterile reservoir	4 Hours
Sterile, ready-to-use feeds, if not decanted	24 Hours
Sterile feeds decanted into a sterile reservoir	24 Hours

EQUIPMENT CARE

Re-usable medical equipment e.g. pumps and stands must be cleaned and serviced according to the manufacturer's instructions. Such equipment should be cleaned using hot water and detergent on a regular basis as part of routine equipment cleaning schedules. Enteral feed solutions can be difficult to remove from equipment if left to dry after spillage so prompt cleaning is recommended.

Equipment requiring servicing or repair should be cleaned, decontaminated and have the necessary documentation accompanying it, (See Decontamination Policy)

PERSONAL PROTECTIVE CLOTHING

To reduce the risk of infection a new set of disposable non-sterile gloves should be used each time the enteral feeding system is handled. Disposable plastic aprons should be worn whenever the feeding system is handled.

CARE OF INSERTION SITE AND ENTERAL FEEDING TUBE

The stoma site should be inspected and washed daily with water and dried thoroughly. Dressings are not necessary once the stoma has healed (following its placement) which is usually after 10 – 12 days. The tube should be rotated 360° regularly to avoid infections related to “buried bumper syndrome”.

To prevent blockage, the enteral feeding tube should be flushed with freshly drawn tap water before and after feeding or administering medications if the service user is immune-competent. For immune-suppressed individuals the flushing water should be either cooled *freshly* boiled water or sterile water from a *freshly* opened container. Do NOT store any water between uses even in the ‘fridge.

Single use syringes (discarded after single use) or single-patient use syringes should be used. The latter should be discarded according to manufacturer’s instructions. If used more than once, single patient use syringes should be washed after use with warm water and detergent (after removing plunger), rinsed thoroughly in clean, running water; shaken to remove excess water; dried as much as possible and stored in a covered container until next use. Replace plunger immediately prior to use – do not replace inside barrel as this will deter drying the inside.

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

ASEPSIS IN MINOR SURGICAL PROCEDURES

INTRODUCTION

This section has been written using the professional guidance issued by the Association for Perioperative Practice and contained in: Standards and Recommendations for Surgery in Primary Care AfPP 2016

Reference should be made to the general principles of aseptic technique listed in page 3 of section A (Aseptic Technique and Care of Invasive Devices). These principles, correctly applied, will help prevent contamination of an open wound (during surgical procedures) or sterile body cavity (e.g. for Joint Injections).

The basic principles of aseptic technique prevent contamination of the open wound, isolate the operative site from the surrounding non-sterile physical environment and create, maintain and promote a sterile field so that surgery can be performed safely.

This section is not intended to provide comprehensive guidance on minor surgical procedures. Its purpose is to highlight those practices that have an impact on the prevention and control of the development of post-operative wound infection.

Guidance on the minor surgery environment is available in section 14 Estates and Facilities Management.

General considerations

All staff involved in the preparation and performance of surgical procedures must receive competency assessed training in aseptic surgical techniques. This should include surgical hand scrub and gown/glove donning procedures. The management and use of sterile instruments should also be taught and assessed.

Staff with infected lesions of the skin or bacterial infections of the upper respiratory tract should not participate in any aseptic technique.

The environment and all working surfaces must be cleaned in accordance with local policies prior to the commencement of any aseptic procedures.

If asepsis is compromised immediate action is required. Contaminated items should be removed and discarded. If the sterile field is compromised then a new field is required. Re-gloving and re-gowning may also be required.

Equipment and medical devices

All pre-sterilised articles must be checked for evidence of sterilisation, damage, integrity of packaging and expiry date prior to use. Any packs found to be in an unsatisfactory condition must be discarded.

Items used within a sterile field *must be sterile*. Any items that fall into an area of questionable cleanliness must be considered non-sterile. This is of particular importance where medical devices contain more than one component part which may involve a disposable element and a

re-usable element e.g. diathermy forceps.

In procedures involving 'knife to skin' a sterile drape is required. This should be handled by the edges only, and applied from surgical site to periphery. Once in situ these should not be rearranged.

Sterile drapes should conform to EN 13795 (European Committee for Standardization 2002) and be used correctly to establish a sterile field.

Sterile drapes should be handled as little as possible. The drapes should be applied from the surgical site to the periphery, avoiding reaching over non-sterile areas. Once placed, drapes should not be repositioned in order to avoid contamination of the sterile field.

Scrubbed personnel

Sterile gloves should be worn for all invasive procedures by the clinician undertaking the procedure and any scrub assistant who manipulates the sterile field or instruments. Sterile Gowns should be worn for all 'knife to skin' procedures. These should comply with standard EN 13795. Care should be taken when donning gowns to avoid contaminating the front of the gown.

Scrubbed personnel should remain close to the sterile field and not leave the immediate area. If personnel leave the sterile field and exit the minor surgery area they must re-scrub before returning to the sterile field. Leaving the sterile field increases the risk for potential contamination.

Personnel participating in sterile procedures must stay within the sterile boundaries; a wide margin of safety should be given between scrubbed and non-scrubbed personnel.

When changing positions or moving between sterile areas, scrub personnel should turn back to back or face to face to avoid contamination.

Scrubbed personnel must keep their arms and hands within the sterile field at all times. Contamination may occur if hands are moved below the level of the sterile field.

Scrubbed personnel should only be seated when the operative procedure is to be performed at that level.

Circulating personnel should not walk between sterile fields (e.g. between a prepared patient and the instrument trolley) and should be aware of keeping an adequate distance from the sterile field.

Special considerations

Dressings must be removed carefully from the wound to prevent scattering of micro-organisms into the air; it is recommended that this is carried out by an assistant wearing gloves and not a member of the scrubbed team. Used and soiled dressings should be discarded immediately and in accordance with local policy.

To reduce the risk of airborne cross infection, talking, movement, opening and closing doors, exposure of wounds, disturbance of clothing and linen and number of personnel in the minor surgery area should be kept to a minimum. Special consideration must be taken to maintain the integrity of the sterile field at all times.

The sterile field should be constantly monitored and maintained, as sterility cannot be assured without direct observation. Any break in sterility must be reported and acted on to ensure patient safety.

Sterile fields should be prepared as close as possible to the time of use.

Procedure trolley

A designated area, which affords sufficient space to open packs whilst maintaining a sterile field, should be identified for this procedure. There should be minimal movement of personnel within this area during the preparation of the trolley.

A trolley of appropriate size is required for sterile instrumentation and products and this may be influenced by the type of procedure / surgery being carried out. This area, along with the wound site, comprises the sterile field. Fields should be protected from contamination by unsterile items or by non scrubbed staff. Care must be taken when opening items onto the sterile field e.g. additional instruments/dressings/fluids, to ensure the field is not compromised.

All trolleys should adhere to the Medical Devices Directive 93/42/ EEC (incorporated into UK law in 2002) and be stable and robust enough for the intended job and in good condition (free of surface abrasions) and in sound working order. Ease of cleaning should be taken into account when making product choice as should ease of movement and height. Trolleys should be included in a planned prevention maintenance programme. Particular attention to wheel mechanisms is required in order to allow free and smooth movement. Trolleys, mayo stands and bowl stands should be made of aluminium, stainless steel or mild steel covered in nylon.

Preparation of sterile trolleys in advance, with the use of sterile sheets to cover them, is not recommended. The trolleys are subject to contamination over time and removal of sheets without contamination cannot be guaranteed. In addition, unless trolleys are continuously monitored, there is a potential for sterility to be compromised

Scrubbed personnel should move draped sterile trolleys by placing hands on the horizontal surfaces only.

To maintain asepsis it is essential that all staff are aware of the correct method of opening different sterile packages to avoid the contamination of contents. Circulating practitioners should open wrapped sterile supplies by opening the wrapper flap furthest away from them first. The nearest wrapper should be opened last. Outer wrappers should be secured when presenting sterile items, to avoid contamination. The scrubbed person opens packs towards themselves first and then away, to avoid contamination of the sterile item.

Sterile items should be presented to the operating or scrubbed practitioner or placed securely on a specific area of the sterile field identified and managed by the operating or scrubbed

practitioner. Items should not be tossed onto a sterile field as they may roll off or cause other items to be displaced.

Sharps and heavy items must be presented to avoid penetration of the sterile field. Sharps should be opened into a container to avoid sharps injury and damage to the sterile field. Needles and scalpel blades may pose a risk to staff during procedures if safe practices are not followed. Sharps items should never be passed from hand to hand, whether used or not. A 'neutral zone' should be identified.

If re-usable blade handles are used, the blade should be removed using a dedicated device to prevent injury to the operator.

When dispensing solutions, the solution vessel should be placed near the trolley edge or held by the operating or scrubbed practitioner. The solution should be poured slowly to avoid splashing which could cause strike-through and compromise the sterile field.

The edge of a container is considered contaminated after the cap is removed and therefore the sterility of its contents cannot be guaranteed if the cap is replaced.

Preparation of sterile trolleys in advance, with the use of sterile sheets to cover them, is not recommended. The trolleys are subject to contamination over time and removal of sheets without contamination cannot be guaranteed. In addition, unless trolleys are continuously monitored, there is a potential for sterility to be compromised.

Trolleys should be positioned close together to ensure that there are no breaks in the sterile field.

The disposal of all equipment, drapes and sharps must be carried out in accordance with local and national guidelines. The scrub person should be considered the person of choice to dispose of all contaminated materials whilst still gowned and gloved.

PATIENT SKIN PREPARATION IN THE REDUCTION OF SURGICAL SITE INFECTION

Skin preparation is the process by which the skin is cleansed to reduce the number of transient and resident skin bacteria before surgical incision. Transient bacteria do not normally colonise the skin and are easily removed, whereas resident bacteria grown on normal skin and are difficult to remove. Most wound infections are associated with the patient's own skin flora and thus skin must be prepared to reduce the risk of surgical site infection.

The purpose of skin preparation is to remove dirt and debris from the skin, reduce the number of micro-organisms, inhibit the re-growth of further micro-organisms and reduce the number of micro-organisms entering the wound site, thus reducing the potential for surgical site infection. Skin preparation should not cause irritation to the skin.

HAIR REMOVAL

Hair removal is often a routine part of pre-operative preparation but staff and patients need to be aware of the evidence and rationale for this practice as research has shown that removal of hair is not always necessary and should only be undertaken after assessment of the individual patient. The removal of hair is only necessary if it will directly interfere with access to the incision site or if there is a risk it will contaminate the wound site. Systematic review (Tanner et al 2006) has shown no difference in surgical site infection rates among patients who have had hair removal prior to surgery and those who have not.

If hair removal is undertaken the following is recommended:

- Patient consent must be obtained with a full explanation of the method to be used and why it is necessary
- Method of hair removal should be decided between the patient and the clinician performing the procedure
- Hair removal should take place as close to the time of surgery as possible to minimise the risk of bacterial contamination of the skin surface
- Hair removal should be carried out by an experienced practitioner in a clean area of the surgical suite with good lighting, affording patient privacy at all times
- Details including who performed the hair removal, the area from where the hair was removed and the method used should be documented.

Methods of hair removal

Depilatory creams are effective and may be carried out by the patient. However, skin irritation can occur and a patch test is recommended prior to use. This method is also costly.

Clipping – using an electric or battery-powered clipper with a disposable or re-usable head (that can be disinfected) – is a simple and less irritating method than shaving.

Wet shaving causes the most trauma to skin and carries the highest risk of postoperative wound infection. Wet shaving should not be used unless other methods are not suitable.

SOLUTIONS USED FOR SKIN PREPARATION

Antiseptics have to be effective against transient and resident micro-organisms. They should have a broad spectrum of microbial activity with a fast and lasting effect against Gram negative and Gram positive bacteria, as well as viruses and fungi. They should be resistant to inactivation by organic matter, be non-toxic and acceptable cosmetically.

Antiseptics should be supplied in ready-to-use, single use containers or sachets as sterility is not guaranteed once open and there is a risk of contamination from using multi-use containers. Solutions must be licensed as a preparation suitable for skin disinfection prior to a surgical procedure

Skin preparation fluid should not be kept on the sterile trolley following skin preparation. This supports good infection prevention principles and patient safety. Only solutions that are in labelled containers should remain on the sterile trolley (NHS England 2015)

Types of skin preparation include:

- Povidone-iodine alcoholic solution
- Povidone-iodine aqueous solution
- Chlorhexidine 0.5% in 70% industrial methylated spirit (IMS)
- Iodine 1% in IMS
- 70% iodine in spirit
- Chlorhexidine gluconate 0.015% and cetrimide solution
- 70% IMS

Alcohol solutions are deemed to be more efficient than aqueous solutions

Decisions regarding the preparation to be used should be influenced by the area which requires preparation, the condition of the skin and patient allergies.

Delicate areas, such as eyes and ears may require special or diluted solutions. Chlorhexidine is not recommended for facial prep and iodine may cause corneal damage if introduced into the eye. If solutions enter the inner ear they may cause sensorineural deafness. Chlorhexidine gluconate and alcohol or alcohol-based solutions should also be avoided on mucous membranes.

When using an alcohol-based solution, it is imperative that skin is allowed to dry completely after each application and before applying electrocautery or laser treatment. Spontaneous combustion can occur when flammable solutions are exposed to an ignition source when oxygen is present.

Skin solutions should be kept in a locked cupboard and particular attention should be paid to storage of flammable solutions according to The control of substances hazardous to health (COSHH) regulations 2002.

ISOLATION OF INFECTIOUS PATIENTS IN GENERAL PRACTICE

INTRODUCTION

The aim of isolation is to contain and prevent the spread of potential or known pathogenic or epidemiologically important organisms in order to reduce the risk of transmission of infection to and from service users, visitors or staff.

Identifying patients with suspected infections in General Practice can be a challenge as many patients visiting the surgery will not be aware of their potentially infectious status or may not communicate this in advance to staff.

Patients visiting the practice with known infections (or colonisation with transmissible organisms such as MRSA) provide less of a challenge but consideration is still required to ensure that risks are reduced / avoided during their practice visit.

Patients being visited in their own homes (or in other community environments, such as care homes) also pose a risk to staff attending to provide care.

INDICATIONS FOR ISOLATION IN GENERAL PRACTICE

There are a number of circumstances in which suspected or known infections may present in general practice. These include (but are not limited to):

- Diarrhoea and / or vomiting e.g. norovirus; *C. difficile* diarrhoea; bacterial food-poisoning
- Suspected or clinically proven infection which may be transmitted through the respiratory / airborne route e.g. influenza, chickenpox, measles, TB, group A streptococcal sore throat, MRSA etc.
- Suspected or clinically proven infection which may be transmitted via the contact route e.g. MRSA, group A streptococcal infection in wounds

COMMUNICATION

Informing patients of their responsibilities to limit the spread of infection is difficult. Posters and other visual means of identifying risks (inc. translation into local languages) can help to inform patients of the need to communicate symptoms when they attend the practice.

Reception staff should be trained (at induction and during mandatory IPC training) to identify those symptoms which may indicate transmissible infections e.g. rash, diarrhoea, vomiting etc. when receiving calls from patients requesting appointments or visits. Reception staff are NOT expected to take any other action other than to notify medical or nursing staff if they are concerned about patients' symptoms so that effective arrangements can be put in place e.g. being placed in (and examined / treated) in a separate room.

Medical and nursing staff visiting patients in their own home or in residential care environments should be trained (as part of professional update training) to identify risks that may require

additional precautions to be taken e.g. single room isolation in a care home for a resident with diarrhoea/vomiting).

All professional staff attending patients with suspected / known infections should be familiar with the route of spread (of infection) and standard infection prevention and control precautions to be used whilst providing care. Staff should also be familiar with additional interventions such as enhanced cleaning of equipment and the environment which may be required following care (and which may require additional time to undertake).

SEGREGATION OF PATIENTS IN GENERAL PRACTICE

WHILST AWAITING APPOINTMENT

Patients with suspected / known infections *spread by the respiratory / airborne route* should be segregated whilst awaiting their appointment and **ideally** should be examined in the same room to minimise the risk of environmental spread to other clinical areas. A separate consulting room is ideal. Chlorine clean is required after the consultation to reduce the risk to other patients.

Patients with diarrhoea / vomiting should ideally not attend surgery. However, in such circumstances patients should also be segregated as above. In addition, *where possible*, a separate toilet should be made available together with items such as disposable bowls and wipes. Patients should be examined at the earliest opportunity. Any toilet facilities used by symptomatic patients should be thoroughly cleaned with a solution of sodium hypochlorite (bleach) prior to being used by other patients.

Patients with suspected / known infections *spread by the contact route* e.g. those with colonised or infected wounds requiring dressing can wait in general waiting areas. Usually these patients will be returning for regular wound care. In such cases, nursing / medical staff should be encouraged to provide appointments at the end of clinics to allow for additional cleaning of equipment / environment following care.

INTERVENTIONS TO REDUCE RISK

Standard infection prevention and control precautions (SICPs) should be used at all times with all patients. Strict attention to SICPs is necessary whilst examining / treating patients with suspected / known infections in general practice:

- Thorough hand decontamination
- Appropriate use of PPE especially gloves
- Disposal of infectious waste into yellow bags
- Thorough cleaning of ALL medical devices / equipment used
- Use of single use, disposable medical devices where appropriate
- Cleaning of all environmental surfaces in contact with the patient and their immediate environment – chair, couch, trolley, desk, horizontal surfaces etc.
- Ideally, non-essential items of equipment / furniture should be removed from the immediate environment during procedures such as wound dressings to minimise

environmental contamination. This is of particular importance with wounds colonised / infected with MRSA which is spread by both contact and airborne route on skin scales as well as contaminated dressings

- Prompt cleaning of any spillage of body fluids

TRANSPORT OF SERVICE USERS TO OTHER HEALTHCARE ENVIRONMENTS

On occasions, general practice staff may be required to refer a patient for additional healthcare to another provider e.g. for admission to hospital.

Receiving hospital staff must be informed of the potential infection risks prior to the transfer. This should be done at the time of making the referral. Ambulance service staff must be informed of potential / known infection risks so that they can make appropriate arrangements for transportation. All infection risks should be documented in handover documentation.

THIS PAGE HAS BEEN LEFT INTENTIONALLY BLANK

COLLECTION OF MICROBIOLOGICAL SPECIMENS

Please read in conjunction with:
Sample Transport Procedures for the Community

The document describes the procedure for transporting specimens to Barnsley and Rotherham Pathology Services from community premises or any other off-site location and is updated by Pathology Services.

www.barnsleyhospital.nhs.uk/pathology/pathology-general/transport-specimens-laboratory/

INFORMATION REQUIRED

Label the specimen container with the following essential information:

- Patient's Name (first name & surname)
- Date of birth
- NHS number/Hospital number
- Date collected
- Location/ward

In addition, the form should include specific information about the individual service user, including:

Service user's condition: e.g. immuno-compromised; if unusual organisms are suspected; RELEVANT clinical conditions; if part of a suspected outbreak.

Current medication and treatments: current or recent antibiotics, steroids or other immune-suppressive drugs, etc.

Source of specimen: particular body site; type of body fluid including method of collection e.g. MSU, catheter specimen of urine (CSU) etc. Wound swabs must specify the body area from which the specimen is obtained.

Purpose of the specimen: order investigations specifically required and avoid unnecessary tests, which add to budget costs. Discuss with Infection Control Advisor or laboratory prior to collection if necessary.

CONFIDENTIALITY

It is essential that confidentiality is maintained at all times and local arrangements must be in place to ensure sensitive information is not revealed unnecessarily on request cards. This is of particular importance with Blood Borne Viruses (BBVs) and Sexually Transmitted Diseases (STDs).

COLLECTION OF SPECIMENS

All laboratory staff are required to reject specimens that appear to be poorly collected or are inadequately labelled as results may be unreliable and misleading. Poorly collected specimens cause delayed results, results of little or no clinical utility, service user inconvenience, wasted time and increased costs. When collecting specimens observe the following:

- Whenever possible always take specimen prior to commencing antibiotics. If a course of antibiotics has started the specimen should be taken immediately prior to next dose and the antibiotics being given should be documented on the request form
- Provide an adequate quantity of material for examination as this will increase the chances of isolating the causative micro-organism. For example, stool container must be two thirds full.
- Collect fresh materials as free from extraneous contamination as possible and take material only from the required site (not surrounding tissues).
- Use an aseptic technique to avoid inadvertent contamination of the site of the sample or the specimen
- Prior to taking swabs from a dry area the swab should be moistened in sterile normal saline to improve adherence of bacteria
- Ensure that appropriate specimen containers are used especially if containing a transport medium. Using the wrong container may invalidate the specimen. Seek laboratory guidance if uncertain
- Secure lids immediately, to avoid spillage and contamination during transport.
- Write details on container prior to filling.
- Do not overfill containers especially faecal and urine containers. Over filling can increase the risk of leakage.
- All specimens must be placed in a specimen bag with the request form in a separate pocket. An additional 'Danger of Infection' label must be attached to specimens and request forms for known or suspected "high risk" pathogens. (See below)
- Always follow standard infection control precautions when handling specimens e.g. ensure appropriate protective clothing is used and ensure safe disposal of sharps.

SPECIMEN CONTAINERS

No change in the type of containers purchased or used should be made without discussion with the local laboratory.

The individual sending the specimen must ensure that an appropriate container is used, that it is securely closed and not externally contaminated.

HIGH RISK "BIOHAZARD" SPECIMENS (DANGER OF INFECTION)

All high risk specimens must have a danger of infection sticker attached to the request form and the sample.

If in doubt as to whether a specimen should be accompanied by a biohazard sticker, consult your local laboratory tel: 01226 432687 (Monday-Friday 08:30 – 20:00) or the Infection Prevention and Control Team tel: 01226 2825 (Monday-Friday 08:00 – 17:00).

High risk micro-organisms include:

Category 3

Human Immuno-Deficiency Virus (HIV)
Hepatitis B Virus (HBV)
Hepatitis C Virus (HCV)
Tuberculosis (TB)
Salmonella Typhimurium

Category 4

Viral haemorrhagic fevers
Rabies
Anthrax
CJD

STORAGE OF SPECIMENS

Any fridge that is used for the storage of specimens **MUST NOT** be used for the storage of any food items or drugs including vaccines. The fridge should have a min/max thermometer and be regularly cleaned and serviced.

Urine should ideally be examined in the laboratory within two hours. Otherwise, urine may be stored in the fridge for up to 24 hours. Bacteria will multiply at room temperature giving misleading results.

Sputum should be sent to the laboratory immediately as respiratory pathogens will not survive for prolonged periods.

Stools should be examined within twelve hours unless parasites are suspected when a warm fresh stool is required. Rectal swabs are only of value if they show the presence of faeces but stool specimens are preferred whenever possible. Stools for viral culture e.g. during an outbreak of diarrhoea and vomiting should reach the laboratory as soon as possible after collection as viral particles are rapidly killed.

High vaginal swabs should reach the laboratory within four hours.

Wound swabs should ideally reach the laboratory on the day they are taken. However, they can be stored in a specimen fridge overnight if this is not possible. Wound swabs must be collected using an appropriate transport medium e.g. Stewarts medium.

Do not leave specimens over the weekend or bank holidays even in the fridge.

TRANSPORT OF SPECIMENS

Pathology specimens are potentially infectious and hazardous. Care must therefore be taken to package and transport specimens in compliance with the Carriage of Dangerous Goods by Road (ADR 2017) to minimise the risks to staff, couriers and the general public. Specimens must be transported in an adequate leak-proof primary container; a leak-proof secondary container and an outer box to comply with British Standards

The collection and transportation of specimens is also important in ensuring the quality of results. If there have been any temperature excursions or delays in the transportation of the samples, this should be reported to laboratory reception staff on arrival.

All specimens must be placed in a specimen bag with the request form in a separate pocket.

All specimens should be placed in a designated, secure collection area until ready for collection.

Larger specimens such as 24 hour urine collections should be placed in clear plastic sacks which are tied at the neck. The request form should be attached to the outside of the bag. **DO NOT** use pins or staples to attach the form to the bag.

Specimens to be sent by post to specialist laboratories **MUST** be sent in packaging that conforms to the current transportation of dangerous goods regulations. Usually this is undertaken by the local medical microbiology laboratory and it may be necessary to send specimens there for transportation. Staff should liaise with the local laboratory undertaking diagnostic medical microbiology for further guidance. Under no circumstances must specimens be posted in packaging which does not conform to current regulations.

SPECIMENS CONTAINING RADIOACTIVE MATERIAL

If sending specimens from service users receiving therapeutic doses of unsealed radioactive sources seek advice from the laboratory staff before collecting the specimens.

Specimens from service users who have received tracer doses of radio- pharmaceutical products constitute no radiation risk and no special precautions need to be taken.

SPECIMENS CONTAINING CYTOTOXIC DRUGS

Specimens from service users receiving cytotoxic drugs may contain some unchanged drug or active metabolites. Specimens and request forms should be appropriately labelled and advice should be sought from the clinician/specialist nurse prior to transportation.

SPECIMENS AND CONTAINERS CONTAINING HAZARDOUS REAGENTS

If there is a hazardous reagent (e.g. liquid acid preservatives) present in the container, a hazard label should be attached by the laboratory.

RECEPTION OF SPECIMENS BY LABORATORY

If an unlabelled specimen is sent it will be discarded.

If damaged or unlabelled specimens are received by local laboratories, the recipient should, where possible, inform the sender that specimens have been discarded and should request further specimens are sent.

ACCIDENTS AND SPILLAGE

A bio-hazard spill kit will normally contain appropriate equipment and guidance for dealing with specimen spillage. These kits should be carried in vehicles used to transport specimens on a regular basis.

COLLECTION OF SPECIMENS – REFERENCE GUIDE (MARSDEN MANUAL OF CLINICAL NURSING PROCEDURES 9TH EDITION)

SITE	ACTION
Nose (anterior nares)	Relates to specimens for MRSA carriage. Prior to taking swabs from the nose, moisten with sterile water. One swab should be rotated around just inside both nostrils (do not swab further back into the nose).
Throat	One swab should make contact with one tonsil (or tonsillar fossa). The service user should stick out their tongue whilst the swab is guided down the side of the throat to make contact with the tonsil. A tongue depressor may be required. Do not make contact with any other area of mouth or tongue as this may cause contamination with other organisms.
Perineum	One swab should be rolled over the area between the genitalia and the anus (from front to back). Hygienic cleaning of the area should be undertaken if required prior to swabbing.
Groin	One swab should be rolled along the area of skin on the inner part of the thighs closest to the genitalia. Moisten with sterile saline beforehand.
Eye swabs	The exudate from the eye can be swabbed to identify some bacteria but others need to be identified by conjunctival scrapings which should be taken in an eye clinic. Hold swab parallel to the cornea and gently rub the conjunctiva in the lower eyelid from the nasal side outwards. If both eyes are to be swabbed a separate swab must be used for each eye.
Ear swabs	Ensure no antibiotics or other therapeutic drops have been used in the aural region 3 hours before taking the swab. Place the swab at the entrance of the auditory meatus. Rotate gently once.
Wounds/skin lesions	One swab should be rolled over the area. The wound may be irrigated with saline to remove surface debris before taking the swab if remnants of dressing remain. For large wounds, roll swab in a zig-zag motion to include all wound surface. If the wound is dry, moisten with 0.9% sodium chloride.
Catheter specimen of urine (CSU) 5-10 mls is required	Clamp tubing below rubber cuff (of catheter) to allow urine to collect. Urine specimens should only be taken from the sampling port using a sterile syringe +/- sterile needle (most manufacturers provide needle-less ports). Please note, using a needle increases the risk of sharps injury. Needles should not be used if it is a needle less port. Swab with 70% alcohol and allow to air dry prior to sampling. Aspirate the required amount of urine and remove the needle/syringe. Urine specimens must not be taken from the catheter bag as misleading results will be obtained due to bacteria having multiplied in the previously drained urine.
Mid-stream specimen of urine (MSU) 5-10 mls is required	Male - clean skin around prepuce (after retracting) with soap/water or normal saline. Female – part labia and clean with soap/water or normal saline (from front to back). Use separate swab for each wipe. The first and last part of the urine stream should be discarded and the mid-stream specimen collected into a sterile receiver and poured into a sterile container.
Stool/faecal specimens	Using the integrated spoon, scoop enough faecal matter to fill 2 thirds of the specimen container. Stool specimens can be obtained from a bedpan containing urine. This does not affect results. Only liquid stools (Bristol Stool Chart 6/7) will be examined for <i>C. difficile</i> toxins.
Vaginal Swabs	A sterile vaginal speculum must be used in order to separate the vaginal walls. The swab must be taken from as high in the vagina as possible.
Indwelling devices e.g. PEG site	One swab to be rolled over the area of skin surrounding the device. Pre- moisten swab with sterile water if necessary.

Pus	Pus may be collected using a sterile syringe and transferred into a sterile specimen container.
-----	---

INFECTIONS WITH SPECIFIC ALERT ORGANISMS

INTRODUCTION

This section is designed for professional staff including GPs who may provide healthcare to service users in residential care settings as well as in their own home and in local general practice facilities.

Safe Infection control practice requires knowledge of micro-organisms, the diseases they cause and how they spread between people.

To assist staff, the following list provides basic information on common infectious diseases; causative organism; mode of transmission and specific information relating to clinical care.

Staff should consult the following list to determine the risk posed to others and how to manage service users safely.

Some infections are caused by an individual's own micro-organisms. This is called Endogenous Infection.

Cross infection where micro-organisms have been transmitted between individuals are called Exogenous Infections.

Advice may be sought from the local IPC Team / clinician / GP / PHE on the management of service users or service users' household contacts with these infections.

NOTIFICATION OF INFECTIOUS DISEASES

Public Health Regulations require statutory notification of certain infectious diseases. Notification is the responsibility of a Registered Medical Practitioner. See section IPC Management Policy

INFECTIOUS (COMMUNICABLE) DISEASES - Standard Infection Control Precautions (SICP)

Disease	Mode of Transmission	Comments and Precautions
Acquired immunodeficiency syndrome (AIDS) <i>See sub-section on Blood Borne Viruses (BBV)</i>	By direct contact with infected blood, sexual transmission and vertical transmission (mother to baby)	SICP when in contact with blood or blood stained body fluids. Safe spillage management Safe sharps management
Candidiasis (thrush)	Endogenous spread	SICP. May indicate immunosuppression or recent antibiotic therapy.
Chickenpox (Varicella)	Respiratory droplets and direct contact with vesicle fluid	Incubation period 14 – 16 days Highly infectious until lesions are dry. Potentially harmful to non-immune pregnant women and the immunocompromised. Suspicion of Chicken Pox <i>in a member of staff</i> must be reported immediately to the CCDC (at local HPU) even at weekends. Respiratory isolation.
Chlamydiosis	Sexual transmission	No restrictions
Cholera	Ingestion of contaminated food or water	Incubation period 1- 3 days SICP
<i>Clostridium difficile</i> <i>See separate section</i>	Faecal-Oral This bacteria produces spores which can live in the environment for months or years and requires chlorine releasing disinfectants to destroy	Enteric Isolation (own toilet facilities) Environmental cleaning with chlorine releasing disinfectants. Do NOT use alcohol hand rub as less effective than soap and water. Service users with active <i>Clostridium difficile</i> associated disease may be cared for in hospital but may present in general practice
Cold sores	Contact with vesicle fluid, saliva, sexual contact	SICP especially hand hygiene and glove use

Disease	Mode of Transmission	Comments and Precautions
Common cold	Respiratory droplets and contact spread	Incubation period 1 – 3 days SICP
Cryptosporidiosis	Faecal-oral route	SICP
Cytomegalovirus	Direct contact,	SICP
Diarrhoea	Faecal-oral route, contact spread	SICP. Soap and water for hand hygiene (NOT alcohol rub). Environmental cleaning of toilet facilities after use with chlorine releasing disinfectant
Fifth disease (Erythema infectiosum)	Respiratory secretions (saliva, mucus) Also from infected blood	Also known as Slapped Cheek Syndrome Incubation period 4 – 21 days Infectious before rash appears Women in first trimester at risk of serious complication; also those with sickle cell disease and the immunocompromised
Glandular fever (Infectious mononucleosis)	Contact spread with saliva	Incubation period 28 – 42 days Infection may be transmitted on hands of staff if contaminated with saliva. Good hand hygiene essential
Hand foot & mouth disease (Coxsackie A / Enterovirus 71)	Direct contact with nose and throat secretions; faeces	Incubation period 3 – 7 days Strict attention to hand hygiene Immunocompromised at risk. Severe complications uncommon (neurological mainly) Outbreaks common in nurseries / schools
Hepatitis A	Faecal-oral route	SICP
Hepatitis B <i>See sub-section on Blood Borne Viruses (BBV)</i>	Direct contact with infected blood, sexually transmitted	SICP when in contact with blood or blood stained body fluids. Safe spillage management. Safe sharps management Immunization of all staff in contact with blood.

Disease	Mode of Transmission	Comments and Precautions
Hepatitis C <i>See sub-section on Blood Borne Viruses (BBV)</i>	As above	SICP when in contact with blood or blood stained body fluids Safe spillage management Safe sharps management Avoid contact with lesions
Impetigo	Direct contact with lesions	Young children often highly susceptible Attention to hand hygiene essential Avoid contact with lesions
Infestations <i>See Sub Section below</i>	Body Lice, Hair Lice, Scabies	SICP
Influenza	Respiratory (droplet) and contact transmission	Incubation period is 1-4 days. Transmission risk continues for 3-7 days or until the patient is asymptomatic. At risk patient groups should be immunised according to published guidance – see separate vaccination section. Influenza can cause outbreaks in residential care settings, the HPA may request swabbing of affected residents and ant-viral treatment/prophylaxis. Advice should be sought from HPA. May cause mild self limiting disease however unvaccinated at risk patients may experience severe disease.
Legionnaires' disease	Inhalation of contaminated aerosols	Not spread from person-to-person.
Measles	Respiratory droplets	Incubation period 9 – 12 days Potentially hazardous to the very young (under 1 year), immune-compromised people or non-immune pregnant women. Respiratory isolation.
MRSA <i>See sub-section on MRSA</i>	Contact spread	SICP, strict attention to hand hygiene and principles of asepsis when caring for invasive devices or wounds.
Mumps (pertussis)	Respiratory droplets, direct contact with saliva	Incubation period 7 – 14 days Infectious prior to onset of illness. SICP Respiratory isolation.

Disease	Mode of Transmission	Comments and Precautions
Norovirus	Spread via airborne and contact routes in vomit and faeces	Incubation period 1 – 2 days Usually self-limiting but can cause severe dehydration in infants / elderly Enteric / contact isolation SICPs
Poliomyelitis	Faecal-oral route, direct contact with nasal or oral secretions	Incubation period 7 – 14 days Strict attention to hand hygiene.
PVL <i>Staphylococcus aureus</i>	Contact transmission	Some strains of <i>Staphylococcus aureus</i> (both Meticillin resistant and sensitive) produce Panton Valentine Leukocidin, a toxin which is a virulence factor. Strains can cause skin and soft tissue infections which commonly recur. Rarely this causes severe invasive disease e.g. necrotising haemorrhagic pneumonia. PVL should be suspected in patients presenting with recurrent skin infections e.g. boils. Swabs should be taken and PVL suspicion noted on the request. Positive results should be notified to HPA. who will advise on contact tracing and
Resistant Organisms e.g. VRE, ESBLs/Gram negative enterococci	Many bacteria are developing resistance to antibiotics in addition to MRSA. These include some strains of normal gut flora which can be spread by direct or indirect contact	SICP and application of principles of asepsis.
Rubella (german measles)	Direct contact with respiratory secretions or droplets.	Incubation period 14 – 21 days Potentially hazardous to the very young (under 1 year), immune-compromised people or non-immune pregnant women. Respiratory isolation.
Salmonella	Food-borne – ingestion of contaminated food. Faecal-oral transmission.	Transmission can occur via food handling by infected individual. Enteric precautions (own toilet facilities).
Scabies See sub-section on Parasite infections	Prolonged skin-to-skin contact	Norwegian scabies highly infectious. Dermatology diagnosis recommended. SICP apply. Staff contacts may need treatment.

Disease	Mode of Transmission	Comments and Precautions
Shingles (Varicella Zoster)	Direct contact with lesion exudate	<p>Shingles can occur in people who have had chicken pox when the virus reactivates in sensory nerve cells. People not immune to chicken pox can acquire this from individuals with shingles</p> <p>Keep lesions covered. Strict attention to hand hygiene and glove use when in contact with lesions. Infectious until lesions dry.</p>
Tuberculosis <i>See sub-section on Tb</i>	Inhalation of airborne droplets	Pulmonary disease infectious until after 2 weeks of treatment. Infections at other sites are not normally infectious. Respiratory isolation for first two weeks of treatment.

MRSA

INTRODUCTION

Staphylococcus aureus is an organism which approximately one third of the population carries on their skin and in their nose and throat without any associated problems; this is known as colonisation.

Although *Staphylococcus aureus* is capable of causing infection, most infections are easily treated with antibiotics.

Some strains of *Staphylococcus aureus* have developed resistance to some of the more commonly used antibiotics; these are known as Meticillin Resistant *Staphylococcus aureus* (MRSA). Meticillin is an antibiotic that is closely related to Flucloxacillin.

MRSA WITHIN THE COMMUNITY

Service users within the community setting are at a lower risk from MRSA acquisition as they live in their own environment and less likely to come into contact with other colonised individuals.

With earlier hospital discharge and an increase in the number of service users receiving healthcare in community environments, levels of MRSA acquisition in these environments is rising.

High Risk Service Users

Service users at higher risk of carriage of MRSA include those who are:

- Have an indwelling device, e.g catheter,
- Parental enteral feeding tube (PEG),
- May live in a care home where they come into contact with other users who have a higher colonisation rates.
- May have a problematic wound or a Venous Ulcer.
- May have a history of IV drug abuse, which makes them more vulnerable to develop infections.
- Known to be previously infected or colonised with MRSA
- Frequent re-admissions to any healthcare facility

PREVENTION OF SPREAD

The single most important factor is effective hand hygiene. All staff who have direct contact with the service user or their immediate environment must wash their hands thoroughly using liquid soap and disposable paper/clean hand towels before and after contact. Alternatively, an alcohol hand gel can be used for hand decontamination. Gloves and

Aprons must be worn when handling dressings, infected wounds or specimens.

SERVICE USER SCREENING

The Department of Health has published guidance requiring routine screening of most service users admitted to acute hospitals. This is usually undertaken at the time of admission (emergency admissions) or at pre-assessment clinic (routine, planned admissions). There is no requirement for routine screening in general practice.

STAFF SCREENING

There is no need for routine screening of staff in any healthcare setting.

MANAGING MRSA POSITIVE SERVICE USERS

Standard Infection Control Precautions should be applied with strict attention to the principles of asepsis when caring for invasive devices or undertaking an invasive procedure e.g. wound care.

There is no requirement to routinely decolonise service users with MRSA in general practice, unless they sit in the above high risk criteria. Those individuals sent home from hospital whilst undergoing decolonisation should continue with prescribed treatment. It is advisable for one post treatment screen of these individuals to determine effectiveness of treatment/clearance and reduction of microbial load.

Further advice can be obtained from the IPCT

BLOOD BORNE VIRUSES

INTRODUCTION

Viruses transmitted by blood and blood-stained body fluids are of particular importance to healthcare workers who may be at risk of acquiring infection during the course of their work. The most significant route of spread (in occupational exposure) is via contaminated sharps.

The most important blood borne viruses are:

- Hepatitis B
- Hepatitis C
- HIV

This should be read in conjunction with other policies within this manual:

- Safe handling and disposal of sharps
- Management of Occupational Exposure to Blood Borne Viruses
- Decontamination of Medical Equipment
- Spillages of blood and body fluids

Hepatitis B (HBV)

Most infections caused by this virus are mild, however in a few cases extensive liver damage and liver failure may prove fatal

Between 2-10% of those infected do not completely eliminate the virus and become chronic carriers.

Some groups are at increased risk of acquiring HBV; these include but are not restricted to:

- Service users receiving renal dialysis
- Haemophiliacs
- Intravenous drug users who share needles
- Families of chronic carriers
- Residents of institutions whose behaviour may facilitate transmission e.g. biting

HBV is transmitted by sexual intercourse, perinatally from mother to baby, via inoculation when infected body fluids are inoculated through the skin; on instruments, via damaged skin or through splashing contact with mucous membranes.

HBV has been isolated from almost all body fluids. However, the following body fluids are those most implicated in the transmission of the virus:

- Blood
- Semen
- Vaginal fluids

Healthcare workers are at risk of acquiring HBV from sharp injuries, scratches, bites and from body fluid splash incidents. The virus must have contact with the staff member's blood to cause infection. The virus can live outside on equipment, but only for a limited amount of time.

Hepatitis C (HCV)

Primary symptoms of infection with HCV is often mild, asymptomatic and rarely associated with jaundice. 85 % of those infected become chronic carriers.

The infection has been transmitted by blood transfusion, although in developed countries this has been eliminated by the introduction of blood donor screening.

Transmission is also possible by the following routes:

- Intravenous drug users who share needles
- Sexual intercourse

- Perinatally from mother to baby

Healthcare workers are at risk of acquiring HCV from sharp injuries, scratches, bites and from body fluid splash incidents. The virus must have contact with the staff member's blood to cause infection. The virus can live outside on equipment, but only for a limited amount of time.

Human Immunodeficiency Virus (HIV)

Infection with HIV will persist indefinitely once seroconversion has occurred. Infection can be transmitted to others from an infected individual soon after they have acquired the infection (when the virus is replicating rapidly) but becomes more infectious as immunodeficiency decreases and the amount of virus in the blood increases.

HIV is transmitted via the following routes:

- Sexual intercourse
- By inoculation of infected body fluids
- Through damaged skin or on to mucous membranes e.g. conjunctivae, mouth
- By transfusion of contaminated blood
- Perinatally via the placenta, during delivery and from breast milk

The greatest concentration of virus is found in blood or body fluids containing visible blood.

Occupational transmission of HIV has been reported. Healthcare workers are at risk of acquiring HIV via the following exposures:

- Inoculation of infected blood / body fluid into body tissues by a needle or other sharp device.
- Splash incidents with infected blood / body fluids into the eyes or mouth or through damaged skin.

Service users are at risk of acquiring HIV via inadequately decontaminated medical devices. The virus must have contact with the staff member's blood to cause infection. The virus can live outside on equipment, but only for a limited amount of time.

Managing Service users with Blood-borne Viruses

Service users with blood-borne viruses do not require any additional infection control precautions beyond SICPs.

Protective clothing is necessary only for direct contact with blood or body fluid. Specimens should be labelled as 'high risk' as per specimen collection policy.

With new treatments available, patients who are compliant with treatment can reduce microbial load significantly. This will reduce the risk of cross contamination, but only if the service user is compliant with treatment.

Reducing the Risk of Occupational Exposure

Injury with a contaminated sharp device / instrument is the most likely route of transmission to a care worker, therefore all staff should be aware of safe working practices when handling and disposing of used sharp instruments.

Staff should be made aware of the action to be followed in the event of accidental sharps injury or splash incidents. Staff are advised to wash it, bleed it, cover it and report it.

Staff should adopt standard infection control precautions for all service users regardless of the perceived risk of infection.

TRANSMISSIBLE SPONGIFORM ENCEPHALOPATHIES

INTRODUCTION

Transmissible Spongiform Encephalopathies (TSE's) are rare, chronic fatal degenerative brain diseases of humans and certain other animal species, whose natural mode of transmission is unknown but is thought to be passed by inoculation and in some cases by ingestion of high risk tissue.

The infecting agent is of virus size but is an unconventional protein known as a prion, which replicates extremely slowly with an extended incubation period. The infectious agent is thought to be restricted to the central nervous system and lymphoid tissue. There is now some suggestion that some cells in the blood may also be affected. There appears to be no antibody or other immune response to the infection making it difficult to detect. Diagnosis is by clinical signs and symptoms and characteristic changes in the brain.

These agents are very resistant to both heat and chemical disinfection and could therefore pose a potential risk to staff and service users via contaminated surgical instruments. Stringent management arrangements are required for the re- processing of certain types of surgical instruments and other medical devices that may potentially have been contaminated with TSEs. Such instruments are unlikely to be used routinely in general practice. Some clinicians undertaking enhanced services (dependent on type of procedure) *may* need to take additional precautions with surgical instruments and should seek advice from local Trusts' Decontamination Lead.

There have been no confirmed cases of TSE in health care staff as a result of occupational exposure to an infected service user.

Standard infection control practice should be routinely followed as described in this infection control policy.

Known or suspected TSE service users

Includes those service users with:

- An established diagnosis of classical sporadic Creutzfeld-Jakob Disease (CJD), variant CJD (vCJD), Gerstmann-Straussler-Scheinker Syndrome (GSS), Fatal Familial Insomnia (FFI) or Kuru.
- Service users suspected of having a TSE or a related disorder whose clinical symptoms are suggestive of TSE.

At risk service users

Rarely, asymptomatic service users may be identified as potentially at risk of developing TSE related disorders. This may be iatrogenic in origin and such patients will have been notified.

Care of known, suspected or at risk service users

In general practice Standard Infection Control Precautions are usually all that is required for the care of these service users. Specific advice on individual affected service users should seek advice from local clinicians or HPA.

Additional information on CJD risk assessment and management can be found at

<http://www.dh.gov.uk/ab/ACDP/TSEguidance/index.htm?ssSourceSiteId=en>

TUBERCULOSIS

INTRODUCTION

Tuberculosis (TB) is an infectious disease caused by *mycobacterium spp.* There are a number of bacteria within the Mycobacteria family and these are widely distributed throughout the world but only a few species cause disease in man. The most common tuberculosis infections are caused by *Mycobacterium tuberculosis* although *Mycobacterium avium intracellulare* (or MAI or Mycobacterium complex) is commonly seen in immunocompromised service users.

The most common primary site of infection is the lungs (pulmonary tuberculosis), but bones, joints, the brain and meninges and other internal organs may be affected.

TB infections in sites other than the lung are not normally infectious. Some infections with TB remain dormant (and non-infective) for many years; this is called latent TB Infection. Such service users may develop active infection later in life, often following debilitating illness.

It is not easy to become infected with TB but risk is increased through prolonged close contact (a cumulative period of 8 hours is considered sufficient) with an infected case (e.g household contact) and even then only 30% of healthy people will become infected and of

those only 5-10% will develop active disease. Those more at risk of acquiring disease are:

- Household contacts
- Those living in unhealthy or overcrowded environments
- Prolonged exposure in a country with high incidence of TB
- The very young or elderly
- Immunosuppressed (such as with HIV)
- Those with a history of drug or alcohol misuse or detainment in prison
- The children of parents whose country of origin has a high incidence of TB

Casual contacts such as work colleagues and friends are not considered to be at increased risk. The TB specialist team will, however, make an informed decision on whether to screen work colleagues or friends depending on level of contact and level of infectiousness of the case.

Those affected with pulmonary TB most commonly present with a persistent cough, weight loss, severe night sweats, tiredness and some may present with haemoptysis.

Incubation is most commonly 4 to 12 weeks. For some people the infection will lay dormant (latent) but they may, however, develop active disease later in life if predisposed with another debilitating illness.

Sputum Specimens

3 sputum specimens are required to determine the presence, or absence, of mycobacterial type organisms (AFBs). It is important to ensure the specimen consists of purulent secretions coughed up from the bronchi and is not merely saliva from the mouth. Suitable specimens are best collected during a bout of coughing soon after the service user wakes in the morning.

Care must always be taken with specimens to ensure that the lid of the container is secured tightly and there is no trace of the specimen on the outside of the container. A biohazard label must be attached to the specimen and the request form.

A positive AFB result indicates that someone has tuberculosis, but does not confirm active TB disease. A full culture and sensitivity testing is required which may take several weeks.

Appropriate precautions to prevent the spread of infection

Most service users with tuberculosis are treated at home; a few need hospital admission for severe illness, adverse effects of chemotherapy or social reasons.

Special precautions, other than care in disposal of infected exudates, are only required when the service user has confirmed or suspected open pulmonary tuberculosis i.e. AFB positive service users. Such service users should not attend general practice during the first 14 days of drug therapy unless segregated during their visit. See section 18.

Care protocol for first two weeks

Once a person has been commenced on treatment for suspected tuberculosis, they generally remain infectious only for the first two weeks of treatment unless there is a suspicion of a resistant or multi-resistant strain of tuberculosis or are non-compliant with treatment. In a healthcare environment they will require single room isolation with mechanical ventilation during this period.

The service user will usually remain at home as directed by the Chest Physician or TB Nurse Specialist with restricted visitors until confirmatory diagnosis and up to fourteen days of specific chemotherapy has been completed (advice will be given by the TB Nurse Specialist).

Care protocol throughout the service user's illness

Encourage service user to cough into disposable tissues which are then disposed of immediately. If in a healthcare setting, they should be placed into a clinical waste bag.

Encourage the service user to expectorate into a sputum pot which is kept covered and disposed of on at least a daily basis. This must be disposed of with a secure lid. If in a clinical setting, it should be placed into a clinical waste bag.

NB Service users generating significant volumes of respiratory waste *may* require a clinical waste collection service setting up. See Section 10.

Ensure the service user turns their head away from others when coughing or expectorating.

Ensure the service user undertakes effective hand washing particularly before meals and after coughing.

There is no need for separate crockery or cutlery.

Well ventilated accommodation is preferable e.g. open windows when weather conditions allow.

Medication compliance is essential and will require supervision by the TB Nurse Specialist when the service user is cared for in the community. Early referral to the TB team is therefore essential. Failure to comply with treatment may cause complications and will encourage the development of MDRTB and increase infectivity. Service users in community settings undergoing treatment for TB will have their treatment monitored by the TB specialist team who will provide advice to carers on ensuring compliance.

If a service user dies with active tuberculosis the body should be placed in a body bag, and clearly labelled as a risk of infection to alert the mortuary/undertakers staff to the risks.

Multi-Drug Resistant Tuberculosis (MDRTB)

There are now increasing numbers of service users being identified with a tuberculosis infection which is resistant to more than one of the usually prescribed drugs used for treatment. Once identified these service users are usually cared for in hospital until they are no longer infectious. However it is possible that general practice staff may have had regular contact with the service user prior to admission. In such cases, guidance will be provided by the local Health Protection Unit in collaboration with the local Chest Physician / TB team and Trust Infection Control Advisor.

Staff protection / contact tracing

Staff that are caring for service users with open pulmonary TB within the first two week of treatment are not usually required to wear masks. Close fitting FFP3 masks are required when undertaking sputum producing procedures. If masks are required for any other healthcare activity this would be directed by the TB Specialist team.

All close staff contacts of sputum smear positive (AFB positive) service users will be checked and followed up by the TB nurse specialist as appropriate. A contact list will be compiled in conjunction with the local Trust Infection Control Advisor and the local Health Protection Team

Any employee of the organisation who develops an illness suggestive of tuberculosis should seek medical advice from their own GP as soon as possible.

LICE

ECTOPARASITES

Head Lice, Body Lice, Pubic Lice and the Scabies Mite

Lice live on the skin or inner layers of clothing. Once parted from their host, they soon die, although the nits or eggs may remain viable for long periods. Transmission is by contact either with the hair (head or pubic lice) or clothing (body lice) of the host.

Head Lice (*Pediculus Humanus Capitis*)

The adult louse is approximately 3mm long and lives for about 20 days. The female head louse produces on average 56 eggs after a single insemination, at the rate of approximately six eggs per day. It feeds on human blood. Bites cannot be felt but repeated bites lead to sensitisation and irritation (itching) of the scalp. Irritation to the scalp is also due to an allergic reaction to louse faeces. Once the infected person is sensitised to the bites the itch is continuous. The eggs, which are difficult to see, are glued to individual hairs just above the roots and are tear shaped and approximately 1mm long. They hatch after 7-11 days and reach adult stage within 6-12 days. The empty egg shells (nits) are white and shiny and are harmless. As the hair grows the empty egg shells can be found further along the hair shaft.

The live lice are transmitted by prolonged head to head contact, which must be for at least 30 seconds. Lice cannot jump or fly but crawl quickly in dry hair from one head to another.

Diagnosis

Diagnosis is by identification of a live moving louse on the hair which is most effectively done by the wet combing method (described below). Children aged 4-11 years are the most frequently affected so it is important for control measures that families check their hair for infection regularly and treat appropriately.

Wet combing detection method

Wash the hair in the normal way.

Using a fine-toothed comb and lots of conditioner, firstly comb the tangles out of the hair over a pale surface or paper towel. Clean the comb between each stroke using a piece of tissue. Then repeat the process with a fine-toothed comb, combing a small section of hair from the roots to the end and cleaning the comb after each stroke. Examine the tissue after each combing for traces of lice or eggs. After completing the combing, rinse and dry the hair in the usual way.

If live lice are identified, then an appropriate eradication method should be used.

If lice are found then all other close contacts should be checked for infestation by use of the wet combing method and **only those who are found to be affected should be treated.**

Treatment - Insecticides

There are four main types of insecticide treatments available:

- Carbaryl
- Malathion
- Synthetic pyrethroids, phenothrin and permethrin
- Dimeticone which is not a chemical but works by immobilising the lice

Alcohol based treatments must not be used on babies or people with asthma, when a water based treatment must be used.

Staff should wear a plastic apron and gloves while carrying out the treatment.

Apply lotion according to the instructions and rub gently into scalp, avoiding contact with the eyes. Repeat until hair is thoroughly wet. Allow hair to dry naturally.

After the recommended contact time wash hair with normal shampoo, rinse using lots of conditioner. While the hair is wet, comb with a fine toothed comb, making sure that the

teeth of the comb slot into the roots of the hair every time to remove lice and nits. Clear the comb after each stroke.

An insecticide treatment should be repeated seven days later. This is because the insecticide is not 100% effective at eradicating all the eggs which may then hatch during the following seven days. The second application ensures that the nymph stage lice (young lice) are eradicated before being able to lay eggs.

2 to 3 days after the second application of the insecticide the hair should be combed through with a detection comb. If any adult lice are found this is either due to treatment failure or re-infestation. In either case a second choice of insecticide should be chosen so as to prevent resistance to the treatment occurring. This is called the mosaic approach to treatment.

Wet-combing eradication method

Wet combing should be performed on days 1, 5, 9 and 13 over a fourteen day period and should follow the same method as the eradication method described above. It is important that, between each stroke, lice are cleaned from the comb and that the entire head and length of hair is checked during the process.

Period of communicability

Until case is treated

Exclusion

None

Body Lice

The adult body louse is larger than the head louse and also feeds on human blood. It is associated with poor living conditions, lack of cleanliness and lack of adequate nutrition. The presenting signs are pinpoint lesions, excoriation and pigmentation of the skin. Eggs are laid on the clothing of the host, in the lining, seams and underwear and occasionally on the body hairs. The body louse may be transferred by direct contact, but more often by wearing infested clothing or sleeping in infested bedding.

Treatment

Treatment does not usually require pesticides.

Body lice are seldom found on the skin after clothing has been removed. The louse only transfers in the dark therefore remove clothing in a well lit room.

It is recommended that staff wear gloves and a plastic apron while assisting service users.

Collect clothing and bed linen in water-soluble linen bags.

Clothes should be turned inside out and tumbled dried at 50⁰ C for 30 minutes. This will be sufficient to kill both lice and eggs. Clothes can then be washed in the usual way.

No special environmental measures are required.

The crab louse is generally found in the pubic and perineal region, but may also be in the armpits, hairy chests, beard, eyebrows and eyelashes. It is more firmly attached and less likely to transfer to healthcare staff. It is normally acquired by intimate contact. It feeds on human blood and can be seen as a dark red spot. Bites cannot be felt but irritation occurs and blue/grey skin lesions can be seen.

Treatment

Treatment is with Malathion or Carbaryl lotions or shampoo. An aqueous based lotion should be used on the genital or other areas as necessary.

Clothing should be washed and ironed.

Staff should wear gloves if required to carry out the treatment.

Sexual partners should be treated simultaneously whether infection is confirmed with them or not.

SCABIES

Infection with the scabies mite is currently increasing and there have been a number of cases of resistance to the usual treatments. Extended direct contact (i.e. skin to skin for 3-5 minutes) is required for transmission of the mite.

Scabies is an infestation of the skin by the microscopic mite *Sarcoptes scabiei*, which burrows into the skin. These burrows are often visible as a discoloured, raised line, which may be straight, tortuous or dotted on the wrists, back of the hands and between the fingers.

Infection with scabies presents with intense itching caused by an allergic reaction to the faeces of the mite. The burrowing itself may also cause itching. The mite tends to burrow into warm skin creases so elbows, armpits, beneath the breasts, waist, groin, genitalia, buttocks, knees and ankles are often affected.

Infection with the scabies mite is very difficult to detect until the infested individual becomes allergic to proteins in the excreta of the mite which takes from 2-6 weeks. This causes increasingly intensive itching particularly at night. There are two particular types of scabies to note:

- Classical scabies which presents in otherwise healthy individuals. There are few mites present and few associated complications.
- Norwegian scabies (also known as crusted scabies) which can occur in those with impaired immunity. Infestation is with large numbers of mites, reaching possibly thousands and affecting the entire body. Typical burrows may not be seen and the service user may present with a rash resembling a chronic dermatitis. The classical itch may be absent. This form of scabies is highly infectious and can cause environmental contamination.

Transmission of scabies infection occurs during very close skin-to-skin contact with an infected individual and spreads rapidly under crowded conditions where frequent skin-to-skin contact is unavoidable such as in hospitals, care homes and childcare facilities.

As many elderly people are affected by dry skin it is often extremely difficult to diagnose scabies infestation in the elderly. Referral to a dermatologist for confirmation of diagnosis is often the most effective method of determining an accurate diagnosis particularly if other treatment regimes have failed.

Treatment

Those diagnosed with scabies, as well as their sexual partners and any other close contacts that may have had close prolonged contact within the preceding 6 weeks should be treated. These treatments should be given concurrently on the same day.

A malathion or permethrin based lotion are the current treatments of choice.

Instructions for treatment

- Always follow manufacturers' guidance which will be included in packaging.
- The lotion or cream should be applied from the chin downwards. All areas of the body, including genitalia, must be treated, except for the face and neck. It should be left on for the instructed length of time, after which the service user should bath or shower.
- Any cream washed off during the course of treatment should be re-applied until the treatment time has elapsed.
- Those who are infected will need to receive a second treatment 3-5 days later. Unaffected contacts will only need to receive one treatment.
- All bed linen and clothing worn just before treatment must be washed on a high temperature. If items are not washable then they should be ironed with a hot iron.
- It is important to note that itching may persist for several weeks after treatment. Antihistamines may be recommended to reduce itching.
- Further medical attention should be sought if itching persists after 4 weeks.

The two most common causes for treatment failure are:

- Failure to treat all contacts simultaneously so the chances of re-infection are increased.
- Failure to re-apply the treatment during the treatment phase after washing hands.

A hot bath before treatment is NOT recommended. If the service user is dirty a cool bath may be given, and treatment should then be delayed for at least two hours following the bath. Bathing before treatment increases absorption of the lotion into the bloodstream and away from the skin area which requires treatment.

Staff should wear gloves and plastic aprons for direct contact during treatment.

Expert advice should be sought for the treatment of crusted (Norwegian) scabies as in some rare cases systemic treatments may be necessary.

Recommendations

Staff infected outside the healthcare environment should be excluded from work until 24 hours after completion of the treatment.

Staff infected as a result of occupational exposure from service users they are caring for may return to work after treatment but should not work elsewhere until 24 hours after treatment.

Visitors should be discouraged from close contact with the service user/client until 24 hours after completion of treatment.

Service users should not visit Day Units, Lunch Clubs, Occupational Therapy units etc. until treatment is completed.

If an admission to hospital is required, the receiving department must be informed of the diagnosis and treatments already given.

Seek guidance from PHE or IPCT, if there is the likelihood of more than one case of scabies i.e. an outbreak.

MANAGEMENT OF CLOSTRIDIUM DIFFICILE DIARRHOEA

INTRODUCTION

Clostridium difficile is a gram positive, anaerobic, spore forming bacillus. Approximately 3-5% of the adult population and a larger percentage of infants carry the bacteria in normal flora without disease. This percentage may increase in hospitalised patients and there is evidence of increased carriage in residents of care homes. Once established the bacteria produces toxins which cause the diarrhoea and damage cells lining the bowel.

Clostridium difficile spores shed by affected patients with diarrhoea readily contaminate the environment and can survive for long periods. This has led to outbreaks in hospitals and care settings.

Infection and disease can occur if patients are exposed to the spores and the normal flora is disrupted; thus disease acquisition is a two stage process.

National initiatives to reduce the incidence of this infection in hospitalised patients have had considerable success. Whilst this work continues there is a growing focus on prevention and management of *Clostridium difficile* in community and primary care settings. National initiatives include the setting of annual trajectories of cases of CDI. These trajectories are set for all acute NHS trusts *and since 2011/12 trajectories have also been set for Primary Care Organisations and latterly for Clinical Commissioning Groups (CCGs)*. Trajectories are monitored regularly by NHS England and other commissioners of care. Financial penalties are incurred if trajectories are breached. Toxin positive cases of CDI occurring in samples sent from GP practices are included in the local CCG trajectory.

Expert guidance is available on the management of CDI:

Department of Health & PHE Health Protection Agency (2009): *Clostridium difficile* Infection – how to deal with the problem DH.

This guidance was partially updated in 2013 in line with changes to the evidence base for treatment. All local prescribing policies are expected to comply with the treatment options published in this guidance.

SCOPE OF POLICY

These guidelines should be read and followed by all clinical staff in practices.

SYMPTOMS AND SIGNS OF CLOSTRIDIUM DIFFICILE INFECTION

Symptoms range from mild diarrhoea to explosive severe watery diarrhoea with blood and mucus in the stool which may be green. The diarrhoea may be frequent (as many as 30 times per day) and commonly has a distinctive foul odour.

Clinical signs and symptoms of abdominal disease may be present with pain, fever, nausea and bloating. CRP may be raised and neutrophil leucocytosis may be present. In severe cases damage to the lining of the gut may occur with the presence of Pseudomembranous colitis. Toxic megacolon may occur. Severe cases may require colectomy and may result in death.

PATIENT RISK FACTORS FOR CLOSTRIDIUM DIFFICILE DISEASE

Most cases of this infection occur in the over 65 year age group however disease may be found in younger adults. As stated above, disease may follow both the acquisition of spores and the disturbance of normal gut flora.

Previous antibiotic therapy is associated with most cases. In particular, the use of 3rd generation Cephalosporins and Quinolones and repeat or 'cocktail' courses of broad spectrum antibiotics. All routine antibiotic prescribing in general practice should be in line with local formulary. Patients receiving antibiotics are 'at risk' of acquisition of *C. difficile* for approximately 6-10 weeks as this is the time required for normal flora to become re-established.

Additional risk factors include the administration of Proton Pump Inhibitors, the presence of naso-gastric tubes and underlying bowel disease/non-surgical bowel interventions.

Debilitated or immune compromised patients are particularly vulnerable to severe disease if they are exposed to the spores and normal flora is disturbed.

SAMPLING AND DIAGNOSIS

Expert guidance on sampling has been issued: Department of Health (2012)
Updated DH/ARHA guidance on the diagnosis and reporting of C. difficile

Patients presenting with watery diarrhoea should be assessed for indications of *Clostridium difficile* disease. Normal bowel patterns and alternative causes of diarrhoea should also be considered. Cases may be identified where the bacteria is present as a colonising organism with diarrhoea from alternative causes.

If *Clostridium difficile* disease is suspected a stool sample should be taken and sent to the microbiology laboratory. Only samples of Bristol Stool Grade 5-7 should be sent i.e. stools that take the shape of the container. The request should be for MC&S and *Clostridium difficile* toxin. Samples contaminated with urine may be sent.

Diagnosis is usually made by detecting the presence of toxins using EIA. Secondary tests (GDH/NAAT/PCR) may also be undertaken.

TREATMENT

Cases of *Clostridium difficile* infection should be assessed for severity of disease as per published guidance. The table describing this is reproduced below.

Severity	Stool frequency per day	Stool type	Inflammatory markers	Vital signs	Other features
Mild	>3	5-7			
Moderate	3-5	5-7	WCC $<15 \times 10^9/L$ CRP <150		
Severe	Unreliable indicator		WCC $>15 \times 10^9/L$ acute rising serum creatinine $>50\%$ above baseline	Raised temp. >38.5 .	Evidence of severe colitis
Life threatening	Unreliable indicator			Hypotension	Partial or complete ileus or toxic megacolon

Treatment algorithms are as advised in published guidance. Advice may be sought from the Consultant Medical Microbiologist (CMM). In some cases symptoms may continue despite treatment or patients may relapse after resolution of symptoms. The algorithm includes management of initial episodes and also for recurrent disease.

Wherever possible antibiotics prescribed for other infections should be stopped. Advice may be sought from the Community Medicines Management or Microbiologist if required. Where continuing antibiotic therapy is required this should be assessed on a daily basis. In some cases stopping antibiotic therapy may lead to cessation of diarrhoea symptoms within 48 hrs.

Where diarrhoea is severe and frequent, particularly in the elderly patient, dehydration with electrolyte imbalance may follow. Supportive therapy and blood tests may be required.

Antimotility agents should not be prescribed and patients should be advised not to purchase these over the counter.

The use of PPIs should be reviewed and, if possible, discontinued. Surgical, gastroenterology and nutritional advice may be required. Samples should not be sent to demonstrate clearance. Patients will commonly shed bacteria and toxins for many weeks. Laboratories will not test samples sent within 28 days of a positive result - in line with published guidance.

Where relapse is suspected it is not usually necessary to re-sample. Advice may be sought from the Community Medicines Management. Recurrence of diarrhoea is common occurring in approximately 20% of patients.

There is currently not enough robust evidence to support the use of probiotic therapy.

While patients are symptomatic with diarrhoea normal aperient medication should be stopped.

PREVENTION OF SPREAD

Clostridium difficile spores from diarrhoeal patients will contaminate the environment, equipment and the hands/clothing of healthcare workers. Therefore:-

- Hand Hygiene is essential after all contact with the patient and their environment. **Spores are not reliably killed by alcohol gel therefore hands must be washed with soap and water.**
- Aprons should be worn as appropriate to protect clothing. This would include handling of patient's body fluids and contact with the environment.
- Gloves should be worn for contact with patient's body fluids or contaminated equipment. Hands must always be washed after removal of gloves.
- Medical/Nursing equipment having contact with the patient/environment should ideally be dedicated for that person. Chlorine releasing disinfectants (1,000 parts per million) are recommended for decontamination. (See Decontamination of Medical Equipment). Equipment returned to the Community Loan Store should be decontaminated according to the Loan Store instructions and the accompanying return form should state this is done.
- Where patients with *Clostridium difficile* disease are cared for in a care home setting, primary care staff should satisfy themselves that staff are familiar with processes designed to prevent cross-infection / contamination. This will include environmental cleaning regimes, isolation requirements and the need for dedicated toilet facilities whilst the patient is symptomatic. Home staff should be reminded of the importance of hand hygiene with liquid soap and water.
- If a patient with known or suspected *Clostridium difficile* disease is transferred to another care setting it is essential that the receiving facility is informed of the diagnosis in advance of transfer. Ambulance transport staff must also be informed when the transport is arranged.
- Carers of patients at home with *Clostridium difficile* infection should be taught to wash their hands with liquid soap and water after contact. The patient should also practice, with help as required, good hand hygiene. Enhanced environmental cleaning too high risk areas (toilets, door handles, light switches) with diluted household bleach
- continence waste should be discarded as hazardous waste in orange (or yellow) bags.

INVESTIGATION OF CASES

Cases of *Clostridium difficile* disease from all sources are reported through Public Health England Data Capture system (MESS) as required by DoH Mandatory Surveillance schemes.

Notes of patients diagnosed with *Clostridium difficile* disease are reviewed as part of Root Cause Analysis (RCA) investigation by acute NHS Trusts and local CCGs. This may involve requests to GPs for information and, on occasions to participate in the RCA process. This activity assists in understanding causation and thus developing programmes to reduce the incidence of this disease. Further review may be required if a patient dies with *Clostridium difficile* certified as a primary cause of death.

REVIEW

This Guideline will be reviewed two yearly and as required, in response to new guidance or regulation.

THIS PAGE HAS BEEN LEFT
INTENTIONALLY BLANK

MANAGEMENT OF OCCUPATIONAL EXPOSURE TO BLOOD-BORNE VIRUSES

INTRODUCTION

Due to the need for prompt action following an exposure to blood or blood-stained body fluids, staff must be aware of the action to be taken. Training in relation to the management of a needle-stick injury or blood splash must be provided as a mandatory component of Health and Safety / Infection Control induction training and annual training updates.

This policy section deals with sharps/splash incidents which may result in occupational exposure to Blood Borne Viruses (BBVs).

Blood-borne viruses (BBVs)

Blood-borne viruses include Hepatitis B and C and Human Immunodeficiency Virus (HIV)

All individuals infected with blood borne viruses may be capable of transmitting the virus to others irrespective of whether they are ill or apparently fit and healthy. Infectivity depends on a number of individual risk factors and will vary from individual to individual. Many individuals are unaware that they are infected and thus health care workers should always treat all blood and body substances as if they were infected. Body substances that have been shown to transmit BBVs include:

- cerebrospinal fluid
- peritoneal fluid
- pericardial fluid
- pleural fluid
- synovial fluid
- amniotic fluid
- human breast milk
- semen
- vaginal secretions
- saliva in association with dentistry
- any other body substance containing visible blood, e.g. faeces, urine, sputum
- unfixed tissues and organs
- exudate or other tissue fluid from burns or large skin lesions.

Prevalence of BBVs

The risk to the healthcare worker for each virus is proportional to the prevalence of that infection in the population, the infectious status of the individual source (which may or may not be known) and the risk of a significant occupational exposure occurring during the procedure being undertaken.

The risk of transmission to a healthcare worker from an infected service user following a sharps injury has been shown to be:

- Hepatitis B (e antigen positive) If healthcare worker is non-immune
1:3 Hepatitis C 1:30
- HIV 1:300

(UK Health Departments, 1998)

Certain geographical areas of the world have a higher prevalence of blood-borne viruses than others. Such information is useful in certain situations e.g. when making epidemiological assessments of risk. However, on a day-to-day basis, ethnicity is not used as a determinant of risk.

Transmission of BBVs

BBVs can be transmitted via:

- needle-stick injury with contaminated sharp object
- bite, scratch or other skin puncture with contaminated blood or bloodstained body fluids
- exposure of non-intact skin / mucous membranes to blood / body fluids
- unprotected sexual intercourse with an infected person
- infected mother to baby either via the placenta or at the time of delivery, or through breast-feeding
- exposure prone procedures (when infected health care workers can infect service users)
- sharing contaminated sharps/"works" of injecting drug abusers
- contaminated blood or blood products (not usually a risk in the UK but may occur if receiving blood in other countries)

Occupational acquisition of BBVs

A number of factors are associated with an increased risk of occupationally acquired BBV infection:

- deep injury
- visible blood on the device which caused the injury
- injury with a needle which had been placed in an artery or vein
- high levels of circulating virus in the source – as in late stage AIDS or during sero-conversion in the early stages of infection

These factors will be taken into consideration when assessing the risk of BBV transmission following a sharps injury. Such an assessment will usually be undertaken by either the local Occupational Health provider or local A& E / minor injury unit.

The risk of HIV transmission after percutaneous exposures involving larger volumes of blood, particularly if the source viral load is likely to be high, may exceed the average risk. This may occur if injury is sustained with a large hollow-bore needle when the needle contains a large volume of blood from either an artery or vein.

Risk of infection from cutaneous exposure from infected blood / or contaminated body fluids will depend on the infectivity of the material and the size of the exposed area e.g. people with large areas of psoriasis or eczema could be at higher risk of acquiring these infections if in contact with infectious material when splashed.

The highest risk of contamination from cutaneous exposure relates to splashes involving mucous membranes such as conjunctivae and mouth. Hence the requirement for staff to wear appropriate PPE when undertaking splash-inducing procedures

Sharps/Splash Incidents

There are three types of exposures in health care settings associated with significant risk. These are:

- percutaneous injury (from used needles, scalpel blades, lancets and other pointed instruments or equipment; bone fragments, significant bites which break the skin, etc)
- exposure of broken skin (abrasions, cuts, eczema, etc) to blood and/or blood stained body fluids
- exposure of mucous membranes, including the eyes, nose and mouth, to splashes of blood and/or blood stained body fluids

Management arrangements following occupational exposure to BBVs

It is essential that a risk assessment is undertaken at the earliest possible opportunity as delay in receiving prophylaxis (if required) could affect outcome i.e. the possibility of sero-conversion. This needs to be undertaken at the time of the injury NOT at the end of the shift. Current guidance states that HIV prophylaxis should be commenced within one hour of the incident, but can still be given after that time (up to 72 hours post-injury). Risk assessment should be carried out by a qualified and competent health care professional. This is usually either an occupational health professional or staff at local A/E ED or Minor Injury Unit. Local GPs may be assumed to provide this service (as health care professionals) but the registered provider must confirm that this is the case and document that arrangement (see next paragraph).

All registered care providers must have a comprehensive policy in place that details the precise process for staff to follow when sustaining a sharps injury / significant splash with potentially contaminated blood or blood-stained body fluids. The policy should clearly state how staff can access prompt professional risk assessment and treatment. All organisations must have either 24-hour access to an Occupational Health Service and / or a Service Level Agreement in place with a local NHS Trust if local provision of risk assessment and

treatment is not available or is only available during working hours i.e. if no out-of-hours occupational health service is available.

First aid:

- Encourage bleeding from the wound. Do not suck
- Wash the area thoroughly with warm running water and soap
- Cover with water-proof dressing
- Eyes or mouth - irrigate with copious amounts of saline or water

Report

ALL sharps injuries and splash incidents must be reported to the senior nurse or manager on duty (dependent on place of work) as soon as possible, but do not delay seeking guidance on the need for prophylaxis if a manager cannot be contacted.

If the affected staff member has access to local Occupational Health services then contact should be made soon after injury. Alternatively, the local emergency department can be contacted by phone. If telephone support is not immediately available then the injured staff member should attend emergency department at the earliest opportunity for risk assessment.

Record details

Complete an accident and incident report form which must be provided by the registered provider

If the exposure is from a Hepatitis B, C or HIV positive source, RIDDOR form 2508 will be completed by the Occupational Health Physician once confirmation of the test results are known

Try to identify the source service user. This is not essential but can assist in assessing risk.

Complete the Checklist Form - following sharps/splash incident (Sample in Appendix C) to help with the risk assessment and take it to the nearest emergency department. / Occupational Health.

The Checklist will help to establish if the member of staff has had a significant exposure to a high risk body fluid and provides guidance on the important questions that will be asked by the assessing clinician when undertaking risk assessment.

Attending for risk assessment / treatment

When attending for risk assessment, the staff member affected should take the completed Checklist to ensure that appropriate information is available to the clinician undertaking the

risk assessment. The staff member may be required to have a blood sample taken and stored for further testing if necessary. *This blood is not routinely tested* but is stored for future testing should the HCW demonstrate possible evidence of infection. Blood will not be tested without the individual HCWs permission. Results of staff testing must be sent to either Occupational Health or the individual's GP. He/she may also be required – dependent on the risk assessment – to have medication or immunisation to reduce the likelihood of sero- conversion. Any concerns due to exposure, drug treatment or employment etc. can be discussed in confidence at this time

If the source is identified the clinician undertaking the risk assessment will arrange for testing of that service user's blood via the service user's GP/clinician. No blood sample should ever be taken by the staff member or care provider.

The Occupational Health Department or emergency department will arrange the co-ordination of results and follow-up and determine whether further blood tests will be required at 3, 6 and 12 months. This will be undertaken the following working day after injury.

Post-exposure Prophylaxis (PEP)

Hepatitis B – vaccination/prophylaxis

All health care workers at risk of exposure to blood/body fluids as part of their work must be offered vaccination against Hepatitis B at the commencement of employment.

A primary course consists of 3 injections at 0, 1 and 6 month intervals followed by a blood test to determine antibody levels. Some people may not develop antibodies even after further doses of vaccination.

Following a significant exposure, Hepatitis B specific immunoglobulin may also be required within 24 hours of injury to prevent acquisition.

For staff that sustain an injury/exposure and have *not* received a primary course of HBV immunisation then an accelerated course of immunisation may be recommended. This consists of injections at 0, 1 and 2 month intervals.

HIV post exposure prophylaxis (PEP)

Although there is no protective vaccine for exposure to HIV there are certain drugs which, if taken soon after exposure, offer some protection to the exposed individual. Ideally, this should be received within 1 – 2 hours of injury but can still be administered for up to 72 hours post-injury. Healthcare organisations that provide PEP following injury e.g. acute NHS trusts are required to have in place a comprehensive PEP policy for the administration of medication and for the initial assessment and pre-and post-test counselling of injured staff /service users.

Hepatitis C (HCV) prophylaxis

There is currently no vaccine available for the prevention of Hepatitis C infection. Specialist advice and management would be made available to staff at risk of HCV acquisition following exposure

Bloodborne virus-infected healthcare workers

Healthcare workers who are known to have a BBV are restricted from certain aspects of service user care. These restrictions are in place in order to reduce the risk of transmission of BBVs to service users from infected healthcare workers whilst carrying out certain procedures known as Exposure Prone Procedures (EPP's).

EPP's are those invasive procedures where there is a risk that injury to the worker may result in the exposure of the service user's open tissues to the blood of the worker (bleed-back). These include procedures where the worker's gloved hand may be in contact with sharp instruments, needle tips or sharp tissues (e.g. shards of bone or teeth) inside a service user's open body cavity, wound or confined anatomical space. However, other situations, such as pre-hospital trauma care and care of service users where the risk of biting is predictable (e.g. such as with a disturbed and/or violent service user) should be avoided by healthcare workers restricted from performing EPP's.

Staff are under legal, professional and ethical duties to protect the health and safety of their service users.

Any member of staff who knows themselves to be infected (or at risk of infection) with a BBV must seek guidance for advice and management of risk reduction.

Appendix C

CHECKLIST FOLLOWING SHARPS/SPLASH INJURY

To be completed by staff member who has sustained the sharps/splash injury and then taken to Accident & Emergency Department and/or Occupational Health.

Personal details	
Name:	Date of birth:
Post:	Place of work:
Telephone number:	Manager:
Home:	
Work:	
Date:	Time of accident/incident:

Details of the injury
<p>Brief description of the incident with blood/blood stained body fluids (please tick box if applicable)</p> <p>Sharps injury: Needle/scalpel blade or other sharp instrument Scratch Bite Cut Bone Other</p> <p>Skin exposure: Abrasion Eczema Psoriasis Other</p> <p>Exposure to mucous membrane Eye Other</p> <p>Which high risk body substance?</p> <p>Blood Blood stained body fluid Vaginal secretions Saliva (if visibly blood stained e.g. in association with dentistry)</p> <p>Other please specify</p>

Local arrangements for risk assessment / management of injury
<p>Occupational Health:</p> <p>During surgery hours please contact:</p>
<p>Accident and Emergency Department:</p> <p>Out of surgery hours please contact:</p> <p>Identify yourself as a Healthcare Worker who has sustained a sharps injury.</p>

MANAGEMENT OF INFECTIONS IN STAFF

INTRODUCTION

From time to time, health care staff may develop infections which could expose some service users and colleagues to the risk of infection.

Symptoms or signs of infection can appear trivial to staff who are usually fit and well, but can cause severe problems in vulnerable service users.

REPORTING

Early reporting and implementation of suitable control measures can prevent cross-infection and subsequent outbreaks of infection.

Confirmed or suspected transmissible infections in health care staff should be reported by the staff member to the Practice Manager or lead clinician. In addition, advice can be sought from the local Infection Control Advisor / HPU / Consultant Medical Microbiologist if there is concern regarding spread to other staff and/or service users.

TREATMENT

If necessary, treatment should only be undertaken by the Occupational Health provider (OH) or the individual's General Practitioner (GP), as appropriate.

EXCLUSION FROM WORK

The necessity for exclusion from work should be discussed with the lead clinician and in liaison with the Public Health England / Consultant Medical Microbiologist / Environmental Health Officer (EHO) as necessary.

Staff with gastro-intestinal infections who handle or prepare food in the course of their work may be required to stay off work until their stool specimens are free of micro-organisms. Guidance must be sought from Occupational Health or the individual's GP who will make the decision regarding return to work after liaising with a medical microbiologist/CCDC where necessary.

Although not an exhaustive list, the following table summarises the risks to service users from staff with some infectious diseases.

INFECTIOUS DISEASES AND ADVICE TO STAFF

INFECTION	SERVICE USER RISKS	ADVICE TO STAFF
BLOOD BORNE VIRUSES (BBV) including Hepatitis B Hepatitis C HIV	The risk of transmission of a blood borne virus from a HCW to a service user is extremely low. Not all staff will be aware of their possible infectious status therefore standard infection control practice should be applied at all times.	Staff should seek confidential advice from their GP or local clinician as soon as possible following diagnosis, or if concerned that they may have been exposed to a BBV. An assessment will be made regarding further clinical management, in consultation with the HPU. If a staff member is diagnosed with a BBV some modification of working practices may be necessary in some situations.
INFECTED SKIN LESIONS or skin conditions, i.e. psoriasis, eczema, impetigo etc.	A bacterial infection is the usual cause which can then be spread to service users. Particularly vulnerable service users are those with open lesions, surgical or traumatic wounds, the immuno-compromised or elderly.	Staff suffering with these infections may be required to remain off duty until the infection has resolved unless it can be covered by an occlusive dressing. Antibiotics are often required.
CHICKEN POX (varicella)	Non-immune and immune-suppressed service users may require active protection e.g. immunisation and guidance should be sought from the service users GP immediately exposure is confirmed or suspected.	Non-immune health care staff, i.e. those who have not had the disease or vaccination, should seek immediate medical advice and may be medically suspended from clinical work from day 8-21 post- exposure. Non-immune pregnant staff (particularly < 20 weeks pregnant or in last 3 weeks of pregnancy) must discuss with their Obstetrician urgently. Immune-suppressed staff who have had contact with an infectious case must discuss their exposure with their clinician and / or Occupational Health provider immediately. Immunisation against varicella (chickenpox) is now widely available for non-immune individuals. See section – Vaccination Programme for Staff
COLD SORES and GENITAL HERPES INFECTIONS	Caused by the herpes simplex virus, which may expose some service users who are immuno- compromised, neonates and pregnant women to particular risks. Viral encephalitis may ensue in these susceptible service users.	Depending on working environment staff may need to remain off duty until resolution of symptoms and lesions are dry. Seek medical guidance. Do not touch lesions, wash hands thoroughly.

INFECTION	SERVICE USER RISKS	ADVICE TO STAFF
DIARRHOEA and/or VOMITING	These may be symptoms of food poisoning or viral infection, which can result in cross infection causing outbreaks. Viral outbreaks spread rapidly & vulnerable service users are at particular risk especially babies and the elderly.	Staff must remain off duty until 48 hours after resolution of the symptoms. Notify the Practice Manager / lead clinician if more than 2 staff affected.
INFLUENZA	A viral infection which usually spreads to service users and other staff if prompt action is not taken. It can cause high morbidity and mortality rates, particularly in the elderly.	Staff should remain off duty until resolution of symptoms. Uptake of influenza vaccine is recommended for both care workers and vulnerable service users.
MEASLES, MUMPS and RUBELLA	Cases are highly infectious.	Non-immune staff must inform Practice Manager / lead clinician of exposure to an infectious source. Non-immune pregnant staff, i.e. those who have no history of disease and/or no positive antibody test must seek medical guidance especially in the first trimester of pregnancy.
SCABIES	Staff may be infected by skin to skin contact with service users. Scabies is often difficult to diagnose in the elderly. Service users remain contagious until 24hrs post-treatment. If > 1 service user affected, treatment will need to be undertaken simultaneously.	Staff contacts of infested service users may require treatment but this is unlikely to occur in General Practice. If staff member is affected, family contacts will also require treatment. Contact IC/HPU for further guidance.
SORE THROATS	These may have many causes but are usually viral. Bacterial causes e.g. streptococcal infections can cause severe infections in vulnerable service users. simultaneously.	Staff should remain off duty until resolution of symptoms, if unwell and with a severe sore throat associated with pyrexia. Notify the Practice Manager / lead clinician if more than one member of staff is affected.

TUBERCULOSIS	Physical isolation is only required for those who are pulmonary smear positive for AFBs (acid fast bacilli). Isolation should continue until at least 14 days after commencing appropriate anti- tuberculosis therapy and/or until advised by TB specialist/team.	The necessity for exclusion of diagnosed staff members from work will require discussion by the lead clinician in conjunction with the TB specialist team. Contacts will be investigated by the TB nurse specialist and HPU
PARVOVIRUS (FIFTH DISEASE)	Mild, non-febrile viral disease characterized by erythema of cheeks. Most infectious prior to development of rash but not infectious thereafter.	Can cause foetal abnormality. Pregnant staff less than 20 weeks pregnant should seek advice from their obstetrician.

VACCINATION PROGRAMME FOR (1) STAFF AND (2) SERVICE USERS

(1) STAFF

INTRODUCTION

Under the Health and Safety at Work Act (1974) employers, employees and the self-employed have specific duties to protect, so far as reasonably practicable, those at work and others who may be affected by their work activity, such as contractors, visitors and service users. Central to health and safety legislation is the need for employers to assess the risks to staff and others.

The Control of Substances Hazardous to Health (COSHH) Regulations 2002 require employers to assess the risks from exposure to hazardous substances, including micro-organisms (called biological agents in COSHH) and to bring into effect the measures necessary to protect workers and others from those risks as far as is reasonably practicable.

A local Occupational Health policy for immunisation and vaccination of health care staff should be available *and is a requirement of the Code of Practice (criterion 10)*. This should be supported by employment health assessment which should include a review of immunisation needs. Employers need to be able to demonstrate that an effective employee immunisation programme is in place, and they have an obligation to arrange and pay for this service. It is recommended that immunisation programmes are managed by occupational health services with appropriately qualified specialists.

This section draws on published guidance available from the Department of Health *Immunization against infectious diseases* (the Green Book) Chapter 12: Immunisation of healthcare and laboratory staff (2013) together with relevant chapters relating to individual infections.

IMPORTANCE OF IMMUNISATION

Any vaccine-preventable disease that is transmissible from person to person poses a risk to both healthcare workers (HCWs) and their service users. HCWs have a duty of care towards their patients which includes taking reasonable precautions to protect them from communicable diseases. Immunisation of HCWs may therefore:

- Protect the individual and their family from an occupationally-acquired infection
- Protect service users including those that are vulnerable and who may not respond well to their own immunisation
- Protect other healthcare staff
- Allow for the efficient running of services without disruption.

RECOMMENDATIONS BY STAFF GROUPS

The objective of occupational immunisation is to protect workers at high risk of exposure and their families; to protect patients and other staff from exposure to infected workers and to sustain the workforce. Potential exposure to pathogens, and therefore the type of immunisation required, may vary from workplace to workplace.

Staff working in primary care may be exposed to unknown pathogens on a regular basis.

STAFF INVOLVED IN DIRECT PATIENT CARE

This includes staff who have regular clinical contact with patients and who are directly involved in patient care. This includes doctors, nurses, support workers / healthcare assistants, paramedics, professionals allied to medicine and volunteers who work with patients.

ROUTINE VACCINATION

All staff should be up to date with their routine immunisations e.g. tetanus, diphtheria, polio and MMR (measles, mumps and rubella). The MMR vaccine is especially important in the context of the ability of staff to transmit measles or rubella infections to vulnerable groups. Satisfactory evidence of protection would include documentation of having received two doses of MMR or having had positive antibody tests for measles and rubella.

SELECTED VACCINES

HEPATITIS B

Department of Health guidance on Hepatitis B immunisation must be followed and incorporated into a local Occupational health policy for staff immunisation.

Hepatitis B vaccination is recommended for healthcare workers who may have direct contact with patients' blood or blood-stained body fluids. This includes any staff that are at risk of injury from blood-contaminated sharp instruments or of being deliberately injured or bitten by patients.

The primary course (of HBV immunisation) consists of three injections given over six months. Antibody titres for hepatitis B should be checked one to four months after the completion of a primary course of vaccine. Such information allows appropriate decisions to be made concerning post-exposure prophylaxis following known or suspected exposure to the virus.

Alternatively staff may be required to show satisfactory evidence of immunity.

N.B. staff undertaking Exposure Prone Procedures (EPP's) will be required to provide an

Identity Validated Sample (IVS) from a UK laboratory demonstrating serological evidence of immunity to Blood Borne Viruses (BBV's) including HBV,HCV and, in some cases, HIV

In the event of a sharps injury or significant exposure to blood or blood- contaminated body fluids e.g. conjunctival splash, the degree of risk will be assessed and appropriate prophylaxis will be provided. It is essential that all accidental exposures are reported as soon as possible, as in certain circumstances it is necessary to give Hepatitis B immunoglobulin and/or a booster dose of vaccine, even though the individual has been immunised, to minimise the risk of acquiring infection. Refer to section – Management of Occupational Exposure to BBV's.

BCG (TUBERCULOSIS)

The major source of tuberculosis is from individuals who have active pulmonary TB and are AFB (acid fast bacilli) smear positive and during the first 2 weeks of compliant treatment. It is uncommon for health care staff in good health to acquire tuberculosis from patients. However, all staff in regular contact with patients/clients who are / may be infectious and those who handle material which may contain tubercle bacilli i.e. sputum specimens, are at risk.

All relevant staff should be screened soon after taking up employment. A history will be taken regarding past BCG vaccination and/or any previous Tuberculin /Mantoux test results and any recent chest x-rays. BCG immunisation is not given more than once due to the risk of adverse reactions. There is also no data on protection afforded to individuals older than 35 years receiving BCG for the first time and thus it is not routinely given. Clinical risk assessment of individual staff cases should be undertaken if considered appropriate.

INFLUENZA

Offering influenza vaccine to health care staff is recognised as a cost-effective way of ensuring that key clinical contact staff remain influenza free during the Influenza season and reduces the likelihood of transmission of influenza to vulnerable patients (by staff).

Influenza vaccination is therefore recommended for healthcare workers directly involved in patient care, who should be offered influenza immunisation on an annual basis.

See the Influenza Immunisation Guidelines (Dept. of Health website) for guidance on the annual patient protection programme.

CHICKENPOX (VARICELLA)

Chickenpox is highly infectious and varicella vaccine is recommended for susceptible healthcare workers who have direct patient contact.

A history should be taken regarding previous history of chickenpox or herpes zoster

(shingles) acquisition, documentary evidence of vaccination or documentary evidence of immunity to chickenpox. Healthcare workers with a negative or uncertain history of chickenpox or herpes zoster should be serologically tested. Relevant staff found to be non-immune may be offered vaccination unless there are contra-indications.

NON-CLINICAL STAFF

This includes ancillary staff that may have social contact with patients but are not involved in direct patient care. This group includes receptionists, clerks, porters and housekeepers / cleaners.

ROUTINE VACCINATION

Exactly the same requirements as for clinical staff (above)

SELECTED VACCINES

BCG (Tb) – not required

HEPATITIS B – required if exposed to similar risks as clinical staff (as above)

VARICELLA – required if susceptible (as above) and with regular patient contact

INFLUENZA – not routinely recommended (unless directed by Department of Health) **HEPATITIS A**

Certain groups of staff may be at risk of acquiring Hepatitis A infection, which is spread via the faecal-oral route. This includes staff working in mental health and learning disabilities environments together with maintenance workers involved in procedures likely to involve contact with raw sewage, such as drain cleaning/unblocking and laundry staff handling contaminated laundry.

ADDITIONAL CONSIDERATIONS

Staff who are travelling abroad and are unsure of the vaccination requirements for the country that they are visiting should contact their GP or the Foreign Office website for advice. Travel clinics are also an excellent source of up to date information. Staff have a responsibility to the employing organisation and its clients to ensure that they are appropriately immunised prior to and during travel. If, following return from overseas travel, staff are unwell they should seek a medical opinion at the earliest opportunity both to ensure their appropriate treatment and to minimise the risk of disease transmission to their work colleagues and clients. They should also report their illness to the lead clinician for an assessment of their fitness to work.

HEALTH CLEARANCE AND IMMUNISATION FOR AGENCY, LOCUM AND VISITING STAFF

All agency, locum, visiting and voluntary staff working for the organisation must comply with the requirements for vaccination cover in the same manner as that of permanent

staff members. This is a requirement under Regulation R4 of the Management Regulations (1999) and the associated Approved Code of Practice (ACOP). This requirement states that it is the responsibility of the employer to specify to employment agencies the minimum requirements for temporary workers under Regulation 12.1 and agencies have a duty to comply.

Agencies are responsible for ensuring that their employees meet these requirements.

FURTHER ADVICE

All staff seeking guidance on vaccination requirements should contact Occupational Health in the first instance.

(2) SERVICE USERS

Registered providers are expected to provide assurance that local policies and procedures are in place in relation to the immunisation status of service users. Dependent on age and risk, service users may require a range of immunisations to protect them. These will range from childhood immunisations to the annual requirement for influenza vaccination of the elderly and others at risk. The decision regarding immunisation is the responsibility of the service users' Registered Medical Practitioner and / or clinicians involved in the health care management of individual service users as well as local Public Health departments, Commissioners and Immunization Co-coordinators.. Decisions will be made on the basis of guidance published by the Department of Health (and regularly updated) entitled *Immunisation against infectious disease* ("The Green Book"). All service users' immunisations must be documented.

THIS PAGE HAS BEEN LEFT
INTENTIONALLY BLANK

DEFINITIONS OF TERMINOLOGY

Asepsis-is recognised as the state of being free from pathogenic microorganisms

Aseptic technique- is defined as a means of preventing or minimising the risk of introducing harmful micro-organisms into sterile/key sites of the body when undertaking clinical procedures

Aseptic Non-Touch Technique (ANTT) is a specific type of aseptic technique. The overriding principle is that key sites e.g. wound, must not come into contact with any item (hand, equipment, solution) that is not sterile. Sterile gloves are not always required for standard ANTT. Each procedure must be risk assessed. Whether sterile or non-sterile gloves are worn depends if you can avoid touching the sterile parts of equipment which will come into contact with the service users susceptible areas e.g. their wound

Clostridium Difficile Infection (CDI) –Anaerobic, gram positive spore forming bacillus. These spores are resistant to exposure to air, drying, heat and survive in the environment. Following antibiotic therapy the intestinal flora is altered which allows any C diff bacteria to proliferate. The bacteria produce 2 toxins:

1. Toxin A which irritates the colon and causes what is commonly known as antibiotic associated diarrhoea
2. Toxin B which is predominately cytotoxic

Cross infection –The transmission of disease from one person to another because of a breach in barrier.

Decontamination-The process used to remove organic matter and micro-organisms from an item and render it safe for use. There are three levels of decontamination: cleaning, disinfection and sterilisation.

Hand decontamination-The physical removal of blood, body fluids and transient micro-organisms from the hands e.g. hand washing.

Hand Hygiene-A general term that applies to either: hand washing, antiseptic handwash, antiseptic hand rub or surgical antisepsis

Healthcare Associate Infection (HCAI) - Infections that occur as a result of contact with the healthcare system in its widest sense-in community and hospital settings

Healthcare Worker- Any person employed by the health service, social services authority or agency to provide care for sick, disabled or elderly people.

Infection Prevention and Control (IP&C) –To reduce to an acceptable minimum the risk of the acquisition of an infection amongst service users, healthcare workers and any others in the healthcare environment.

Infection Prevention and Control Team (IP&CT) - A specialist team employed by the organisation to provide expert reactive and proactive information and advice about the management of healthcare associated infections and incidents.

Methicillin Resistant Staphylococcus Aureus (MRSA) - is a variant of Staphylococcus aureus which has developed resistance to commonly used antibiotics and is considered endemic in both hospitals and the community setting and may be more difficult to treat because of limited treatment options.

Personal Protective Equipment (PPE) - All equipment which is intended to be worn or held by a person to protect them from risks to health and safety whilst at work. Examples of PPE include gloves, aprons and eye and face protection.

Public Health England (PHE) - Is an executive agency of the Department of Health and Social Care in the UK which exists to protect and improve the nation's health and wellbeing and reduce health inequalities

Resident Organisms - Micro-organisms that colonise the deeper crevices of the skin and hair follicles as they have adapted to the hostile environment. They are not readily transferred to other people or objects and are not easily removed by the mechanical actions of soap and water, but can be reduced in number with the use of an antiseptic solution.

Risk Assessment – Making a suitable and sufficient assessment of risks-this will involve identifying the hazards (something with the potential to do harm) and evaluating the extent of risks (the likelihood that the harm from a particular hazard is realised) and identifying measures needed to comply with legal requirements.

Sharps injury/ Incident - When intact skin is breached by a sharp object (needle instrument, or bone) This also includes human bites and scratches that break the skin.

Transient Organisms - Micro-organisms acquired on the skin through contact with surfaces. The hostile environment of the skin means that they can usually only survive for a short time, but they are readily transferred to other surfaces touched. They can be removed by washing with soap and water or inactivated by alcohol hand rub

BIBLIOGRAPHY

INFECTION PREVENTION & CONTROL MANAGEMENT

Health Protection Legislation (England) 2010 Guidance London DH Health & Social Care Act 2008 Regulated Activities Regulations 2010

Department of Health (2010) Code of Practice for health & adult social care on the prevention and control of infections and associated guidance

Department of Health (2006) Essential Steps to safe, clean care: reducing healthcare-associated infections London: DH

Department of Health (2004) Towards cleaner hospitals and lower rates of infection: a summary of action London: DH

Department of Health (2003) Winning ways: working together to reduce healthcare associated infection in England. Report from the Chief Medical Officer. London: DH

Department of Health (2002) Getting ahead of the curve: a strategy for combating infectious diseases (including other aspects of health protection). A report by the Chief Medical Officer London: DH

Department of Health (1995) HSG (95) 10: Hospital infection control. London: DH

NICE CG139; 2012 Infection Control

Antimicrobial prescribing

Department of Health (2007) Saving Lives: reducing infection, delivering clean and safe care: antimicrobial prescribing London DH

Department of Health, Standing Medical Advisory Committee, Sub-Group on Antimicrobial Resistance (1998) The path of least resistance. London: DH.

Audit

Infection Prevention Society (2016) Audit tools for monitoring infection control guidelines within the community setting. Infection Prevention Society

Care of deceased persons

Health and Safety Executive (2003) Safe working and the prevention of infection in the mortuary and post-mortem room London: HSE

Clinical practice and patient management

Department of Health (2003) Winning ways: working together to reduce healthcare associated infection in England. Report from the Chief Medical Officer. London: DH

Department of Health (2007) Essential Steps to safe, clean care: reducing HCAs in care homes etc. London DH

National Institute for Clinical Excellence NICE (2012) Infection Control: Prevention of healthcare associated infections in primary and community care (Partial update of NICE Clinical Guideline 2: 2003) London NICE

National Patient Safety Agency (2008) Patient Safety alert: clean hands save lives. London: NPSA

Pratt RJ, Pellowe CM, Wilson JA, Loveday HP et al (2007) epic2: National evidence-based guidelines for preventing healthcare associated infections in NHS hospitals in England. Journal of Hospital Infection 65 (Supplement)

World Health Organisation WHO (2009) Guidelines on hand hygiene in health care Geneva WHO

Control of infections associated with specific alert organisms

Acinetobacter and other antibiotic – resistant bacteria

Health Protection Agency (2006) Working party guidance on the control of multi-resistant acinetobacter outbreaks London: HPA

Clostridium difficile

Department of Health / Health Protection Agency (2009) *Clostridium difficile* infection: how to deal with the problem London: DH

Diarrhoeal infections

Health Protection Agency (2004) Preventing person-to-person spread following gastrointestinal infections: guidelines for public health physicians and environmental health officers, Communicable Disease and Public Health 7(4): 362-84

Health Protection Agency (2010) A report on the management of diarrhoea in care homes London HPA

Public Health England (2012) Guidelines for the management of norovirus outbreaks in acute and community health and social care settings

Meticillin-resistant Staphylococcus Aureus (MRSA)

Coia JE, Duckworth GJ, Edwards DI, Farrington M, et al, for the joint working party of the British Society for Antimicrobial Chemotherapy, the Hospital Infection Society, and the Infection Control Nurses Association (2006), Guidelines for the control and prevention of meticillin-resistant Staphylococcus aureus (MRSA) in healthcare facilities Journal of Hospital Infection (Supplement): 63 S1-44

MRSA screening

Department of Health (2008) MRSA screening – operational guidance London: DH
Department of Health (2008) MRSA screening – operational guidance 2 London: DH

Department of Health (2006) Screening for MRSA colonisation: a summary of best practice, Professional letter from the Chief Medical Officer and the Chief Nursing Officer. London: DH

Panton-Valentine leukocidin (PVL) associated and community associated

Staphylococcus aureus

Health Protection Agency (2008) Guidance on the diagnosis and management of PVL-associated Staphylococcus aureus infection (2nd edition). London: HPA

Respiratory Infections

Health Protection Agency (2012) Respiratory Infections Guidance HPA

National Institute for Health and Clinical Excellence (2008) Respiratory tract infections-antibiotic prescribing NICE Clinical Guideline 69 London: NICE

Department of Health (2006) Immunisation against infectious disease ('The Green Book') London: DH

Steer J A et al (2012) Guidelines for the prevention and control of Group A streptococcal infection in acute healthcare and maternity settings in the UK Journal of Hospital Infection 64 1-18

Creutzfeldt-Jakob disease (CJD) and other human prion diseases

Advisory Committee on Dangerous Pathogens TSE Working Group guidance Transmissible spongiform encephalopathy agents: safe working and the prevention of infection – guidance from the Advisory Committee on Dangerous Pathogens.

London: DH

Tuberculosis

National Institute for Health and Clinical Excellence (2006) Tuberculosis: clinical diagnosis and management of tuberculosis, and measures for its prevention and control. NICE Clinical Guideline 33 World Health Organisation (2009) WHO Policy on TB infection control in healthcare facilities, congregate settings and households WHO

Viral haemorrhagic fevers

Advisory Committee on Dangerous Pathogens (1996) Management and control of viral haemorrhagic fevers ACDP

Decontamination of reusable medical devices

Department of Health (2007) Clarification and policy summary – decontamination of reusable medical devices in the primary, secondary and tertiary care sectors London: DH

Department of Health (2007) HTM 01-01: Decontamination of reusable medical devices: Part A – Management and environment; Part B – Additional management guidance and common elements; Part C – Sterilizers; Part D – Washer-disinfectors and ultrasonic cleaners London: DH

Department of Health (2007) Clarification and Policy Summary – Decontamination of re-usable medical devices in the primary, secondary and tertiary care sectors DOH

Medical Devices Regulations 2002

Medical Devices (Amendment) Regulations 2008

Medicines and Healthcare products Regulatory Agency (2006) DB 2006 (05): Managing Medical Devices. London: MHRA

Medicines and Healthcare products Regulatory Agency (2006) Sterilization, disinfection and cleaning of medical equipment: Guidance on decontamination from the Microbiology Advisory Committee to department of Health. London: NHRA

Education of Care workers

NHS Core Learning Unit (2005) Infection control e-learning programme for healthcare and social care staff. Available from: www.infectioncontrol.nhs.uk

Skills for Care (2005) Common Induction Standards Social Care (Adults, England) Leeds: Skills for care

Skills for Care (2005) Knowledge set for infection prevention and control Social Care (Adults, England). Leeds: Skills for care

Environment (Estate)

Department of Health (2011) Facilities for primary and community care services: planning and design manual DOH

Department of Health (2011) Facilities for primary and community care services: policy and service context manual DOH

Health Building Note (HTM) 00-10 Performance Requirements for building elements used in healthcare facilities 8941:0.6 England DOH

NHS Estates (2002) HFN 30: Infection control in the built environment: design and planning

Environmental Cleaning

National Patient Safety Agency (2009) Revised healthcare cleaning manual. London: NPSA

National Patient Safety Agency (2007) National specifications for cleanliness in the NHS: a framework for setting and measuring performance outcomes. London: NPSA

National Patient Safety Agency (2007) safer practice notice 15: Colour coding hospital cleaning materials and equipment NPSA

National Patient Safety Agency (2010) National Specifications for cleanliness in the NHS: guidance on setting and measuring performance outcomes in primary care medical and dental premises NPSA

Planned preventive maintenance

NHS Estates (2002) HEN 30: Infection control in the built environment: design and planning DOH

Healthcare waste

Controlled Waste Regulations England and Wales 2012

Department of Health (2012) HTM 07-01: Environment and sustainability: safe management of healthcare waste v1.8 DOH

Department of Health (2003) NHS Standard Service Level Specifications: Service specific specification-waste management

Health and Safety Executive (2009) Managing offensive / hygiene waste London: HSE

Pest control

Department of Health (2003) NHS Standards Service Level Specifications: Service specific specification – pest control version 2

Management of water supplies

Department of Health (2006) HTM 04-01: Water systems: the control of Legionella, hygiene, “safe” hot water, cold water and drinking water systems. Part A: Design, installation and testing and Part B: Operational management

Health and Safety Executive (2000) Legionnaires Disease Approved Code of Practice L8 3rd edition HSE

Food Hygiene and Safety

Food Hygiene (England) Regulations (2006) Statutory Instrument SI 2006/14

Department of Health (1996) HSG (96) 20: Management of food hygiene and food services in the NHS. London: DH

Health and safety

Health and Safety Executive (2008) Blood-borne viruses in the workplace, Guidance for employers and employees – INDG342 London: HSE.

Health and Safety Executive (2006) Five steps to risk assessment – INDG163 (rev2) London: HSE

Health and Safety Executive (2005) COSHH: a brief guide to the regulations: what you need to know about the Control of Substance Hazardous to Health Regulations 2002 (COSHH). London: HSE

Health and Safety Executive (2005) Respiratory protective equipment at work: a practical guide. HSG53. London: HSE

Health and Safety Executive (1999) Management of Health and Safety at Work Regulations: Management of health and safety at work. Approved code of practice and guidance Statutory Instrument No. 3242

Health and Safety Executive (1999) A guide to the reporting of injuries, diseases and dangerous occurrences regulations (RIDDOR) 1995. London: HSE

Health and Safety Executive (1992) Personal Protective Equipment at Work Regulations London: HSE

Health and Safety at Work etc Act 1974

Healthcare workers infected with a blood-borne virus

Department of Health (1993) HSG 93 (40): Protecting health care workers and patients from hepatitis B. London: DH

Department of Health (1993) Protecting health care workers and patients from hepatitis B: recommendations of the Advisory Group on Hepatitis. London: DH

Department of Health (1996) Addendum to HSG 93 (40): Protecting health care workers and patients from hepatitis B

Department of health (2000) HSC 2000/020: Hepatitis B infected health care workers. London: DH

Department of Health (2000) Hepatitis B infection health care workers. Guidance on implementation of Health Service Circular 2000/020 London: DH

Department of Health (2007) Hepatitis B infected healthcare workers and antiviral therapy. London: DH

Department of Health (2002) HSC 2002/010: Hepatitis C infected health care workers.

London: DH

Department of Health (2002) Hepatitis C infected health care workers. Guidance on implementation of Health Service Circular 2002/010 London: DH

Department of Health (2005) HIV-infected health care workers: guidance on management and patient notification. London: DH

Health and Safety Executive (2008) Blood-borne viruses in the workplace, Guidance for employers and employees – INDG342 London: HSE

Immunisation

Department of Health (2006) Immunisation against disease ('The Green Book') London: DH

Isolation of service users with an infection

Department Of Health (2008) Guide to best practice: isolation of patients. London: DH

Linen, laundry and dress (uniform)

Department of Health (2003) NHS Standard Service Level Specifications: Service specific specification – linen

Department of Health (1995) HSG (95) 18: Hospital laundry arrangements for used and infected linen. London: DH

Department of Health (2007) Uniforms and work wear: an evidence-based for developing local policy London DH

Department of Health (2010) Uniforms and work wear: guidance on uniform and work wear policies for NHS employers London DH

Management of occupational exposure to blood-borne viruses and post-exposure prophylaxis

Department of Health (2008) HIV post-exposure prophylaxis: guidance from the UK Chief Medical Officers' Expert Advisory Group on AIDS. 4th edition London: DH

Health Protection Agency (2005) Reporting of occupational exposure to blood borne viruses – history and how to report London: HPA

Medical devices directives / regulations

Medicines and Healthcare products Regulatory Agency (2006) Bulletin No 17: Medical devices and medicinal products. London: MHRA

Occupational health

NHS Employers (2007) The Healthy workplaces handbook

Department of Health (2007) Health clearance for tuberculosis, hepatitis B, hepatitis C and HIV: New healthcare workers. London: DH

Health and Safety Executive (1985) Reporting of injuries, Disease and dangerous Occurrences Regulations (RIDDOR)

Health and Safety Executive (2005) Control of substance hazardous to health (fifth edition): The Control of Substance Hazardous to Health Regulations 2002 (as amended): Approved Code of Practice and guidance. London: HSE

Outbreaks of communicable infection

Department of Health (2008) Guide to best practice: isolation of patients. London: DH

Department of Health (1995) HSG (95) 10: Hospital infection control. London: DH

Prevention of occupational exposure to blood-borne viruses, including the prevention of sharps injuries

Health Protection Agency (2008) Examples of good and bad practice to avoid sharps injuries London: HPA

Department of Health (1998) Guidance for clinical health care workers: protection against infection with blood-borne viruses. London: DH

Council Directive 2010/32/EU Framework Agreement on prevention from sharp injuries in the hospital and healthcare sector European Union

Advisory Committee on Dangerous Pathogens (1995) Guidance on protection against blood-borne infections in the work place: HIV and hepatitis. PL CO (95)5

Provision of information to the patient, the public and other service providers

National Patient Safety Agency (2005) being open – communicating patient safety incidents with patients and their carers London: NPSA

Single-use devices

Medicines and Healthcare products Regulatory Agency (2006) Single-use Medical Device: Implications and consequences of Reuse. Medicines and Healthcare products Regulatory Agency Device Bulletin DB 2006 904

USEFUL TEXTS

Lawrence J and May D Editors (2003) Infection Control in the community London Churchill Livingstone

Dougherty L and Lister S Editors (2015) The Royal Marsden Hospital Manual of Clinical Nursing Procedures 9th edition Chichester Wiley Blackwell

Chin J Editor (2000) Control of Communicable Diseases Manual 17th edition Washington APHA (American Public Health Association)